



Contribution ID: 125

Type: **Invited Presentation**

High-sensitivity and High-resolution Laser Spectroscopy of $^{76,77,78}\text{Cu}$ at CRIS

Thursday, 1 June 2017 14:15 (15 minutes)

The Collinear Resonance Ionization Spectroscopy experiment (CRIS) at ISOLDE combines the high sensitivity of resonance ionization spectroscopy with the high resolution offered by collinear laser spectroscopy. The first experiments at CRIS demonstrated the ability to reach exotic isotopes, normally out of reach for collinear laser spectroscopy methods based on photon detection, with an intermediate resolution [1]. Further developments have focused on improving the resolving power, to the point where it now matches the resolution of other collinear laser spectroscopy methods [2]. With this performance, the CRIS experiment is ideally suited to study the evolution of nuclear structure in regions far from stability.

Several ISOLDE experiments have been working towards the region around the doubly magic ^{78}Ni . Previous laser spectroscopy work [3-7] clearly demonstrated the inversion of the $\pi f_{5/2}$ and the $\pi p_{3/2}$ orbitals between ^{73}Cu and ^{75}Cu as the $vg_{9/2}$ orbital is filled. This inversion is currently understood in terms of the tensor interaction between the neutrons and protons [8] which could potentially result in a quenching of the $Z=28$ shell gap towards $N=50$ [9].

This contribution will focus on the application of the high-resolution CRIS technique to the study of neutron-rich copper isotopes in the vicinity of $N=50$. The g -factors, quadrupole moments and charge radii of these neutron rich copper isotopes will provide additional information to gauge the robustness of the magicity of the $Z=28$ shell in ^{78}Ni . During the last campaign in April 2016, measurements have been performed on 15 Cu isotopes, including for the first time high resolution measurements of the very exotic isotopes $^{76,77,78}\text{Cu}$, where ^{78}Cu was produced at a rate of only 20 ions/s. These measurements provide information on the spin, magnetic moment, quadrupole moment and charge radius. The obtained data will be compared to large scale shell model calculations.

References:

- [1] K.T. Flanagan et al., PRL 111, 212501 (2013)
- [2] R.P. de Groote et al., PRL 115, 132501 (2015)
- [3] K.T. Flanagan, PRL 103, 103, 142501 (2009)
- [4] P. Vingerhoets, PRC 82, 064311 (2010)
- [5] P. Vingerhoets, PLB 703, 1 (2013)
- [6] K. T. Flanagan, PRC 82, 041302(R) (2010)
- [7] U. Köster et al, PRC 84, 034320 (2011)
- [8] T. Otsuka et al, PRL 95, 232502 (2005)
- [9] K. Sieja et al., PRC 81, 061303(R) (2010)
- [10] J. Hakala et al., PRL 101, 052502 (2008)
- [11] Z. Y. Xu et al., PRL 113, 032505 (2014)

Primary author: Mr DE GROOTE, Ruben (KU Leuven)

Presenter: Mr DE GROOTE, Ruben (KU Leuven)

Session Classification: Breakout 2