



MEETING OF THE AMERICAN PHYSICAL SOCIETY DIVISION OF PARTICLES AND FIELDS

Contribution ID: 172

Type: **Presentation**

## Constraints on the astrophysical flux and the dark matter decay with IceCube HESE data

*Monday, 31 July 2017 11:05 (20 minutes)*

The IceCube detection of High Energy Starting Events (HESE) and the upward muon track events (6 year data) are presently hard to explain with the single power-law astrophysical flux for energies above 30TeV. We investigate the possibility that a significant component of the additional neutrino flux originates due to the decay of a very heavy dark matter particle via several possible channels into standard model particles. We perform a full 4 parameter fit to IceCube data in which we vary astrophysical flux normalization, power-law index, dark matter mass, dark matter lifetime and dark matter decay mode. We show that that dark matter with mass in the PeV range and the lifetime around  $10^{27}$ s provides much better fit to IC data than the best-fit astrophysical flux alone. We also find dark matter lifetime limits which are much stronger than those obtained from gamma-ray data for all channels except  $b\bar{b}$ , which is also disfavored by the IceCube HESE data.

**Primary author:** Prof. SARCEVIC, INA (UNIVERSITY OF ARIZONA)

**Presenter:** Prof. SARCEVIC, INA (UNIVERSITY OF ARIZONA)

**Session Classification:** Cosmology and Astrophysics

**Track Classification:** Cosmology and Astrophysics