




K. Scholberg, Duke University  
On behalf of the COHERENT collaboration  
August 2, 2017  
DPF 2017, Fermilab

# Coherent elastic neutrino-nucleus scattering (CEvNS)

$$\nu + A \rightarrow \nu + A$$

A neutrino smacks a nucleus via exchange of a  $Z$ , and the nucleus recoils as a whole;  
**coherent** up to  $E_\nu \sim 50$  MeV




- Important in **SN processes & detection**
- Well-calculable cross-section in SM:  
SM test, probe of neutrino NSI
- Dark matter direct detection background
- Possible applications (reactor monitoring)


$$\frac{d\sigma}{d\Omega} = \frac{G^2}{4\pi^2} k^2 (1 + \cos \theta) \frac{(N - (1 - 4 \sin^2 \theta_W) Z)^2}{4} F^2(Q^2)$$

$$\propto N^2$$

The cross-section is *large*




**Large cross section, but never observed  
due to tiny nuclear recoil energies:**



➔ but **WIMP dark matter detectors** developed over the last ~decade are sensitive to ~ keV to 10's of keV recoils


# CEvNS from natural neutrinos creates ultimate background for direct DM search experiments



Understand nature of background (& detection response)

Clean SM prediction for the rate  $\rightarrow$  measure  $\sin^2\theta_{W\text{eff}}$  ;  
**deviation probes new physics**

$$\sigma \sim \frac{G_f^2 E^2}{4\pi} (N - (1 - 4 \sin^2 \theta_W) Z)^2$$




Example: hypothetical dark Z mediator (explanation for g-2 anomaly)

CEvNS sensitivity is @ low Q;  
 need sub-percent precision to compete w/  
 electron scattering & APV, but **new channel**

# Non-Standard Interactions of Neutrinos:

new interaction **specific to  $\nu$ 's**

$$\mathcal{L}_{\nu H}^{NSI} = -\frac{G_F}{\sqrt{2}} \sum_{q=u,d} [\bar{\nu}_\alpha \gamma^\mu (1 - \gamma^5) \nu_\beta] \times (\varepsilon_{\alpha\beta}^{qL} [\bar{q} \gamma_\mu (1 - \gamma^5) q] + \varepsilon_{\alpha\beta}^{qR} [\bar{q} \gamma_\mu (1 + \gamma^5) q])$$




Can improve ~order of magnitude beyond CHARM limits with a first-generation experiment (for best sensitivity, want **multiple targets**)

# Oscillations to sterile neutrinos w/CEvNS


(NC is flavor-blind): a potential new tool;

look for deficit and spectral distortion vs L,E

Examples:



Multi- $\pi$ DAR sources at different baselines (20 & 40 m)




B. Dutta et al, arXiv:1511.02834

# Neutrino magnetic moment

Signature is **distortion at low recoil energy E**

$$\frac{d\sigma}{dE} = \frac{\pi\alpha^2\mu_\nu^2 Z^2}{m_e^2} \left( \frac{1 - E/k}{E} + \frac{E}{4k^2} \right)$$



→ requires low energy threshold

See also Kosmas et al., arXiv:1505.03202

# Nuclear physics with coherent elastic scattering

If systematics can be reduced to ~ few % level,  
we can start to explore nuclear form factors

P. S. Amanik and G. C. McLaughlin, J. Phys. G 36:015105


K. Patton et al., PRC86 (2012) 024612

$$\frac{d\sigma}{dT}(E, T) = \frac{G_F^2}{2\pi} M \left[ 2 - \frac{2T}{E} + \left( \frac{T}{E} \right)^2 - \frac{MT}{E^2} \right] \frac{Q_W^2}{4} F^2(Q^2)$$


Form factor: encodes information  
about nuclear (primarily neutron)  
distributions

Fit recoil ***spectral shape*** to determine the  $F(Q^2)$  moments  
(requires very good energy resolution, good systematics control)

Example:  
tonne-scale  
experiment  
at  $\pi$ DAR source




# Tonne-scale underground DM detectors can measure **solar and supernova neutrinos**



Billard et al., arXiv:1409.0050

**Solar neutrinos:**  
rule out sterile oscillations  
using CEvNS (NC)




Horowitz et al., PRD68 (2003) 023005

**Supernova neutrinos:**  
~ handful of events per tonne  
@ 10 kpc: sensitive to  
***all flavor components of the flux***




Why use the 10's of MeV neutrinos from  $\pi$  decay at rest?

→ higher-energy neutrinos are advantageous, because both **cross-section and maximum recoil energy increase with  $\nu$  energy**





Reactor experiments (RICOCHET, CONNIE, CONus etc.) can take advantage of very large flux ( $\sim$ factor of  $10^4$ ) but require very low energy thresholds, where background can be daunting; radioactive source experiments require even lower thresholds

# Stopped-Pion ( $\pi$ DAR) Neutrinos




2-body decay: monochromatic 29.9 MeV  $\nu_\mu$   
PROMPT




3-body decay: range of energies  
between 0 and  $m_\mu/2$   
DELAYED ( $2.2\mu\text{s}$ )

# Stopped-Pion Sources Worldwide



# Comparison of pion decay-at-rest $\nu$ sources


from duty cycle





# Spallation Neutron Source

Oak Ridge National Laboratory, TN



Proton beam energy: 0.9-1.3 GeV


Total power: 0.9-1.4 MW

Pulse duration: 380 ns FWHM


Repetition rate: 60 Hz

Liquid mercury target

# The SNS has **large, extremely clean** DAR $\nu$ flux




SNS flux (1.4 MW):  
 **$430 \times 10^5 \nu/\text{cm}^2/\text{s}$**   
**@ 20 m**



Note that contamination  
from non  $\pi$ -decay at rest  
(decay in flight,  
kaon decay,  $\mu$  capture...)  
is **down by several  
orders of magnitude**

# Time structure of the SNS source


60 Hz *pulsed* source



**Background rejection factor  $\sim \text{few} \times 10^{-4}$**

# The COHERENT collaboration

<http://sites.duke.edu/coherent>



~80 members,  
18 institutions  
4 countries




arXiv:1509.08702



# COHERENT Detectors

Nuclear Target	Technology	Mass (kg)	Distance from source (m)	Recoil threshold (keVr)
<b>CsI[Na]</b>	Scintillating Crystal	14.6	20	6.5
<b>Ge</b>	HPGe PPC	10	22	5
<b>LAr</b>	Single-phase	22	29	20
<b>NaI(Tl)</b>	Scintillating crystal	185*/2000	28	13



Multiple detectors for  $N^2$  dependence of the cross section




# Siting for deployment in SNS basement

(measured neutron backgrounds low,  
~ 8 mwe overburden)

View looking  
down “Neutrino Alley”




# Expected recoil signals



Prompt defined as first  $\mu$ s; note some contamination from  $\nu_e$  and  $\bar{\nu}_\mu$

# COHERENT Detector Status

Nuclear Target	Technology	Mass (kg)	Distance from source (m)	Recoil threshold (keVr)	Data-taking start date
CsI[Na]	Scintillating Crystal	14.6	20	6.5	9/2015
Ge	HPGe PPC	10	22	5	2017
LAr	Single-phase	22	29	20	12/2016
NaI[Tl]	Scintillating crystal	185*/2000	28	13	*high-threshold deployment summer 2016



- CsI installed in July 2015
- 185 kg of NaI installed in July 2016
- LAr single-phase detector installed in December 2016,  
upgraded w/TPB coating of PMT & Teflon; commissioning underway
- Ge detectors to be installed late 2017

**CsI results soon: embargoed until Aug 3, 2 pm EST**

# Currently measuring *neutrino-induced neutrons* in lead, (iron, copper), ...




↓  
1n, 2n emission



↓  
1n, 2n,  $\gamma$  emission




- potentially a non-negligible background, especially in lead shield
- valuable in itself, e.g. HALO SN detector



Talk by Brandon Becker next!

# Potential upgrades


- additional Ge detectors
- larger LAr (up to few 100 kg)
- up to 7 ton NaI
- additional targets/detectors



# Summary

- **CEvNS** never before measured
- Multiple physics motivations
  - DM bg, SM test, astrophysics, nuclear physics, ...
- Now within reach with WIMP detector technology and neutrinos from pion decay at rest

**COHERENT@ SNS** going after this  
with multiple targets, extremely clean neutrino flux




Talk by Phil Barbeau Fri morning plenary

# **Extras/backups**

# CEvNS from natural neutrinos creates ultimate background for direct DM search experiments

J. Billard, E. Figueroa-Feliciano, and L. Strigari, arXiv:1307.5458v2 (2013).



Understand nature of background (& detector response)

# Neutron Backgrounds

Several background measurement campaigns have shown that Neutrino Alley is neutron-quiet

