

DMUK Meeting - UCL

Wednesday, 18 January 2017 - Wednesday, 18 January 2017

University College London

Book of Abstracts

Contents

Welcome	1
Dark Matter Heating	1
Dark Matter searches at the LHC	1
Axions and non-WIMP searches	1
WIMP Direct Detection	1
Results from the LUX experiment	1
The SABRE dark matter search experiment	1
Indirect dark matter searches and CTA	2
Direct Detection of Nuclear Dark Matter Using Tonne-Scale Experiments	2
Directional dark matter searches with the DRIFT and CYGNUS-TPC experiments	2
The DEAP-3600 experiment at SNOLab	2
Low-background radio-assay capability in the UK	2
The LZ experimental hardware systems	2
Software development for the LZ experiment	3
Directional detectors with polarised targets	3

5

Welcome

Author: Chamkaur Ghag¹

¹ *UCL*

Corresponding Author: c.ghag@ucl.ac.uk

6

Dark Matter Heating

Corresponding Author: j.read@surrey.ac.uk

7

Dark Matter searches at the LHC

Corresponding Author: sarah.malik@imperial.ac.uk

8

Axions and non-WIMP searches

Corresponding Author: e.daw@sheffield.ac.uk

9

WIMP Direct Detection

Corresponding Author: asher.kaboth@rhul.ac.uk

10

Results from the LUX experiment

Corresponding Author: sally.shaw.13@ucl.ac.uk

11

The SABRE dark matter search experiment

Corresponding Author: f.froberg@imperial.ac.uk

12

Indirect dark matter searches and CTA

Corresponding Author: anthony.brown@durham.ac.uk

13

Direct Detection of Nuclear Dark Matter Using Tonne-Scale Experiments

Corresponding Author: alistairbutcher@gmail.com

Summary:

Nuclear dark matter models propose a possible composite form of dark matter, dark matter nuclei, which are analogues to Standard Model nuclei. We present possible nuclear dark matter direct detection signals in the DEAP-3600 and XENON1T experiments for a particular class of nuclear dark matter. The number of events required to distinguish between this case and a standard point-like WIMP state is presented for each experiment. We find that, in the most favourable regions of the parameter space, it is possible to distinguish nuclear dark matter from WIMPs at the 3σ level using both experiments in combination, while at best a 2σ distinction is possible individually.

14

Directional dark matter searches with the DRIFT and CYGNUS-TPC experiments

Corresponding Author: a.scarff@sheffield.ac.uk

15

The DEAP-3600 experiment at SNOLab

Corresponding Author: franco.lazia.2014@live.rhul.ac.uk

16

Low-background radio-assay capability in the UK

Corresponding Author: paul.scovell@physics.ox.ac.uk

17

The LZ experimental hardware systems

Corresponding Author: b.lopez-paredes@imperial.ac.uk

18

Software development for the LZ experiment

Corresponding Author: j.dobson@ucl.ac.uk

19

Directional detectors with polarised targets

Corresponding Author: tarsfranarin@gmail.com