New Perspectives 2017



Contribution ID: 29

Type: Oral Presentation

Decomposition Methods for the ν_e Appearance analysis in the NOvA Near Detector

Tuesday, 6 June 2017 10:00 (15 minutes)

NOvA is a long-baseline neutrino oscillation experiment that is designed to probe the neutrino mass hierarchy and mixing structure. It uses two functionally identical liquid scintillator detectors 14mrad off-axis from the NuMI beamline at Fermilab, allowing a tightly focused neutrino flux peaked at around 2 GeV. The Near Detector is located 100 m underground and is used to characterize the neutrino beam before oscillations. Since the beam components are affected differently by oscillations, the data collected at the Near Detector needs to be broken down into these components. This enables a precise prediction to be made of the beam-induced backgrounds to the ν_e appearance signal at the Far Detector, 810 km from the neutrino source. Various data-driven techniques are employed for this purpose, in particular by constraining the flux yields and correcting for observed neutrino interaction characteristics. In this talk, I will present an overview of the methodology used to decompose the Near Detector data and predict the FD spectrum in the latest ν_e appearance analysis, utilizing an accumulated exposure of 6.05×10^{20} protons-on-target.

Primary author: Mr NAYAK, Nitish (University of California-Irvine)
Presenter: Mr NAYAK, Nitish (University of California-Irvine)
Session Classification: Long Baseline Neutrino Program