

WA105


Effect of LEM borders, screw holes and HV connectors on collection charge efficiency

Philippe COTTE

March 23, 2017

Main question

WA105 LEM geometry




Study


Border, screw holes, HV connectors : What is their **impact** on **charge collection** and **charge resolution**?

Main question

Motivation : LAr LEM-TPC




A. Badertscher, ArXiv:1301.4817v1 [physics.ins-det] 21 Jan 20



A. Badertscher, ArXiv:1301.4817v1 [physics.ins-det] 21 Jan 2013

A. Badertscher et al. JINST 8 (2013) P04012

200L DLAr LEM-TPC
40 × 76 cm² readout



Rectangular LEM with 1.6 mm gap between each LEM ⇒ Absence of charge at LEM border



1. **Simulate the electric field** close to the border, screw holes and HV connectors (done with ANSYS)
2. **Simulate the electrons drifting toward the LEM** in those region, compute collection efficiency depending on initial electron position (done with GarField)
3. **Create an efficiency map** for a typical LEM of $49.95 \times 49.95 \text{ cm}^2$ (done with Root)
4. **Simulate events in $6 \times 6 \times 6$** and compare charge and charge resolution with/without efficiency (done with QScan)

Geometry : example of 2D model








Figure: 2D modelisation of a LEM border


Geometry : example of 2D model

field, potential and drift of 10 electrons






End point of electron path
used as condition :

- ▶ Consider any electron reaching amplification zone as **collected**
- ▶ **Kill** electron reaching bottom LEM on dead zone (border, screw, HV connector)
- ▶ Compare **initial position of all electrons** to initial position of **collected electrons** (ratio of histograms collected/generated)




Geometry : 3D models

Border, corner, screw hole and HV connector




← Screw Hole HV connector →




Drift of 10000 electrons on LEM border

uniformly distributed at bottom of the geometry



Drift of 10000 electrons on LEM border


uniformly distributed at bottom of the geometry



Fit function : $\sim \arctan(-x)$

Results for 3D models



Border, corner, screw hole and HV connector



HV connector

Results for 3D models



Border, corner, screw hole and HV connector




HV connector

Results for 3D models


Border, corner, screw hole and HV connector



Screw Hole




HV connector




- ▶ Pixel efficiency computed by integrating histogram pixel by pixel
(pixel = $3.125 \times 3.125 \text{ mm}^2$)


Real LEM




Efficiency map for one LEM



Originally :



Now :



And then again split
in strips (since it
is what the detector
will eventually give
us)



- ▶ Possibility to specify efficiency pixel by pixel (computation time +15%)



- ▶ Possibility to specify efficiency pixel by pixel (computation time +15%)
- ▶ Efficiency map read as TH2D once for each event (no impact on computation time)




- ▶ Possibility to specify efficiency pixel by pixel (computation time +15%)
- ▶ Efficiency map read as TH2D once for each event (no impact on computation time)
- ▶ Possibility to record charge with or without efficiency map



- ▶ Possibility to specify efficiency pixel by pixel (computation time +15%)
- ▶ Efficiency map read as TH2D once for each event (no impact on computation time)
- ▶ Possibility to record charge with or without efficiency map
- ▶ Compare charge distribution with "perfect" (**no efficiency map**) and "real" (**with efficiency map**) LEM for different particles and momenta

1000 particles simulated






Electrons, muons, kaons, pions and protons



- ▶ Initial momentum fixed
- ▶ Direction random in 4π
- ▶ Initial position at $z = 0$
- ▶ and $(x, y) \in [-50; 50]cm$

Results

Charge distribution for **electrons** at different momenta




Results

Charge difference of **electrons** for different momenta




Mean charge










Relative dispersion of collected charge



Results

Charge distribution for **negative pions** at different momenta




Results

Charge difference of **negative pions** for different momenta




Mean charge






Relative dispersion of collected charge






Results

Charge difference of different particles



Results

Charge dispersion of different particles





- ▶ Effect on resolution small on most particles above $1\text{ GeV}/c$
Except electrons : resolution doubled



- ▶ Effect on resolution small on most particles above $1\text{ GeV}/c$
Except electrons : resolution doubled
- ▶ Effect on total charge can be well known



- ▶ Effect on resolution small on most particles above $1\text{ GeV}/c$
Except electrons : resolution doubled
- ▶ Effect on total charge can be well known
- ▶ Dead zone between LEMs : $4 + 4 + 0.5 = 8.5\text{ mm}$, screw holes : 6 mm , HV connectors : 10 mm \Rightarrow vertex happening in those zones can be difficult to analyse, especially showers

The End



Thank you!