



Contribution ID: 64

Type: poster

## Improving Uncertainties in the $^{17}\text{O}(\alpha, n)^{20}\text{Ne}$ cross section

*Wednesday, 23 May 2018 16:30 (1h 30m)*

The  $^{17}\text{O}(\alpha, n)$  reaction recycles neutrons for the most efficient neutron absorber ( $^{16}\text{O}(n, \gamma)^{17}\text{O}$ ) during both the core He-burning and shell C-burning phases of the weak s-process. In order to improve uncertainties in the reaction rate, we are performing  $(\alpha, n)$  and  $(\alpha, n'\gamma)$  cross section measurements at the 5U accelerator of the Notre Dame Nuclear Science Laboratory. We will report on a recent measurement that was performed with  $\text{LaBr}_3:\text{Ce}$  gamma-ray detectors from the HAGRID array in combination with a deuterated liquid scintillator neutron detector. The employed digital electronics allow for pulse shape discrimination in both detector types, and spectral unfolding is used to obtain neutron energy information from the observed light output spectrum in the liquid scintillation detector.

**Primary authors:** BOELTZIG, Axel (University of Notre Dame); MACON, Kevin (University of Notre Dame)

**Co-authors:** MICHAEL, Febraro (Oak Ridge National Lab); JONES, Kate (University of Tennessee Knoxville); Prof. WIESCHER, Michael (University of Notre Dame); Dr DEBOER, Richard J. (University of Notre Dame); Dr MOSBY, Shea (P27 Los Alamos National Lab); Dr PEREIRA-LOPEZ, Xesús (University of Tennessee Knoxville)

**Presenter:** MACON, Kevin (University of Notre Dame)

**Session Classification:** Poster Session