

Meson electromagnetic form factors from lattice QCD

Tuesday, 24 July 2018 18:45 (2 hours)

Lattice QCD can provide direct determination of the electromagnetic form factors of mesons as a prediction to be compared to upcoming experiments at Jefferson Lab. At the same time we can compare to expectations from perturbative QCD, which take a very simple form at high Q^2 .

We will show recent results from HPQCD, building on the work in 1701.04250.

We give predictions for π and K mesons up to 4 GeV^2 that include calculations for physical u/d quarks and provide accurate predictions for Jefferson Lab.

We also extend the Q^2 range up to 20 GeV^2 by studying mesons made of heavier quarks. At these values of Q^2 , discrepancies (both in the values and the Q^2 -dependence) with perturbative QCD raise issues of the reliability of the assumptions going in to the perturbative QCD calculations. This has wider implications also for other processes.

Primary author: Prof. DAVIES, Christine (University of Glasgow)

Co-authors: Mr ZIMERMAN-SANTOS, Andre (University of Sao Paulo); Dr LYTLE, Andrew (University of Glasgow); Dr KOPONEN, Jonna (INFN sezione di Roma Tor Vergata); Prof. LEPAGE, Peter (Cornell University)

Presenter: Prof. DAVIES, Christine (University of Glasgow)

Session Classification: Poster reception

Track Classification: Hadron Structure