

$\mathcal{O}(a)$ improved quark mass renormalization for a non-perturbative matching of HQET to three-flavor QCD

Thursday, July 26, 2018 10:10 AM (20 minutes)

The use of Heavy Quark Effective Theory (HQET) on the lattice as an approach to B-physics phenomenology is based on a non-perturbative matching of HQET to QCD in finite volume. As a first step to apply the underlying strategy in the three-flavor ($N_f = 2 + 1$) theory, we determine the renormalization constant and improvement coefficients relating the renormalized current and subtracted quark mass of (quenched) valence quarks in $\mathcal{O}(a)$ improved $N_f = 3$ lattice QCD. We present our strategy and first results for the relevant parameter region towards weak couplings along a line of constant physics, which corresponds to lattice resolutions $a \leq 0.02$ fm and fixes the physical extent of the matching volume to $L \approx 0.5$ fm.

Primary authors: Dr HEITGER, Jochen (Institut für Theoretische Physik, Universität Münster); Dr FRITZSCH, Patrick (Theoretical Physics Department, CERN); KUBERSKI, Simon (Institut für Theoretische Physik, Universität Münster)

Presenter: KUBERSKI, Simon (Institut für Theoretische Physik, Universität Münster)

Session Classification: Standard Model Parameters and Renormalization

Track Classification: Standard Model Parameters and Renormalization