

Calculating the ρ radiative decay width with lattice QCD

Wednesday, July 25, 2018 3:20 PM (20 minutes)

We present the results of our lattice QCD study of the $\pi\gamma \rightarrow \pi\pi$ process, where the ρ resonance appears as an enhancement in the transition amplitude. We use Nf=2+1 clover fermions on a lattice of L=3.6 fm and a pion mass of 320 MeV. Using a combination of forward, stochastic and sequential propagators we calculate the two-point and three-point functions required in the determination of the $\pi\gamma \rightarrow \pi\pi$ amplitude and determine the $\pi\gamma \rightarrow \pi\pi$ matrix elements in a region of invariant mass s and momentum transfer q^2 . To fit the q^2 and s dependence of the amplitude we are exploring a set of general models based on a Taylor expansion and their description of the data. By analytic continuation to the complex pole corresponding to the ρ resonance we determine the resonant form factors and calculate the radiative decay width of the ρ resonance.

Primary author: Dr LESKOVEC, Luka (University of Arizona)

Co-authors: POCHINSKY, Andrew (MIT); Prof. ALEXANDROU, Constantia (Cyprus University); Mr RENDON SUZUKI, Jesus Gumaro (University of Arizona); Prof. NEGELE, John Negele (MIT); Dr PETSCHLIES, Marcus (University Bonn); Mr PAUL, SRIJIT (The Cyprus Institute); Prof. SYRITSYN, Sergey (Stony Brook University (SUNY)); MEINEL, Stefan (University of Arizona)

Presenter: Dr LESKOVEC, Luka (University of Arizona)

Session Classification: Hadron Spectroscopy and Interactions

Track Classification: Hadron Spectroscopy and Interactions