

# Scattering in Euclidean formulations of relativistic quantum theory

*Friday, 27 July 2018 14:40 (20 minutes)*

Scattering theory can be formulated given a representation of the quantum mechanical Hilbert space and a set of self-adjoint Poincaré generators satisfying cluster properties. These are both provided by the Osterwalder-Schrader reconstruction theorem, where the input is a collection of Euclidean-covariant reflection-positive distributions. In this representation both Hilbert space inner products and matrix elements of the generators can be expressed directly in terms of Euclidean variables, without analytic continuation. A Euclidean version of Haag-Ruelle scattering can be formulated in this representation, which leads to expressions for scattering observables as strong limits. I discuss the construction of one-particle Haag-Ruelle states in this representation and show how these states can be used to construct wave operators. I exhibit toy model calculations of sharp-momentum transition matrix elements over a wide range of energies that suggest the feasibility of formulating numerical methods to compute scattering observables in this Euclidean representation.

**Primary author:** Prof. POLYZOU, Wayne (University of Iowa)

**Co-authors:** Dr AIELLO, Gordon (University of Iowa); Mr KOPP, Philip (University of Iowa)

**Presenter:** Prof. POLYZOU, Wayne (University of Iowa)

**Session Classification:** Theoretical Developments

**Track Classification:** Theoretical Developments