

Computing \hat{q} on a quenched SU(3) lattice

Thursday, July 26, 2018 10:10 AM (20 minutes)

The jet transport coefficient \hat{q} is the leading parameter that controls the modification of hard jets produced in heavy-ion collisions. This coefficient, like other jet coefficients is inherently non-perturbative, and hence, is challenging to compute from first principles. Currently, existing theoretical model to data comparisons require a separate normalization of \hat{q} between RHIC and LHC energies, beyond the obvious T^3 scaling from dimensional arguments. This is known as the jet \hat{q} puzzle. In this talk, we present a pQCD and lattice gauge theory based formulation to study \hat{q} which sheds new light on the non-perturbative nature of \hat{q} and the jet puzzle. For this first attempt, we formulate \hat{q} within a quenched SU(3) lattice. We consider a leading order diagram for a hard parton passing through the thermal medium. The non-perturbative part is expressed in terms of a non-local (two-point) Field-Strength-Field-Strength operator product which can be Taylor expanded after analytic continuation to the Euclidean region. Such an expansion allowed us to write \hat{q} in terms of the expectation of local operators. We also carry out a perturbative analysis both on the lattice and in continuum field theory to understand the scale dependence of the jet transport coefficient.

Primary authors: Dr MAJUMDER, Abhijit (Wayne State University); Mr KUMAR, Amit (Wayne State University)

Presenter: Mr KUMAR, Amit (Wayne State University)

Session Classification: Nonzero Temperature and Density

Track Classification: Nonzero Temperature and Density