

Exploring the phase diagram of finite density QCD at low temperature by the complex Langevin method

Monday, 23 July 2018 15:00 (20 minutes)

Monte Carlo studies of QCD at finite density suffer from the notorious sign problem, which becomes easily uncontrollable as the chemical potential increases for a moderate lattice size. In this work, we attempt to approach the high density low temperature region by the complex Langevin method (CLM). Simulations are performed on an $8^3 \times 16$ lattice using four-flavor staggered fermions with reasonably small quark mass. Unlike previous work with a $4^3 \times 8$ lattice, the criterion for correct convergence is satisfied in the nuclear matter phase without using the deformation technique. In this phase the baryon number density has a plateau with respect to the chemical potential, and it starts to grow rapidly at some point.

Primary author: Dr ITO, Yuta (KEK)

Co-authors: Prof. TSUCHIYA, Asato (Shizuoka University); Dr MATSUFURU, Hideo (KEK); Prof. NISHIMURA, Jun (KEK, SOKENDAI); Dr SHIMASAKI, Shinji (Keio University, KEK); Dr TSUTSUI, Shoichiro (High Energy Accelerator Research Organization)

Presenter: Dr ITO, Yuta (KEK)

Session Classification: Nonzero Temperature and Density

Track Classification: Nonzero Temperature and Density