



Contribution ID: 27

Type: Oral Presentation

Long-baseline searches for sterile neutrinos using neutral current interactions in NOvA

Tuesday, 19 June 2018 17:15 (15 minutes)

This talk discusses an updated analysis of neutrino data from the NOvA experiment to search for sterile neutrino oscillations. NOvA consists of two functionally identical liquid scintillator detectors in Fermilab's NuMI neutrino beam: a 300 ton near detector at a 1km baseline, and a 14,000 ton far detector 810km away in Ash River, MN, 14.6 mrad off the beam's central axis. Sterile neutrino oscillations are constrained by searching for neutrino disappearance in neutral current interactions between the near and far detectors. A covariance matrix approach is utilised to cancel systematic uncertainties between the two detectors over a broad range of Δm^2_{41} parameter space. This analysis uses neutrino data collected between February 2014 and February 2017, corresponding to an exposure equivalent to 8.85×10^{20} protons on target. A summary is provided of methods and results, including limits on the sterile neutrino mixing parameters θ_{24} and θ_{34} as a function of Δm^2_{41} .

Primary authors: Dr HEWES, Jeremy (University of Cincinnati); WALLBANK, Michael (University of Cincinnati)

Presenters: Dr HEWES, Jeremy (University of Cincinnati); WALLBANK, Michael (University of Cincinnati)

Session Classification: Long Baseline Neutrino Program