




GW170817: Implications for the astrophysical site of the r-process



Daniel M. Siegel

NASA Einstein Fellow

Center for Theoretical Physics & Columbia Astrophysics Laboratory, Columbia University

FRIB and the GW170817 kilonova, MSU, July 16-27, 2018


I.

Observations

GW170817 and the fireworks of EM counterparts


- **unique event in astronomy**, unprecedented level of **multi-messenger observations**
- confirms **association of BNS to SGRBs**
 - Archival data: SGRB spectral properties may not be unusual Burns+ 2018
 - VLBI obs.: SGRB morphology may not be unusual Mooley+ 2018b
- **kilonova** provides strong evidence for synthesis of **r-process material**

Lightcurve from *Fermi*/GBM (50 – 300 keV)



SGRB




Gravitational-wave time-frequency map



kilonova

X-rays


radio



The r-process and s-process

Burbidge, Burbidge, Fowler, Hoyle (1957), Cameron (1957):

The heavy elements ($A > 62$) are formed by neutron capture onto seed nuclei



Burbidge, Burbidge, Fowler, Hoyle (“B²FH”)

REVIEWS OF MODERN PHYSICS

VOLUME 29, NUMBER 4

OCTOBER, 1957

Synthesis of the Elements in Stars*

E. MARGARET BURBIDGE, G. R. BURBIDGE, WILLIAM A. FOWLER, AND F. HOYLE

*Kellogg Radiation Laboratory, California Institute of Technology, and
Mount Wilson and Palomar Observatories, Carnegie Institution of Washington,
California Institute of Technology, Pasadena, California*

“It is the stars, The stars above us, govern our conditions”;
(*King Lear*, Act IV, Scene 3)

but perhaps

“The fault, dear Brutus, is not in our stars, But in ourselves,”
(*Julius Caesar*, Act I, Scene 2)

TABLE OF CONTENTS

	<i>Page</i>
I. Introduction.....	548
A. Element Abundances and Nuclear Structure.....	548

1957 February

THE ASTRONOMICAL JOURNAL

9

and aggregates. The 21-cm absorption spectra will be investigated for accessible discrete radio sources. Studies relating to the spiral structure of our galaxy will be limited largely to regional surveys for small sections of the sky; we shall stress in these studies at all times the close interconnection that exists between radio and optical phenomena. Following Heesch's successful detection of 21-cm emission from the Coma cluster of galaxies, we shall attempt to study further 21-cm radiation from beyond our own galactic system, but in these studies we shall be limited to some extent by our electronic equipment, which was designed especially for high-resolution work in our own galaxy.

The new equipment is described in some detail in *Sky and Telescope* for July 1956, and an article is in press in *Nature*.

*Harvard College Observatory,
Cambridge, Mass.*

in the 1100Å to 1340Å detector which included the Lyman α line of hydrogen, 1216Å. The 1220Å to 1340Å tube detected discrete celestial sources. Of the region scanned by this tube the most significant responses were obtained in the Puppis-Vela region.

*Naval Research Laboratory,
Washington, D. C.*

Cameron, A. G. W. On the origin of the heavy elements.

The inverse correlation between the metal abundances and the ages of stars suggests that the elements have been formed in stellar interiors. An analysis of the cosmic abundances of nuclear isobars, and calculations relating to the growth of nuclide abundances by neutron capture, show that the following three mechanisms are necessary, and probably sufficient, to produce the observed cosmic abundances of the nuclides with mass number greater than 70.



Cameron

speculated that **r-process** requires explosive environment of supernovae

The kilonova of GW170817

- **blue** kilonova properties:

$M_{ej} \sim 10^{-2} M_{\text{sun}}$ Kilpatrick+ 2017

$v_{ej} \sim 0.2-0.3c$ Kasen+ 2017

$Y_e > 0.25$ Nicholl+ 2017

$X_{La} < 10^{-4}$ Villar+ 2017

Coughlin+ 2018

- **red/purple** kilonova properties:

$M_{ej} \sim 4-5 \times 10^{-2} M_{\text{sun}}$ Kilpatrick+ 2017

$v_{ej} \sim 0.08-0.14c$ Kasen+ 2017

$Y_e < 0.25$ Kasliwal+ 2017


$X_{La} \sim 0.01$ Drout+ 2017

Cowperthwaite+ 2017

Chornock+ 2017

Villar+ 2017

Coughlin+ 2018



- two (“red-blue”) or multiple components **expected from merger simulations**

- single component models might be possible, Smartt+ 2017
but require fine-tuning of Y_e Waxman+ 2017


II.

Mass ejection

GW170817 phenomenology


Sources of ejecta in NS mergers

dynamical ejecta (~ms)



Hotokezaka+ 2013, Bauswein+ 2013


winds from NS remnant (~10ms-1s)



Dessart+ 2009

Siegel+ 2014
Ciolfi, Siegel+ 2017

accretion disk (~10ms-1s)



Siegel & Metzger 2017, 2018a

tidal ejecta
shock-heated ejecta

$$M_{\text{tot}} \lesssim 10^{-3} M_{\odot}$$

$$v \gtrsim 0.2c$$

neutrino-driven wind

$$\dot{M}_{\text{in}} \sim (10^{-4} - 10^{-3}) M_{\odot} \text{s}^{-1}$$

magnetically driven wind

$$\dot{M}_{\text{in}} \sim (10^{-3} - 10^{-2}) M_{\odot} \text{s}^{-1}$$

combined see Metzger 2018

thermal outflows

$$M_{\text{tot}} \gtrsim 0.3 - 0.4 M_{\text{disk}}$$

$$v \sim 0.1c$$

Overall ejecta mass per event:

$$\lesssim 10^{-3} - 10^{-2} M_{\odot}$$

strongly depends on EOS and mass ratio

Bauswein+ 2013
Radice+ 2016, 2017
Sekiguchi+ 2016
Palenzuela+2015
Lehner+2016
Ciolfi, Siegel+2017

Siegel & Metzger 2017, 2018

$$\gtrsim 10^{-2} M_{\odot}$$

lower limit


Dynamical ejecta and winds



Movie: BNS merger showing dynamical ejecta and winds from remnant NS


Ciolfi, Siegel+2017

Accretion disk formation



Movie: BNS merger and formation of
post-merger accretion disk [Radice+ 2016](#)

Accretion disk outflows




Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a

Movie: [long-term evolution](#) of post-merger accretion disk, $M_{\text{BH}}=3M_{\text{sun}}$ (spin: 0.8), $M_{\text{disk}}=0.02M_{\text{sun}}$

Sources of ejecta for kilonova in GW170817


dynamical ejecta (~ms)



$$M_{\text{tot}} \lesssim 10^{-3} M_{\odot}$$

$$v \gtrsim 0.2c$$


winds from NS remnant (~ms-ls)



$$\dot{M}_{\text{in}} \sim (10^{-3} - 10^{-2}) M_{\odot} \text{s}^{-1}$$

$$v \lesssim 0.1c$$

accretion disk (~10ms-ls)



$$M_{\text{tot}} \gtrsim 10^{-2} M_{\odot}$$

$$v \sim 0.1c$$


blue KN in GW170817

- requires large amount of **shock heated ejecta** to obtain high $Y_e > 0.25$
- requires **metastable NS phase**
- requires **EOS with small NS radius (~12 km)**

red KN in GW170817

- produces the heavy r-process elements in GW170817 ($Y_e < 0.25$)

Scenario for GW170817



blue kilonova
with $10^{-2}M_{\text{sun}}$

presence of red/purple kilonova
absence of energy injection by NS

Margalit &
Metzger 2017

III.

**How are the heavy elements
produced?**

Post-merger accretion disks: MHD turbulence





Fig.: **disk properties**; contours: optical depth for electron neutrinos

Fig.: **magnetic fields in the disk**; contours: rest-mass density

Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a

properties very similar to
Ciolfi, Siegel+ 2017, Kiuchi+ 2015

Post-merger accretion disks: MHD turbulence

average radially for space-time diagram





Fig.: **disk properties**; contours: optical depth for electron neutrinos


Fig.: **magnetic fields in the disk**; contours: rest-mass density

Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a


properties very similar to
Ciolfi, Siegel+ 2017, Kiuchi+ 2015

Accretion disk dynamo: butterfly diagram




magnetic energy is generated in the mid-plane

- migrates to higher latitudes
- dissipates into heat off the mid-plane



→ “hot corona”

hot corona launches thermal outflows (neutron-rich wind)




NS post-merger accretion disk are cooled from the mid-plane by neutrinos (rather than from the EM photosphere)!

Siegel & Metzger 2018a


Disk outflow properties

Siegel & Metzger 2018a



ejecta composition

$$Y_e \approx 0.1 - 0.3$$



ejecta velocities


$$v_\infty \approx 0.1c$$

corresponds to $\sim 8\text{MeV}$ per baryon in **nuclear binding energy release**


mass ejection due to **combination of thermal winds** from hot corona plus **nuclear binding energy release**

Why are the disk outflows neutron-rich?


t = 40ms



t = 130ms




t = 250ms



Siegel & Metzger 2018a


Neutron-rich conditions favor:



How can the overall Y_e of the outflow stay low ($\sim 0.1-0.2$)?

(and produce 3rd peak r-process elements?)

Self-regulation: keeping a neutron-rich reservoir





Neutrino-cooled accretion disks self-regulate themselves to mild degeneracy (low Y_e matter):



Beloborodov 2003, Chen & Beloborodov 2007, Metzger+ 2009

- viscous heating via magnetic turbulence
- neutrino cooling

charged-current processes:



pair annihilation:



plasmon decay:






Fig.: disk properties; contours: rest-mass density

Self-regulation: keeping a neutron-rich reservoir



Neutrino-cooled accretion disks self-regulate themselves to mild degeneracy (low Y_e matter):

Beloborodov 2003, Chen & Beloborodov 2007, Metzger+ 2009

- viscous heating via magnetic turbulence
- neutrino cooling

→ balance with feedback mechanism:

higher degeneracy μ_e/kT



fewer e^- , e^+ (lower Y_e)



less neutrino emission, i.e., cooling



higher temperatures



lower degeneracy μ_e/kT

direct evidence of self-regulation

Fig.: disk properties; contours: rest-mass density


Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a

r-process nucleosynthesis in disk outflows

nuclear reaction
network
(SkyNet)

- neutron captures
- photo-dissociations
- α -, β -decays
- fission





Movie: r-process nucleosynthesis from NS merger remnant disks

r-process nucleosynthesis in disk outflows

Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a



- robust 2nd and 3rd peak r-process!
- including neutrino absorption: additional good fit to 1st & 2nd peak elements




production of all r-process elements!


IV.

**constraints on r-process
enrichment**

Constraints on r-process nucleosynthesis




Constraints on r-process nucleosynthesis



post-merger disk outflows are a promising site for the r-process!




What about galactic chemical evolution?



late-time Eu/Fe decrease appears to be in tension
with NS merger paradigm Côté+ 2017, 2018, Hotokezaka+ 2018a

Conclusions

- ▶ Multiple kilonova components generally expected due to various ejecta mechanisms in NS mergers
- ▶ GW170817 with its exquisite kilonova data allowed us to address the **origin of the heavy elements**, which has been an **enduring mystery** for more than 70 years
- ▶ **First-principle simulations key** to understand kilonova properties and heavy-element formation (identify the site, production processes, abundance pattern etc.)
- ▶ **Simulations + GW170817 + EM (kilonova)** point to **post-merger accretion disk winds** as promising site (**ubiquitous phenomenon!**)



Relative abundances, total ejecta mass, measured BNS merger rate provide
yet strongest evidence for NS mergers being the prime production site for the r-process

Appendix

Disk simulations: numerical setup




Fig.: **disk properties**; contours: optical depth for electron neutrinos

Siegel & Metzger 2017, PRL

Siegel & Metzger 2018a

First self-consistent simulations modeling r-process nucleosynthesis from disk outflows from first principles:

- **GRMHD**: magnetic instabilities (**MRI**) mediating turbulence (transport of angular momentum) in the disk (*Einstein Toolkit*, *GRHydro*; Loeffler+ 2012, Moesta+ 2014)
- **weak interactions** in GRMHD
- new **enhanced methods for primitive recovery** to support **evolved microphysics** Siegel+ 2018
Siegel & Moesta 2018
- **approximate neutrino transport** (leakage scheme)
- **realistic EOS** (Helmholtz EOS) valid at low temperatures and densities, capturing nuclear binding energy release from **alpha-particle formation**
- **full r-process network calculations** on disk outflows using 10^4 tracer particles (*SkyNet*; Lippuner & Roberts 2017)

Previous 2D Newtonian alpha-disk simulations:

Fernandez & Metzger 2013, 2015, 2017

Just+ 2015