



Contribution ID: 382

Type: Poster

## CUPID-Mo, first sensitivity estimates for $2\nu\beta\beta(0\nu\beta\beta)$ decay of $^{100}\text{Mo}$ to excited states

CUPID-Mo is a demonstrator experiment for CUPID, a proposed next generation ton-scale  $0\nu\beta\beta$  search. CUPID-Mo, located in Laboratoire Souterrain de Modane, consists of an array of 20  $\sim 0.2\text{kg}$  enriched  $\text{Li}_2^{100}\text{MoO}_4$  scintillating cryogenic calorimeters with 20 Ge wafer light detectors for alpha and beta/gamma particle discrimination. Whilst CUPID-Mo aims to have leading sensitivity for  $0\nu\beta\beta$  to the ground state of  $\text{Ru1}$ , there is also the potential to measure  $2(0)\nu\beta\beta$  to excited states by looking at coincidences with gamma de-excitations. An excellent energy resolution of 2 - 7 keV over the full spectral range, along with good radiopurity leads us to expect a significant potential for new results. Here, we present a study of the signal efficiency and the predicted background index for various event signatures using a detailed Geant4 MC simulation.

### Mini-abstract

CUPID-Mo sensitivity to double beta decay of  $^{100}\text{Mo}$  to excited states

### Experiment/Collaboration

CUPID-Mo

**Primary author:** Mr DIXON, Toby (UC Berkeley)

**Presenter:** Mr DIXON, Toby (UC Berkeley)

**Session Classification:** Poster session 4