

# MPD exclusive channels

Chris Marshall

Lawrence Berkeley National Laboratory


21 October, 2019



# Needs for MPD physics studies

- Why do we need the HPgTPC/MPD and not just a cheap muon spectrometer?
- Demonstrate some physics channel that
  - Impacts LBL oscillation sensitivity/bias
  - Can't be determined by LAr alone
  - Is easily demonstrated with HPgTPC samples
- One idea we have discussed: pion multiplicity


# CC0 $\pi$ energy smearing



- LAr ND pseudo-reconstructed “FD TDR sample”
- Hadronic energy is determined by summing visible energy
- Cut on energy near the edges ensures reasonable containment


# CC2 $\pi$ energy smearing

- For events with 2 pions, energy is underestimated on average, by about 2 pion masses




# CC3 $\pi$ energy smearing

- For events with 3 pion, energy is underestimated on average, by about 3 pion masses





# CC1 $\pi$ energy smearing

- For events with 1 pion, energy is underestimated on average, by about 1 pion mass




# $CC1\pi^\pm$ vs. $CC1\pi^0$



- Charge of the pion has a significant impact on LAr energy reconstruction – we typically see the pion mass when it is neutral

# CC acceptance in LAr

- Muon must be contained or HPgTPC matched, hence the dip
- Hadronic energy must be contained  $\rightarrow$  better efficiency when the same energy is distributed among nucleons instead of mesons





# Potential for LBL bias

- If the mix of  $0\pi/1\pi/2\pi$  in our model is incorrect, then our model-based energy unsmearing will be wrong, which could lead to biases
  - This could be demonstrated by creating a mock data sample by re-weighting based on pion multiplicity and/or kinematics, and fitting
- If there is any dependence on neutrino energy, then the LAr ND cannot resolve this bias for FD because the fluxes are different
- LAr ND may be able to address this by selecting exclusive final state samples, but this could be challenging due to thresholds, hadronic showers, etc.
- If HPgTPC efficiency is  $\sim$ flat vs. number of pions, neutrino energy, etc. then these acceptance curves could be directly measured

# To do

- Make all these plots for HPgTPC samples
- Demonstrate potential LBL bias by constructing a mock data sample
  - What will probably happen is FD-only will give good fit with wrong oscillation parameters
  - FD + LAr ND will give terrible fit with nuisance parameters pulled all over the place
  - Show that HPgTPC sample allows you to identify *how* the model is wrong, potentially fix it

# Other ideas?

• ...