



Contribution ID: 86

Type: **not specified**

## Detecting Dark Matter with a Qubit

*Monday, July 18, 2022 8:40 PM (20 minutes)*

To perform experimental searches for low mass bosonic dark matter such as the hidden photon or axion, our group works to employ dielectric photonic bandgap cavities with a high quality factor to coherently accumulate the axion signal for readout using qubit-based single photon detectors. The advantage of the qubit-based detector is in overcoming the standard quantum noise limit through repeated quantum non-demolition measurements [1]. Other techniques being studied include preparing the dielectric cavity in a higher photon-number ( $n$ ) Fock state to enhance the dark matter signal amplitude by a factor of  $(n+1)$ . Given the large parameter space still unexplored by current dark matter searches, methods of tuning a resonant cavity by electronically controlling the magnetic field seen by a loop of two Josephson junctions is currently being studied.

[1] Dixit et al., Phys. Rev. Lett. 126, 141302 (2021)

### **In-person or Virtual?**

In-person

**Primary authors:** Dr DIXIT, Akash (University of Chicago); AGRAWAL, Ankur (University of Chicago); ZHAO, Fang (Fermi National Accelerator Laboratory); LYNN, Morgan (University of Chicago/Fermilab)

**Co-authors:** CHOU, Aaron (Fermi National Accelerator Laboratory); Prof. SCHUSTER, David (University of Chicago)

**Presenter:** LYNN, Morgan (University of Chicago/Fermilab)

**Session Classification:** Poster Session