

Theory Overview of Charged Lepton Flavor Violation in Heavy Particle Decays

Wolfgang Altmannshofer (UCSC)

waltmann@ucsc.edu



mainly based on Whitepaper 2205.10576 with:

Cecile Caillol (CERN), Mogens Dam (NBI), Stefania Xella (NBI),
Yongchao Zhang (Nanjing)

Snowmass Summer Meeting 2022, Seattle, July 17 - 26, 2022

Heavy State CLFV Decays and New Physics

- In the SM, lepton flavor violating decays of the Z , Higgs, and top are suppressed by the tiny neutrino mass splittings

$$\text{e.g. } \text{BR}(Z \rightarrow \mu e) \sim \text{BR}(Z \rightarrow \mu\mu) \left| \frac{g^2}{16\pi^2} \frac{m_\nu^2}{m_W^2} \right|^2 \sim 10^{-50}$$

- Any observation in the foreseeable future would be an unambiguous sign of new physics.

Comparison with Low Energy Probes

- Consider LFV decays of the Z boson, the Higgs, the top in the presence of generic New Physics

$$\frac{\text{BR}(Z \rightarrow \mu e)}{\text{BR}(Z \rightarrow \mu\mu)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4, \quad \frac{\text{BR}(H \rightarrow \tau\mu)}{\text{BR}(H \rightarrow \tau\tau)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4$$

$$\frac{\text{BR}(t \rightarrow c\mu e)}{\text{BR}(t \rightarrow Wb)} \sim \frac{g_{\text{NP}}^2}{16\pi^2} \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4$$

Comparison with Low Energy Probes

- Consider LFV decays of the Z boson, the Higgs, the top in the presence of generic New Physics

$$\frac{\text{BR}(Z \rightarrow \mu e)}{\text{BR}(Z \rightarrow \mu \mu)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4, \quad \frac{\text{BR}(H \rightarrow \tau \mu)}{\text{BR}(H \rightarrow \tau \tau)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4$$

$$\frac{\text{BR}(t \rightarrow c \mu e)}{\text{BR}(t \rightarrow W b)} \sim \frac{g_{\text{NP}}^2}{16\pi^2} \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4$$

- Compare to low energy probes (e.g. muon decays, tau decays)

$$\frac{\text{BR}(\mu \rightarrow 3e)}{\text{BR}(\mu \rightarrow e \nu_\mu \bar{\nu}_e)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4$$

- Same dependence on NP couplings and scale, but much less Z, Higgs, top available in experiments
- Note: these are extremely generic/naive expectations; situation can be very different in concrete models.

Framing the Discussion Model Independently: SMEFT

1 : X^3	2 : H^6	3 : $H^4 D^2$	5 : $\psi^2 H^3 + \text{h.c.}$
Q_G $f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H $(H^\dagger H)^3$	$Q_{H\Box}$ $(H^\dagger H)\Box(H^\dagger H)$	Q_{eH} $(H^\dagger H)(\bar{l}_p e_r H)$
$Q_{\tilde{G}}$ $f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$		Q_{HD} $(H^\dagger D_\mu H)^* (H^\dagger D_\mu H)$	Q_{uH} $(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W $\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			Q_{dH} $(H^\dagger H)(\bar{q}_p d_r H)$
$Q_{\tilde{W}}$ $\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			

4 : $X^2 H^2$	6 : $\psi^2 XH + \text{h.c.}$	7 : $\psi^2 H^2 D$
Q_{HG} $H^\dagger H G_\mu^A G^{A\mu\nu}$	Q_{eW} $(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{Hl}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
$Q_{H\tilde{G}}$ $H^\dagger H \tilde{G}_\mu^A G^{A\mu\nu}$	Q_{eB} $(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{Hl}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
Q_{HW} $H^\dagger H W_\mu^I W^{I\mu\nu}$	Q_{uG} $(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{He} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$Q_{H\tilde{W}}$ $H^\dagger H \tilde{W}_\mu^I W^{I\mu\nu}$	Q_{uW} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
Q_{HB} $H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{H\tilde{B}}$ $H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG} $(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
Q_{HWB} $H^\dagger \tau^I H W_\mu^I B^{\mu\nu}$	Q_{dW} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$Q_{H\tilde{W}B}$ $H^\dagger \tau^I H \tilde{W}_\mu^I B^{\mu\nu}$	Q_{dB} $(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$ $i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$

8 : $(\bar{L}L)(\bar{L}L)$	8 : $(\bar{R}R)(\bar{R}R)$	8 : $(\bar{L}L)(\bar{R}R)$
Q_{ll} $(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee} $(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le} $(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu} $(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu} $(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qq}^{(3)}$ $(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd} $(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld} $(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$ $(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu} $(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe} $(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$ $(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed} $(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
	$Q_{ud}^{(1)}$ $(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
	$Q_{ud}^{(8)}$ $(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
		$Q_{qd}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$

8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$	8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$
Q_{ledq} $(\bar{l}_p^j e_r)(\bar{d}_s q_{tj})$	$Q_{quqd}^{(1)}$ $(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k d_t)$
	$Q_{quqd}^{(8)}$ $(\bar{q}_p^j T^A u_r) \epsilon_{jk} (\bar{q}_s^k T^A d_t)$
	$Q_{lequ}^{(1)}$ $(\bar{l}_p^j e_r) \epsilon_{jk} (\bar{q}_s^k u_t)$
	$Q_{lequ}^{(3)}$ $(\bar{l}_p^j \sigma_{\mu\nu} e_r) \epsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$

2499 baryon number conserving
dim. 6 operators in total
Grzadkowski et al. 1008.4884

Framing the Discussion Model Independently: SMEFT

1 : X^3	2 : H^6	3 : $H^4 D^2$	5 : $\psi^2 H^3 + \text{h.c.}$
Q_G $f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H $(H^\dagger H)^3$	$Q_{H\Box}$ $(H^\dagger H)\Box(H^\dagger H)$	Q_{eH} $(H^\dagger H)(\bar{l}_p e_r H)$
$Q_{\tilde{G}}$ $f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$		Q_{HD} $(H^\dagger D_\mu H)^* (H^\dagger D_\mu H)$	Q_{uH} $(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W $\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			Q_{dH} $(H^\dagger H)(\bar{q}_p d_r H)$
$Q_{\tilde{W}}$ $\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			

4 : $X^2 H^2$	6 : $\psi^2 X H + \text{h.c.}$	7 : $\psi^3 H^3 D$
Q_{HG} $H^\dagger H G_\mu^A G^{A\mu\nu}$	Q_{eW} $(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{H1}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
$Q_{H\tilde{G}}$ $H^\dagger H \tilde{G}_\mu^A G^{A\mu\nu}$	Q_{uB} $(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{H1}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
Q_{HW} $H^\dagger H W_\mu^I W^{I\mu\nu}$	Q_{uG} $(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{H2} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$Q_{H\tilde{W}}$ $H^\dagger H \tilde{W}_\mu^I W^{I\mu\nu}$	Q_{uW} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
Q_{HB} $H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{H\tilde{B}}$ $H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{BG} $(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
Q_{HWB} $H^\dagger \tau^I H W_\mu^I B^{\mu\nu}$	Q_{dW} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$Q_{H\tilde{W}B}$ $H^\dagger \tau^I H \tilde{W}_\mu^I B^{\mu\nu}$	Q_{dB} $(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$ $i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$

2499 baryon number conserving
dim. 6 operators in total
Grzadkowski et al. 1008.4884

4 fermion interactions

8 : $(\bar{L}L)(\bar{L}L)$	8 : $(\bar{R}R)(\bar{R}R)$	8 : $(\bar{L}L)(\bar{R}R)$
Q_{ll} $(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee} $(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le} $(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qe}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu} $(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu} $(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qe}^{(3)}$ $(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd} $(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld} $(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$ $(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu} $(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe} $(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$ $(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed} $(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
	$Q_{ud}^{(1)}$ $(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
	$Q_{ud}^{(8)}$ $(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
		$Q_{qd}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$
8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$	8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$	
$Q_{le dq}$ $(\bar{l}_p^j e_r)(\bar{d}_s^k q_{tj})$	$Q_{qq\Box}^{(1)}$ $(\bar{q}_p^j u_r) e_{jk} (\bar{q}_s^k d_t)$	
	$Q_{qq\Box}^{(8)}$ $(\bar{q}_p^j T^A u_r) e_{jk} (\bar{q}_s^k T^A d_t)$	
	$Q_{le qu}^{(1)}$ $(\bar{l}_p^j e_r) e_{jk} (\bar{q}_s^k u_t)$	
	$Q_{le qu}^{(8)}$ $(\bar{l}_p^j \sigma_{\mu\nu} e_r) e_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$	

Framing the Discussion Model Independently: SMEFT

1 : X^3	2 : H^6	3 : $H^4 D^2$	5 : $\psi^2 H^3 + \text{h.c.}$
Q_G $f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H $(H^\dagger H)^3$	$Q_{H\Box}$ $(H^\dagger H)\Box(H^\dagger H)$	Q_{eH} $(H^\dagger H)(\bar{l}_p e_r \tilde{H})$
$Q_{\tilde{G}}$ $f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$		Q_{HD} $(H^\dagger D_\mu H)^\dagger (H^\dagger D_\mu H)$	Q_{uH} $(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W $\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			Q_{dH} $(H^\dagger H)(\bar{q}_p d_r \tilde{H})$
$Q_{\tilde{W}}$ $\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			

4 : $X^2 H^2$	6 : $\psi^2 X H + \text{h.c.}$	7 : $\psi^3 H^3 D$
Q_{HG} $H^\dagger H G_\mu^A G^{A\mu\nu}$	Q_{eW} $(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{H1}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
$Q_{H\tilde{G}}$ $H^\dagger H \tilde{G}_\mu^A G^{A\mu\nu}$	Q_{uB} $(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{H1}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
Q_{HW} $H^\dagger H W_\mu^I W^{I\mu\nu}$	Q_{uG} $(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{H2} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$Q_{H\tilde{W}}$ $H^\dagger H \tilde{W}_\mu^I W^{I\mu\nu}$	Q_{uW} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
Q_{HB} $H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{H\tilde{B}}$ $H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{BG} $(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
Q_{HWB} $H^\dagger \tau^I H W_\mu^I B^{\mu\nu}$	Q_{dW} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$Q_{H\tilde{W}B}$ $H^\dagger \tau^I H \tilde{W}_\mu^I B^{\mu\nu}$	Q_{dB} $(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$ $i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$

8 : $(\bar{L}L)(\bar{L}L)$	8 : $(\bar{R}R)(\bar{R}R)$	8 : $(\bar{L}L)(\bar{R}R)$
Q_l $(\bar{l}_p \gamma_\mu l_r)(\bar{l}_r \gamma^\mu l_e)$	Q_{ee} $(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le} $(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qe}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu} $(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu} $(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qe}^{(3)}$ $(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd} $(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld} $(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$ $(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu} $(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe} $(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$ $(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed} $(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
	$Q_{ud}^{(1)}$ $(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(3)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
	$Q_{ud}^{(8)}$ $(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
		$Q_{qd}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$

8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$	8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$
$Q_{le dq}$ $(\bar{l}_p^j e_r)(\bar{d}_s q_{tj})$	$Q_{qq qd}^{(1)}$ $(\bar{q}_p^j u_r)(\bar{q}_s^k d_{tj})$
	$Q_{qq qd}^{(8)}$ $(\bar{q}_p^j T^A u_r)(\bar{q}_s^k d_{tj})$
	$Q_{le qu}^{(1)}$ $(\bar{l}_p^j e_r)(\bar{q}_s^k u_{tj})$
	$Q_{le qu}^{(3)}$ $(\bar{l}_p^j \sigma_{\mu\nu} e_r)(\bar{q}_s^k \sigma^{\mu\nu} u_{tj})$

2499 baryon number conserving
dim. 6 operators in total

Grzadkowski et al. 1008.4884

4 fermion interactions

dipole transitions

Framing the Discussion Model Independently: SMEFT

1 : X^3	2 : H^6	3 : $H^4 D^2$	5 : $\psi^2 H^3 + \text{h.c.}$
Q_G $f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H $(H^\dagger H)^3$	$Q_{H\Box}$ $(H^\dagger H)\Box(H^\dagger H)$	Q_{eH} $(H^\dagger H)(\bar{l}_p e_r \tilde{H})$
$Q_{\tilde{G}}$ $f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$		Q_{HD} $(H^\dagger D_\mu H)^\dagger (H^\dagger D_\mu H)$	Q_{uH} $(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W $\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			Q_{dH} $(H^\dagger H)(\bar{q}_p d_r \tilde{H})$
$Q_{\tilde{W}}$ $\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			

4 : $X^2 H^2$	6 : $\psi^2 X H + \text{h.c.}$	7 : $\psi^3 H^2 D$
Q_{HC} $H^\dagger H G_\mu^{A\nu} G_\nu^{A\mu}$	Q_{eW} $(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{H1}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
Q_{HG} $H^\dagger H \tilde{G}_\mu^{A\nu} G_\nu^{A\mu}$	Q_{uB} $(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{H1}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
Q_{HW} $H^\dagger H W_\mu^{I\nu} W_\nu^{I\mu}$	Q_{uG} $(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{H2} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$Q_{H\tilde{W}}$ $H^\dagger H \tilde{W}_\mu^{I\nu} W_\nu^{I\mu}$	Q_{uW} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
Q_{HB} $H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{H\tilde{B}}$ $H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{BG} $(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \tilde{H} G_{\mu\nu}^A$	Q_{Hu} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
Q_{HWB} $H^\dagger \tau^I H W_\mu^{I\nu} B_\nu^{\mu}$	Q_{dW} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \tilde{H} W_{\mu\nu}^I$	Q_{Hd} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$Q_{H\tilde{W}B}$ $H^\dagger \tau^I H \tilde{W}_\mu^{I\nu} B_\nu^{\mu}$	Q_{dB} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tilde{H} B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$ $i(\tilde{H}^\dagger D_\mu \tilde{H})(\bar{u}_p \gamma^\mu d_r)$

8 : $(\bar{L}L)(\bar{L}L)$	8 : $(\bar{R}R)(\bar{R}R)$	8 : $(\bar{L}L)(\bar{R}R)$
Q_{ll} $(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee} $(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le} $(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qe}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu} $(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu} $(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qe}^{(3)}$ $(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd} $(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld} $(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$ $(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu} $(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe} $(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$ $(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed} $(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
	$Q_{ud}^{(1)}$ $(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
	$Q_{ud}^{(8)}$ $(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
		$Q_{qd}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$

8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$	8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$
$Q_{le dq}$ $(\bar{l}_p^j e_r)(\bar{d}_s^k q_t^j)$	$Q_{qq uq}^{(1)}$ $(\bar{q}_p^j u_r)(\bar{q}_s^k d_t^j)$
	$Q_{qq uq}^{(8)}$ $(\bar{q}_p^j T^A u_r)(\bar{q}_s^k d_t^j)$
	$Q_{le qu}^{(1)}$ $(\bar{l}_p^j e_r)(\bar{q}_s^k u_t^j)$
	$Q_{le qu}^{(3)}$ $(\bar{l}_p^j \sigma_{\mu\nu} e_r)(\bar{q}_s^k \sigma^{\mu\nu} u_t^j)$

2499 baryon number conserving
dim. 6 operators in total
Grzadkowski et al. 1008.4884

4 fermion interactions

dipole transitions

"Z-penguins"

Framing the Discussion Model Independently: SMEFT

1 : X^3	2 : H^6	3 : $H^4 D^2$	5 : $\psi^2 H^3 + \text{h.c.}$
Q_G $f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_H $(H^\dagger H)^3$	$Q_{H\Box}$ $(H^\dagger H)\Box(H^\dagger H)$	Q_{eH} $(H^\dagger H)(\bar{l}_p e_r H)$
$Q_{\tilde{G}}$ $f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$		Q_{HD} $(H^\dagger D_\mu H)^* (H^\dagger D_\mu H)$	Q_{uH} $(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
Q_W $\epsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			Q_{dH} $(H^\dagger H)(\bar{q}_p d_r H)$
$Q_{\tilde{W}}$ $\epsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$			

4 : $X^2 H^2$	6 : $\psi^2 X H + \text{h.c.}$	7 : $\psi^3 H^3 D$
Q_{HC} $H^\dagger H G_\mu^A G^{A\mu\nu}$	Q_{eW} $(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$Q_{H1}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
Q_{HG} $H^\dagger H \tilde{G}_\mu^A G^{A\mu\nu}$	Q_{uB} $(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$Q_{H1}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
Q_{HW} $H^\dagger H W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG} $(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	Q_{H2} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$Q_{H\tilde{W}}$ $H^\dagger H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$Q_{Hq}^{(1)}$ $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
Q_{HB} $H^\dagger H B_{\mu\nu} B^{\mu\nu}$	Q_{uB} $(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$Q_{Hq}^{(3)}$ $(H^\dagger i \overleftrightarrow{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{H\tilde{B}}$ $H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{BG} $(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	Q_{Hu} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
Q_{HWB} $H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW} $(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	Q_{Hd} $(H^\dagger i \overleftrightarrow{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$Q_{H\tilde{W}B}$ $H^\dagger \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB} $(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	$Q_{Hud} + \text{h.c.}$ $i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$

8 : $(\bar{L}L)(\bar{L}L)$	8 : $(\bar{R}R)(\bar{R}R)$	8 : $(\bar{L}L)(\bar{R}R)$
Q_{ll} $(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee} $(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le} $(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qe}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu} $(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu} $(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qe}^{(3)}$ $(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd} $(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld} $(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$ $(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu} $(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe} $(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$ $(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed} $(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
	$Q_{ud}^{(1)}$ $(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
	$Q_{ud}^{(8)}$ $(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$ $(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
		$Q_{qd}^{(8)}$ $(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$

8 : $(\bar{L}R)(\bar{R}L) + \text{h.c.}$	8 : $(\bar{L}R)(\bar{L}R) + \text{h.c.}$
$Q_{le dq}$ $(\bar{l}_p^j e_r)(\bar{d}_s^k q_t^j)$	$Q_{qq uq}^{(1)}$ $(\bar{q}_p^j u_r)(\bar{q}_s^k d_t^j)$
	$Q_{qq uq}^{(8)}$ $(\bar{q}_p^j T^A u_r)(\bar{q}_s^k d_t^j)$
	$Q_{le qu}^{(1)}$ $(\bar{l}_p^j e_r)(\bar{q}_s^k u_t^j)$
	$Q_{le qu}^{(3)}$ $(\bar{l}_p^j \sigma_{\mu\nu} e_r)(\bar{q}_s^k \sigma^{\mu\nu} u_t^j)$

2499 baryon number conserving
dim. 6 operators in total
Grzadkowski et al. 1008.4884

4 fermion interactions

dipole transitions

"Z-penguins"

"Higgs penguins"

LFV Z Decays

Existing/Expected Bounds

- Results from the LHC: ATLAS (139 fb^{-1})

Phys.Rev.Lett. 127 (2022) 271801; Nature Phys. 17 (2021) 7, 819-825; ATLAS-CONF-2021-042

$$\text{BR}(Z \rightarrow \mu e) < 3.04 \times 10^{-7}$$

$$\text{BR}(Z \rightarrow \tau e) < 5.0 \times 10^{-6}$$

$$\text{BR}(Z \rightarrow \tau \mu) < 6.5 \times 10^{-6}$$

- Better than LEP for all decay modes.
- In all searches there are backgrounds \Rightarrow expect sensitivities to improve with $\sqrt{\mathcal{L}}$, i.e. \sim factor of 5 at the HL-LHC.
- At Tera-Z factories expect $\text{BR}(Z \rightarrow \mu e) \sim 10^{-8} - 10^{-10}$ and $\text{BR}(Z \rightarrow \tau \ell) \sim 10^{-9}$ (Dam 1811.09408)

- Z couplings are protected by $SU(2)$ gauge symmetry

⇒ generic expectation for a new physics effect

$$\frac{\text{BR}(Z \rightarrow \ell\ell')}{\text{BR}(Z \rightarrow \ell\ell)} \sim g_{\text{NP}}^2 \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4 \sim 4 \times 10^{-7} \times g_{\text{NP}}^2 \left(\frac{10 \text{ TeV}}{\Lambda_{\text{NP}}} \right)^4$$

⇒ sensitivity to New Physics at scales of

$$\Lambda_{\text{NP}} \sim 10 \text{ TeV at the HL-LHC}$$

$$\Lambda_{\text{NP}} \sim 50 \text{ TeV at FCC-ee/CEPC}$$


LFV Z Decays in the EFT Framework

- Parameterize New Physics in a systematic and controlled way:
in terms of dim-6 operators of the SMEFT

dipoles

$$\mathcal{O}_{dW} = (\bar{\ell} \sigma^{\mu\nu} \tau^a P_R \ell') H W_{\mu\nu}^a$$

$$\mathcal{O}_{dB} = (\bar{\ell} \sigma^{\mu\nu} P_R \ell') H B_{\mu\nu}$$




“Z
penguins”

$$\mathcal{O}_{hl}^{(3)} = (H^\dagger i \overleftrightarrow{D}_\mu^a H) (\bar{\ell} \gamma^\mu \tau^a P_L \ell')$$


$$\tilde{\mathcal{O}}_{hl}^{(1)} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{\ell} \gamma^\mu P_L \ell')$$

$$\mathcal{O}_{he} = (H^\dagger i \overleftrightarrow{D}_\mu H) (\bar{\ell} \gamma^\mu P_R \ell')$$




Complementarity with Low Energy Probes

- ▶ Many flavor violating **low energy processes** will be affected as well.
- ▶ Severe indirect constraints on $Z \rightarrow \mu e$ from $\mu \rightarrow e\gamma$, $\mu \rightarrow 3e$, $\mu \rightarrow e$ conversion (barring accidental cancellations).



Complementarity with Low Energy Probes

- **Complementary** sensitivity in the case of taus.



LFV Higgs Decays

► Results from the LHC

ATLAS 1907.06131 ($\sim 36 \text{ fb}^{-1}$), ATLAS 1909.10235 ($\sim 139 \text{ fb}^{-1}$),
CMS 2105.03007 ($\sim 137 \text{ fb}^{-1}$)

$$\text{BR}(H \rightarrow \mu e) < 6.1 \times 10^{-5}$$

$$\text{BR}(H \rightarrow \tau e) < 0.22\%$$

$$\text{BR}(H \rightarrow \tau \mu) < 0.15\%$$

- Expect sensitivities to **improve by ~ 1 order of mag.** at the HL-LHC
- Expect sensitivities at future e^+e^- colliders that are at least as good
(Qin et al. 1711.07243)

The Higgs and Flavor

$$\mathcal{L}_{\text{Yukawa}} = \lambda_{ij} \bar{\Psi}_i \Psi_j H$$

In the **Standard Model** the Yukawa couplings are the only sources of flavor and CP violation

→ the couplings of the Higgs to fermion mass eigenstates are **flavor diagonal and CP conserving**

$$\frac{1}{v} \begin{pmatrix} m_{u,d,e} & 0 & 0 \\ 0 & m_{c,s,\mu} & 0 \\ 0 & 0 & m_{t,b,\tau} \end{pmatrix}$$

The Higgs and Flavor

$$\mathcal{L}_{\text{Yukawa}} = \lambda_{ij} \bar{\Psi}_i \Psi_j H + \frac{\tilde{\lambda}_{ij}}{\Lambda^2} \bar{\Psi}_i \Psi_j H^3$$

In the **Standard Model** the Yukawa couplings are the only sources of flavor and CP violation

→ the couplings of the Higgs to fermion mass eigenstates are **flavor diagonal and CP conserving**



$$\frac{1}{v} \begin{pmatrix} m_{u,d,e} & 0 & 0 \\ 0 & m_{c,s,\mu} & 0 \\ 0 & 0 & m_{t,b,\tau} \end{pmatrix} + \frac{v^2}{\Lambda^2} \begin{pmatrix} \star & \star & \star \\ \star & \star & \star \\ \star & \star & \star \end{pmatrix}$$

- 1) **New Physics** can modify the **flavor diagonal** Higgs couplings
- 2) **New Physics** can lead to **flavor and CP violating** Higgs couplings

Phenomenological parameterization: $\mathcal{L}_{\text{CLFV}} = -Y_{\ell\ell'} \bar{\ell} P_{R\ell'} h + \text{h.c.}$

Bounds on Flavor Violating Higgs Couplings

WA, Caillol, Dam, Xella, Zhang 2205.10576




- Weak indirect constraints from $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$.
- $\mu \rightarrow e\gamma$ strongly constrains $\text{BR}(H \rightarrow \mu e)$ and $\text{BR}(H \rightarrow \tau\mu) \times \text{BR}(H \rightarrow \tau e)$

Blankenburg, Ellis, Isidori 1107.1216; Harnik, Kopp, Zupan 1209.1397; Davidson, Verdie 1211.1248

LFV Higgs Decays in the EFT Framework

$$\frac{C_{\ell\ell'}}{\Lambda_{\text{NP}}^2} \mathcal{O}_{eh} = \frac{C_{\ell\ell'}}{\Lambda_{\text{NP}}^2} (H^\dagger H) (\bar{\ell} P_R \ell') H$$




- Gives flavor changing Higgs couplings

$$Y_{\ell\ell'} = \frac{C_{\ell\ell'}}{\sqrt{2}} \frac{v^2}{\Lambda_{\text{NP}}^2} \sim 4 \times 10^{-4} \left(\frac{10 \text{ TeV}}{\Lambda_{\text{NP}}} \right)^2$$

- Expected sensitivities at future machines probe **new physics at $\Lambda_{\text{NP}} \sim 10 \text{ TeV}$.**

LFV Higgs Decays in the EFT Framework

$$\frac{C_{\ell\ell'}}{\Lambda_{\text{NP}}^2} \mathcal{O}_{eh} = \frac{C_{\ell\ell'}}{\Lambda_{\text{NP}}^2} (H^\dagger H) (\bar{\ell} P_R \ell') H$$



- Gives flavor changing Higgs couplings


$$Y_{\ell\ell'} = \frac{C_{\ell\ell'}}{\sqrt{2}} \frac{v^2}{\Lambda_{\text{NP}}^2} \sim 4 \times 10^{-4} \left(\frac{10 \text{ TeV}}{\Lambda_{\text{NP}}} \right)^2$$

- Expected sensitivities at future machines probe **new physics at $\Lambda_{\text{NP}} \sim 10 \text{ TeV}$.**
- Comment: need to be very careful when calculating loops outside the EFT framework. Results might be gauge dependent.

(WA, Gori, Hamer, Patel arXiv:2009.01258)

LFV Higgs Decays in NP Models


In new physics models one often encounters strong constraints:
The physics that generates the LFV Higgs coupling, will typically also
give **direct contributions to radiative decays** (Dorsner et al. 1502.07784)



Contributions to lepton Yukawa couplings (a) , electromagnetic dipole (b)

LFV Higgs Decays in NP Models

In new physics models one often encounters strong constraints:
The physics that generates the LFV Higgs coupling, will typically also give **direct contributions to radiative decays** (Dorsner et al. 1502.07784)



Contributions to lepton Yukawa couplings (a) , electromagnetic dipole (b)

handwavy upper bound in many models

(assuming that the Wilson coefficient of the dipole is $\frac{1}{16\pi^2} \times$ the Wilson coefficient of the Higgs penguin)

$$\text{BR}(h \rightarrow \tau\mu) \sim 26 \times \text{BR}(\tau \rightarrow \mu\gamma) \lesssim 10^{-6}$$

WA, Gori, Kagan, Silvestrini, Zupan 1507.07927

\Rightarrow Observation of a LFV Higgs decay with expected exp. sensitivities
likely implies an additional source of EW symmetry breaking

LFV Top Decays

EFT Setup and Sensitivity to New Physics


3 body decays that **violate lepton and quark flavor** $t \rightarrow q \ell \ell'$

(Davidson, Mangano, Perries, Sordini 1507.07163)

$$\mathcal{O}_{LL} = (\bar{q} \gamma_\mu P_L t) (\bar{\ell} \gamma^\mu P_L \ell')$$

$$\mathcal{O}_{RR} = (\bar{q} \gamma_\mu P_R t) (\bar{\ell} \gamma^\mu P_R \ell')$$

+ many other Dirac structures



The decays are competing with an unsuppressed 2 body decay $t \rightarrow Wb$

$$\text{BR}(t \rightarrow c \mu e) \sim \frac{g_{\text{NP}}^2}{16\pi^2} \left(\frac{v}{\Lambda_{\text{NP}}} \right)^4 \sim 2 \times 10^{-5} \times g_{\text{NP}}^2 \left(\frac{1 \text{ TeV}}{\Lambda_{\text{NP}}} \right)^4$$

- Strong indirect bounds from B meson decays if left handed quarks are involved.
- For right handed quarks, LHC has the best sensitivity.


Experimental Sensitivity

- ▶ Look for $t\bar{t}$ production followed by a rare top decay $t \rightarrow q\mu e$ and also for non-standard single top production $gq \rightarrow t\mu e$.
- ▶ Main background from $t\bar{t}$, which gives two b-jets
- ▶ Signal has only a single b-jet
- ▶ Translation into top branching ratio depends on the Dirac structure of the operator

$$\text{BR}(t \rightarrow u\mu e) \lesssim 10^{-7}$$

$$\text{BR}(t \rightarrow c\mu e) \lesssim 10^{-6}$$

- ▶ Expect factor of ~ 5 improvement at HL-LHC
- ▶ For further improvement need FCC-hh



LFV New Physics Resonances

- Many BSM scenarios contain neutral resonances that can have lepton flavor violating couplings

e.g. Z' bosons, or additional neutral Higgs bosons H .

- Obvious approach: extend the Z and Higgs searches to higher (and lower!) masses

$$pp \rightarrow Z' \rightarrow e\mu, e\tau, \mu\tau, \quad pp \rightarrow H \rightarrow e\mu, e\tau, \mu\tau$$

$$e^+e^- \rightarrow Z' \rightarrow e\mu, e\tau, \mu\tau, \quad e^+e^- \rightarrow Z + H \rightarrow Z + e\mu, e\tau, \mu\tau$$


- In contrast to standard high-mass di-lepton resonance searches, no irreducible background from Drell-Yan

Exotic Scenarios

- Can imagine exotic scenarios: e.g. a Z' that couples dominantly in a flavor violating way to $\tau\mu$ (can give a viable explanation of $(g-2)_\mu$)
- Currently weakly constrained, but could give **spectacular same sign lepton pair signatures** at lepton colliders

$$\text{e.g. } e^+e^- \rightarrow Z'\tau^+\mu^- \rightarrow \tau^+\tau^+\mu^-\mu^-$$

WA, Caillol, Dam, Xella, Zhang 2205.10576 (update of WA, Chen, Dev, Soni 1607.06832)






More Exotic Scenarios

- Similar results are found for additional Higgs boson that have only flavor violating couplings

$$\text{e.g. } e^+e^- \rightarrow H\mu^+e^- \rightarrow \mu^+\mu^+e^-e^-$$

Dev, Mohapatra, Zhang 1711.08430



- Model building challenge: construct a model in which these exotic Z' or Higgs bosons with only flavor violating couplings arise.

- ▶ Lepton flavor violating decays of Z , Higgs, top are clear signatures of NP.
- ▶ With the expected experimental sensitivities one can probe NP scales of 10 TeV or even higher.
- ▶ Often strong indirect constraints from low energy lepton flavor violating processes ($\mu \rightarrow e\gamma$ etc.), but in many cases there is complementary sensitivity to the NP.
- ▶ BSM resonances can also give prominent lepton flavor violating signatures. Worthwhile to extend the searches to a as broad mass range as possible.