

CF03: Cosmic Probes of Dark Matter Physics

SnowMass2021

Conveners:

Alex Drlica-Wagner

Chanda Prescod-Weinstein

Hai-Bo Yu

July 23, 2022

CF03: Cosmic Probes of Dark Matter


- Cosmological and astrophysical measurements provide the **only** robust, positive empirical measurements of dark matter.
- Cosmic probes are **unique** in that they do not rely on the assumption that dark matter has interactions with normal matter beyond gravity; thus they are the most “expansive” (and could be the **only** viable) approach to the dark matter problem.
- Cosmic probes is **an emergent field** that requires strong synergy among particle theorists, dynamists, simulators, observers, and experimentalists; need a **new mechanism** to support these emerging, collaborative efforts.
- Cosmic probes are highly relevant and complementary to search efforts in CF1, CF2, CF7 and other frontiers, and there is strong **experimental synergy** with cosmological probes of dark matter, dark energy, and inflation (CF4, CF5, CF6) **and theory frontier**

Halo Measurements




[2203.07354](#)

Extreme Environments




Particle-Like

Indirect Detection




Direct Detection




Wave-Like

Axion Coupling



Cosmic Probes

Self-Interaction




Gravitational Coupling




Particle-Like

Indirect Detection




Direct Detection




Wave-Like

Axion Coupling



Cosmic Probes

Self-Interaction



Accelerator Searches

Gravitational Coupling



Interaction Strength

Dark Matter Mass



Ultra-Light

Axion-Like Particles

QCD Axion

Self-Interacting

Dark Sectors

Thermal Relic

Compact
Objects

Dark Matter Mass



Interaction Strength

Ultra Light

Self-Interacting

Compact
Objects

Axion-Like Particles

Dark Sectors

QCD Axion

Thermal Relic

Cosmic Probes

Particle-Like

Wave-Like

Delve Deep, Search Wide!

Letters of Interest and Solicited White Papers

- CF03 received **~75 Letters of Interest** from the community.
- Through a series of discussions (including the Community Planning Meeting), we arrived at a list of **5 solicited white papers** with designated facilitators. All have been submitted.
- CF03 has received **5 additional white papers** (to date). Other relevant white papers include **~15 white papers** submitted to other CF topical groups and other frontiers.

THANK YOU
white paper facilitators and authors!

Authorship extended to white paper facilitators and major contributors.
Please contact us if you have contributed and want to be an author.

Cosmic Probes of Dark Matter

Conveners: A. Drlica-Wagner, C. Prescod-Weinstein, H.-B. Yu

Contributors: A. Albert, A. Banerjee, S. Birrer, R. Caputo, D. Croon, D. Gilman, Y.-Y. Mao,
E. O. Nadler, K. Perez, A. H. G. Peter, N. Sehgal, J. D. Simon, ...

(ADW: Authorship has been extended to white paper facilitators and major contributors. Please contact us if you feel that your contributions merit authorship.)

Draft of CF03 Report (v20220718): [link](#)

Comments and Feedback: [link](#)

Cosmic Probes of Dark Matter	1
3.1 Executive Summary	3
3.2 Introduction	5
3.3 Dark Matter Halo Measurements	7
3.4 Dark Matter Simulations	10
3.5 Primordial Black Holes, Dark Matter, and the Early Universe	14
3.6 Dark Matter in Extreme Astrophysical Environments	17
3.7 Facilities for Cosmic Probes of Dark Matter	18
3.7.1 Current/Near-Future Facilities	18
3.7.2 Future Facilities	20
3.8 Tools for Cosmic Probes of Dark Matter Physics	21
3.9 Roads to New Physics	22
3.9.1 Warm and Self-Interacting Dark Matter	23
3.9.2 Primordial Black Holes	24
3.9.3 Axion-like Dark Matter in Extreme Environments	25
3.10 Summary and Outlook	26

3.1 Three Core HEP Community Priorities

- Current/near-future HEP cosmology experiments have direct sensitivity to dark matter particle physics [1–3]. **Cosmological studies of dark matter should be supported as a key component of the HEP Cosmic Frontier program due to their unique ability to probe dark matter microphysics and link the results of terrestrial dark matter experiments to cosmological measurements.**
- **The construction of future cosmology experiments is critical for expanding our understanding of dark matter physics.** Proposed facilities across the electromagnetic spectrum, as well as gravitational waves, can provide sensitivity to dark matter physics, as well as physics of dark energy and the early universe [4]. HEP involvement will be essential in constructing and operating these facilities, and optimizing their sensitivity to dark matter physics should be a core consideration in their design.
- Cosmic probes provide robust sensitivity to the microphysical properties of dark matter due to enormous progress in theoretical modeling, numerical simulations, and astrophysical data. **Theory, simulation, observation, and experiment must be supported together to maximize the efficacy of cosmic probes of dark matter physics.**


3.1 Five Major Science Opportunities

1. The Standard Model of particle physics and cosmology can be tested at unprecedented levels of precision by measuring the cosmic distribution of dark matter. These measurements span an enormous range of scales from the observable universe to sub-stellar-mass systems (e.g., the matter power spectrum, the mass spectrum of dark matter halos, dark matter halo density profiles, and abundances of compact objects) [7, 12, 13]. The fundamental particle properties of dark matter (e.g., particle mass, production mechanism, and interaction cross sections) can lead to observable changes in the distribution of dark matter. **Measurements of the distribution of dark matter should be supported as a key element of the HEP Cosmic Frontier program to understand the fundamental nature of dark matter.**
2. The Λ CDM model makes the strong, testable prediction that the mass spectrum of dark matter halos extends below the threshold at which galaxies form [5]. Sub-galactic dark matter halos are less influenced by baryonic processes making them especially clean probes of fundamental physics of dark matter. We are on the cusp of detecting dark matter halos that are devoid of baryons through several cosmic probes (e.g., strong lensing, the dynamics of stars around the Milky Way). **The HEP community should pursue the detection of dark matter halos below the threshold of galaxy formation as an exceptional test of fundamental dark matter properties.**

3.1 Five Major Science Opportunities



3. Extreme astrophysical environments provide unique opportunities to explore dark matter couplings to the Standard Model that are inaccessible with terrestrial experiments [8]. **Instruments, observations, and theorists that study extreme astrophysical environments should be supported as an essential means to constrain the expanding landscape of dark matter models.**
4. Numerical simulations of structure formation and baryonic physics play a key role in addressing particle physics questions about the nature of dark matter. **HEP computational resources and expertise can be combined with astrophysical simulation expertise to rapidly advance numerical simulations of dark matter physics.**
5. The interdisciplinary nature of dark matter research calls for interagency mechanisms that support a comprehensive pursuit of scientific opportunities cutting across traditional disciplinary boundaries.

3.2 - CF3 in a single figure...




3.3 Dark Matter Halos

2203.07354



Dark matter **particle physics** changes the astrophysical **structure and distribution** of dark matter

3.3 Complementarity with CF1 (Halos)




Dark matter-nucleon scattering cross section vs dark matter mass.

Projections come from assuming sensitivity to 10^5 Msun halos.

Still in progress

3.4 Numerical Simulations




Bullock & Boylan-Kolchin 2017

Numerical simulations are critically important to interpret cosmic observations in the context of specific dark matter particle models.

Collaborations between computation and particle theory is critical. Problem is well-matched to HEP computational resource and expertise.

[2203.07049](#)

Measuring Dark Matter Physics using Cosmological Simulations



- Need #1:** Collaboration between simulators and particle theorists
- Need #2:** Algorithm development and code comparison tests
- Need #3:** Hydrodynamic simulations for observational targets
- Need #4:** Compare simulations to data in observable parameter space
- Need #5:** Fast realizations of observed systems to constrain dark matter
- Need #6:** Provide guidance to observers about dark matter signatures

Example pipeline for translating between particle physics and observations.

Astrophysics > Cosmology and Nongalactic Astrophysics

[Submitted on 20 May 2022 (v1), last revised 6 Jul 2022 (this version, v2)]

Tidal disruption of solitons in self-interacting ultralight axion dark matter

Noah Glennon, Ethan O. Nadler, Nathan Musoke, Arka Banerjee, Chanda Prescod-Weinstein, Risa H. Wechsler

Ultralight axions (ULAs) are promising dark matter candidates that can have a distinct impact on the formation and evolution of structure on nonlinear scales relative to the cold, collisionless dark matter (CDM) paradigm. However, most studies of structure formation in ULA models do not include the effects of self-interactions, which are expected to arise generically. Here, we study how the tidal evolution of solitons is affected by ULA self-interaction strength and sign. Specifically, using the pseudospectral solver UltraDark.jl, we simulate the tidal disruption of self-interacting solitonic cores as they orbit a $10^{11} M_{\odot}$ Navarro-Frenk-White CDM host halo potential for a range of orbital parameters, assuming a fiducial ULA particle mass of 10^{-22} eV. We find that repulsive (attractive) self-interactions significantly accelerate (decelerate) soliton tidal disruption. We also identify a degeneracy between the self-interaction strength and soliton mass that determines the efficiency of tidal disruption, such that disruption timescales are affected at the $\sim 50\%$ level for variations in the dimensionless ULA self-coupling from $\lambda = -10^{-92}$ to $\lambda = 10^{-92}$.

Comments: 15 pages, 10 figures. This paper was made to be as similar to the PRD version as possible

Subjects: **Cosmology and Nongalactic Astrophysics (astro-ph.CO)**; High Energy Physics – Phenomenology (hep-ph)

Cite as: arXiv:2205.10336 [astro-ph.CO]

(or arXiv:2205.10336v2 [astro-ph.CO] for this version)


<https://doi.org/10.48550/arXiv.2205.10336> 

Related DOI: <https://doi.org/10.1103/PhysRevD.105.123540> 

3.5 Primordial Black Holes and the Early Universe

Primordial black holes may be our earliest window into the birth of the universe and energies between the QCD phase transition and the Planck scale.

Even if PBHs are a small fraction of dark matter, their discovery would have far-reaching implications




3.6 Extreme Environments in one Figure...


[2203.07984](#)

Extreme astrophysical environments (black holes, stellar remnants, and stellar interiors) provide unique opportunities to explore dark matter couplings to the Standard Model that are inaccessible with terrestrial experiments.

Support experiments, observations, and theoretical interpretation that probe extreme environments.



3.6 Complementarity with CF2 (Extreme Environments)



Axion-photon coupling vs dark matter mass.

Projections come from a range of places.

Still in progress

3.7 Facilities for Cosmic Probes of Dark Matter


Current/Near-Future Facilities

Rubin LSST



Atacama, Chile

DESI





CMB-S4




Future Facilities


Stage 5 Spectroscopy



CMB-HD



Gravitational Wave Facilities



Gamma-ray/ X-ray Telescopes

Precise Radial Velocity

Radio Facilities

3.7 Facilities for Cosmic Probes of Dark Matter

- How do we best address the need from the facilities community?


Dark matter physics associated with current and near-future facilities, such as DESI, Rubin, and CMB-S4, is extremely rich. Dark matter science should be supported within these projects on intermediate scales in parallel to studies of dark energy and inflation. Such a program will fully leverage the unprecedented capabilities of these facilities. On large scales, the construction of future cosmology experiments is critical for expanding our understanding of dark matter physics. HEP involvement will be essential for the design and construction of these facilities, and dark matter physics should be a core component of their scientific mission.

- We need to highlight the relevance of technology and expertise of the HEP community.
- We need provide more specific goals and quantitative estimates; these exist for some, but not all facilities.


3.8 Tools for Cosmic Probes of Dark Matter Physics

- **Collaborative Infrastructure** - Support through existing HEP Projects (DESI, Rubin, CMB-S4)
- **New Support Mechanisms** - Cross-disciplinary support initiatives (future DMNI, cross-disciplinary funding)
- **Artificial Intelligence/Machine Learning** - Large complex data sets; need new tools to analyze them.
- **Cosmology Data Preservation** - Large legacy data sets; want to re-analyze for decades to come


3.9 Roadmap to New Physics



SIDM/WDM w/ Rubin LSST and future



Primordial black holes w/ Rubin/Roman



Axion-like particles w/ EHT

Astrophysical uncertainties can be **controlled** and the **extraction of fundamental dark matter properties** is possible.

Snowmass 2013 did not have a CF3... The result from the 2014 P5 →

The 2014 P5 report did not identify dark matter as a science driver for the large cosmic survey efforts (LSST, DESI, CMB-S4).

Faced resistance in expanding the scientific scope of these experiments to support dark matter research *even though* dark matter is a DOE mission priority.

We would like to avoid this happening again...

Table 1
Summary of Scenarios

Project/Activity	Scenarios			Science Drivers				
	Scenario A	Scenario B	Scenario C	Higgs	Neutrinos	Dark Matter	Cosmic Acceleration	The Unknown (Frontier)
Large Projects								
Muon program: Mu2e, Muon g-2	Y, Mu2e small reprofile needed	Y	Y					✓ I
HL-LHC	Y	Y	Y	✓		✓		✓ E
LBNF + PIP-II	Y, LBNF components delayed relative to Scenario B	Y	Y, enhanced		✓			✓ I, C
ILC	R&D only	R&D, possibly small hardware contributions. See text.	Y	✓		✓		✓ E
NuSTORM	N	N	N		✓			I
RADAR	N	N	N		✓			I
Medium Projects								
LSST	Y	Y	Y		✓		✓	C
DM G2	Y	Y	Y			✓		C
Small Projects Portfolio	Y	Y	Y		✓	✓	✓	All
Accelerator R&D and Test Facilities	Y, reduced	Y, some reductions with redirection to PIP-II development	Y, enhanced	✓	✓	✓		E, I
CMB-S4	Y	Y	Y		✓		✓	C
DM G3	Y, reduced	Y	Y			✓		C
PINGU	Further development of concept encouraged				✓	✓		C
ORKA	N	N	N					✓ I
MAP	N	N	N	✓	✓	✓		✓ E, I
CHIPS	N	N	N		✓			I
LAr1	N	N	N		✓			I
Additional Small Projects (beyond the Small Projects Portfolio above)								
DESI	N	Y	Y		✓		✓	C
Short Baseline Neutrino Portfolio	Y	Y	Y		✓			I

Summary

- Dark matter should be **one of (if not “the one”) highest priority** across Snowmass.
- **Current/near-future cosmic surveys** provide **direct access to dark matter model space**.
- **Future cosmic survey facilities** provide discovery potential across a wide range of fundamental physics (dark energy, dark matter, inflation, early universe physics).
- **New mechanisms** for cross-disciplinary support are needed to assemble the expertise needed to make concrete advances in cosmic probes of dark matter.
- Cosmic probes of dark matter **complement terrestrial searches** (i.e., probe similar models in different parts of parameter space), **inform terrestrial searches** (i.e., tell us where the dark matter is and how it moves), and **probe unique parameter space** (i.e., self-interactions and gravitational interactions).