


# Analysis of the T-980 experimental data


*V.Previtali*

G. Annala, R.Assmann, N. Mokhov,  
S. G. Peggs, S. Redaelli, D.Still

# Layout of T-980



- Si o-shaped crystal, 5 mm long, 410  $\mu$ rad bending angle
- Pin diode downstream of the crystal, used to measure the inelastic interactions at the crystal location
- Collimator E03 (horizontal, p collimator) 23.7 m downstream
- LE0 BLM counters immediately downstream the collimator: total losses at the collimator location
- E1 scintillating paddles: gated counters for losses at the E03 collimator. They can distinguish between bunched and abort gap beam.




# Standard measurement procedure



Once we have the crystal as leading edge, we perform two different measurements:

- 1. Angular scan:** change the crystal orientation and measure losses at the E03 collimator
- 2. Collimator scan:** keep the angle of the crystal fixed, and change the horizontal position of the collimator E03


# Angular scan: what we expected...



The ideal behaviour:

- clear channeling region, width of  $2 \times$  critical angle (12  $\mu$ rad)
- clear volume reflection region, acceptance = channeling angle (410  $\mu$ rad)
- maybe a bump at the end of the VR region (as foreseen by simulations)

# Angular scan: what we found...




The signal for bunched beam is noisy: a normalization is needed. (how to do it???)

The maximum of the channeling peak is at  $-240 \mu\text{rad}$ .


The measured acceptance of channeling is  $\sim 200 \mu\text{rad}$ : **much larger than expected!** ( $\sim 12 \mu\text{rad}$ ).

Possible reasons are investigated further

# Collimator scan




# Collimator scan



- The collimator scan allows understanding the profile of the incoming beam
- We can measure the displacement between the channeled and the non-channeled beam
- The expected displacement (for 410 $\mu$ rad kick) is 9.5 mm

# Collimator scan

in the middle of channeling peak




The measured displacement (~7 mm) is much lower than expected.

The displacement is evaluated by fitting the channeling signal with an error function

$$f(x) = a \cdot \operatorname{erf}\left(\frac{(x-c)}{\sqrt{2}s}\right) + b$$

Final set of parameters	Asymptotic Standard Error	
=====	=====	
a	= 0.505231	+/- 0.009469 (1.874%)
b	= 0.821607	+/- 0.008615 (1.049%)
c	= -6.84432	+/- 0.5168 (7.551%)
s	= 10.8197	+/- 0.6928 (6.403%)






# What we could not understand:



1. Why is the channeling peak acceptance much larger ( $\sim 200 \mu\text{rad}$ ) than expected ( $\sim 20 \mu\text{rad}$ )?
2. Why we do not see a clear VR effect?
3. Why the measured displacement ( $\sim 7 \text{ mm}$ ) for the channeling peak is lower than the expected one ( $9.6 \text{ mm}$ )?
4. What is the peak at  $\sim 250/300 \mu\text{rad}$  in the angular scan? Is there a third peak around  $\sim 700 \mu\text{rad}$ ?



# Hypothesis




Different attempts have been made to explain these features. Two hypotheses:

## 1. Feature of the beam (momentum offset)

- Off momentum particles have a different incoming angle: can this explain the channeling peak width?
- Off momentum particles have a different displacement at the collimator location: how much is the difference?

## 2. Feature of the crystal (mis-cut angle)

- What is the effect of the mis-cut on the channeling acceptance?
- What is the effect of the mis-cut on the observed displacement at the collimator location?



# Hypothesis




Different attempts have been made to explain these features. Two hypotheses:

## 1. Feature of the beam (momentum offset)

- Off momentum particles have a different incoming angle: can this explain the channeling peak width?
- Off momentum particles have a different displacement at the collimator location: how much is the difference?

## 2. Feature of the crystal (mis-cut angle)

- What is the effect of the mis-cut on the channeling acceptance?
- What is the effect of the mis-cut on the observed displacement at the collimator location?



# Off-momentum particles



Taking into account that:

- the dispersion at the crystal (and at the collimator) is quite high (2m!)
- we are channeling also the abort gap beam
- the abort gap beam has high  $\Delta p/p$  values

We tried to evaluate the effect of dealing with large off-momentum particles

For reference:


1  $\sigma p/p$  in the tevatron is 140 MeV

The RF bucket height is 450 MeV

In the abort gap particles are just outside of the separatrix

**electron lens heating is turned on**

# Off-momentum particles: angular spread



$$n_0(\delta p/p) = n_{cry} - D_{cry} \cdot \frac{\delta p/p}{\sigma_{cry}}$$

g function =  $\alpha + \beta \eta'/\eta$


The **grazing condition** requires that, at the crystal location, the *maximum betatron extension* of the particle plus the *offset given by the dispersion* is equal to the *x coordinate of the crystal's edge*  
=> careful: the synchrotron oscillation is neglected in this first approximation.

$g = 2.9 \cdot 10^{-3}$   
 $\sigma_{p/p} = 2.9 \cdot 10^{-3}$

**The angular spread is only  $\sim 1.6 \mu\text{rad}$  for particles with  $\Delta p/p = 4 \sigma_p$  !**


**The momentum offset cannot explain a  $200 \mu\text{rad}$ -wide channeling peak**

# Off-momentum particles: displacement at the collimator



Every kick changes the amplitude and the phase of the particle. The new amplitude and the phase shift depend on the initial amplitude: different outcomes for particles with different energy!


Particles with higher  $\Delta p/p$ , will have different amplitude/phase shift in comparison with on momentum particles => they will have different displacement at the collimator. How much?



# Off-momentum particles: displacement at the collimator




Assuming the channeling kick of  $410 \mu\text{rad}$



The displacement is higher for larger energy offset.

Anyway the difference is only  $100 \mu\text{m}$ !!!

**The momentum offset cannot explain the reduced displacement observed at E03**



# Hypothesis



Different attempts have been made to explain these features. Two hypotheses:

## 1. Feature of the beam (momentum offset)

- Off momentum particles have a different incoming angle: can this explain the channeling peak width?
- Off momentum particles have a different displacement at the collimator location: how much is the difference?

## 2. Feature of the crystal (mis-cut angle)


- What is the effect of the mis-cut on the channeling acceptance?
- What is the effect of the mis-cut on the observed displacement at the collimator location?



# Influence of the mis-cut angle




- The mis-cut angle of the crystal is very large (1.3 mrad over 0.41 mrad of bending angle)



- Even if we are in the “good” orientation, the mis-cut could affect the particle-crystal interactions. In the following we analyze the problem in details.


# particles aligned for channeling




In this region the particles are aligned: channeling with full channeling angle ( $410 \mu\text{rad}$ ). Impact parameter  $>5 \mu\text{m}$

- Particles are aligned with the crystal planes at the entrance face

# Positive mis-cut angle: crystal aligned for channeling




- Particles are aligned with the crystal planes at the entrance face:
  - The closest point to the beam is the end of the crystal
  - They will have to cross  $\sim 5 \mu\text{m}$  of amorphous layer before being channeled



# Positive mis-cut angle: crystal aligned for VR




1. In this region the particles are not aligned: amorphous layer.  
Impact parameter  $0\mu\text{m} < \lambda_0 < 5\mu\text{m}$



# Positive mis-cut angle: crystal aligned for VR



1. In this region the particles are not aligned: amorphous layer.  
Impact parameter  $0\mu\text{m} < \lambda_0 < 5\mu\text{m}$



3. In this region the particles are aligned for  
Volume Reflection / Volume Capture (low  
probability).  
Impact parameter  $\lambda > \lambda_0$


# Positive mis-cut angle: crystal aligned for VR




1. In this region the particles are not aligned: amorphous layer.  
Impact parameter  $0 \mu\text{m} < \lambda_0 < 5 \mu\text{m}$

2. In this region the particles are aligned:  
channeling with **reduced channeling angle**  
( $< 410 \mu\text{rad}$ ).  
Impact parameter  $\lambda \sim \lambda_0$  parameter.  
The width of this region is  $\Delta\lambda \sim 0.25 \mu\text{m}$

3. In this region the particles are aligned for  
Volume Reflection / Volume Capture.  
Impact parameter  $\lambda > \lambda_0$




# Positive mis-cut angle: crystal aligned for VR



For each orientation of the crystal, there will be an impact parameter  $\lambda_0$  for which the particles are aligned with crystal planes

=> **channeling**, but with a reduced channeling angle! This could explain the reduced displacement at the collimator AND the larger channeling peak.




## Positive mis-cut angle:

# interpretation of measured data




- For each orientation there will be a superposition of the three effects (reduced channeling, VR, VC)
- **We channel in each orientation, but with reduced channeling angles!** We can calculate this reduced angle, and **predict the displacement** at the collimator.



**How to interpret our angular scan?**


NB: Reduced channeling and Volume Capture give the same kick to the particle, but the channeling probability is much higher.






## Positive mis-cut angle:

# interpretation of measured data



-350  $\mu\text{rad}$   
Beginning of channeling:  
full channeling kick is expected

-350  $\mu\text{rad} < \theta < +60 \mu\text{rad}$   
intermediate orientation: “reduced”  
channeling kicks are expected  
(it scales linearly)  
also amorphous and channeling should  
take place




+60  $\mu\text{rad}$   
End of channeling: all the  
crystal coherent effects should  
stop

**Completely new  
interpretation of the  
measured data!**








# Positive mis-cut angle: comparison with measured data



- We selected 6 different orientations for new collimator scans:
  - 320  $\mu\text{rad}$
  - 295  $\mu\text{rad}$
  - 287  $\mu\text{rad}$
  - 237  $\mu\text{rad}$
  - 200  $\mu\text{rad}$
  - 50  $\mu\text{rad}$
- For each point we measure the displacement of the channeled peak, and compare it with the expected displacement



# LHC Collimation




fnal - 3 Dec 2008

# All together...




LE033 losses versus collimator position (E03HCP)



# typical shape of a collimator scan



LE033 losses versus collimator position (E03HCP)




All the E03 scans have the same shape, apart from the scan at  $-50 \mu\text{rad}$ , where the channeling peak is not visible

# All together... Gaussian fits




LE033 losses versus collimator position (E03HCP)



All together...


# Gaussian fits vs theory



# Displacement for different orientations of the crystal



measured and expected




The mis-cut can probably explain the reduced displacement observed at E03

Good agreement

The displacement is larger than expected!

In the experimental error?  
Must be evaluated...






# All together... more questions




What is this behavior? It is common to all the angular scans.  
It is equivalent to a r.m.s. kick of **100  $\mu\text{rad}$**  ! Cannot be amorphous (typical kick 3.2  $\mu\text{rad}$ ) or single VR (-6  $\mu\text{rad}$ ). Cannot be de-channeling (cannot be larger than the channeling kick).  
Maybe **multiple volume reflection?**  
This effect probably covers the “reduced” channeled peak for small channeling kicks!!!  
That’s why we do not observe the correct displacement for the -50  $\mu\text{rad}$  orientation.


... versus collimator position (E03HCP)





# Angular scan: a wider view






# More questions?



- What is the peak we see at 250-300  $\mu\text{rad}$  in the angular scan? Is there a third peak at  $\sim 700 \mu\text{rad}$ ? Are they channeling peaks? Are we almost aligned in vertical position?
- What is the final slope we observe in each collimator scan, which corresponds to a typical kick of  $-100 \mu\text{rad}$ ? Is it multiple volume reflection?
- What is the effect of the electron lens in the particle-crystal dynamic?
- Is the synchrotron oscillation playing an important role? Is the “grazing” assumption valid? Should we evaluate in more details the impacting angle of off momentum particles?



# Even more questions...

(from Steve)



- 1. what would you expect to see in the H8/RD22 single particle line, with this crystal?
- 2. What happens if the electron lens heating is turned off/down?
- 3. What happens if the RF voltage is turned up/down (moving the separatrix)?
- 4. What happens if the RF frequency itself is slightly changed, shifting the underside of the separatrix slightly up/down?
- 5. Synchrotron oscillations **MUST** be playing a vital role
  - a) in providing lots of time for betatron heating as the protons go oh-so-slowly past the unstable fixed point
  - b) in helping to determine the actual spread of impact parameters as the protons go at normal speed (as in a regular synchrotron oscillation) past the bottom of the RF bucket.

# Conclusions



- We trust the characterization of the crystal made in Ferrara (V. Guidi et al.): we assume the bending angle=410  $\mu$ rad.
- We observe a channeling acceptance that is far too large.
- We observe a displacement of the channeled beam at the collimator which is lower than expected (= lower kick).
- Different hypothesis to explain this features:
  - Feature of the beam (momentum offset)
  - Feature of the crystal (mis-cut angle)
- The momentum offset does not have large influence on the displacement.
- **The mis-cut angle can partially explain the results we observe.**
- ***A lot of open questions...***
- Further investigations will be done in the next studies.