

Family Non-universal $U(1)'$ Gauge Symmetries and $b \rightarrow s$ Transitions

Tao Liu

**Enrico Fermi Institute
University of Chicago**

Collider Physics 2009

Joint Argonne and IIT Theory Institute

05/18/2009

- V. Barger, L. Everett, J. Jiang, P. Langacker, T.L. and C. Wagner, “Family Non-universal $U(1)'$ Gauge Symmetries and $b \rightarrow s$ Transitions,” arXiv:0902.4507 [hep-ph].
- V. Barger, L. Everett, J. Jiang, P. Langacker, T.L. and C. Wagner, “ $b \rightarrow s$ Transitions in Family-dependent $U(1)'$ Models,” arXiv:0905.xxxx [hep-ph].

- Recent Experimental Progresses on $b \rightarrow s$ FCNC Effects
- Family Non-universal $U(1)'$ Models and $b \rightarrow s$ Transitions
- Correlated Analysis of $B_s - \bar{B}_s$ Mixing and B_d Decays
- Conclusions

Recent Experimental Progresses on $b \rightarrow s$ FCNC Effects

- Three classes of FCNC processes:
 - $d \rightarrow s$ transitions: e.g., $K - \bar{K}$ mixing
 - $b \rightarrow d$ transitions: e.g., $B_d - \bar{B}_d$ mixing
 - $b \rightarrow s$ transitions: e.g., $B_s - \bar{B}_s$ mixing
- Absent at tree level in the SM and sensitive to UV physics.
⇒ A useful tool to probe or constrain NP models.

A. $\Delta B = 2$ processes via $b \rightarrow s$: $B_s - \bar{B}_s$ mixing

Key observables: off-diagonal element of the mixing matrix

$$M_{12}^{B_s} = (M_{12}^{B_s})_{\text{SM}} C_{B_s} e^{2i\phi_{B_s}^{\text{NP}}}$$

SM predictions: $C_{B_s} = 1$ and $\phi_{B_s}^{\text{NP}} = 0$

Table: The fit results for the $B_s - \bar{B}_s$ mixing parameters (**UTfit Collaboration' 08**), obtained by combining the analyses of $B_s \rightarrow \psi\phi$ by (**CDF Collaboration' 08**) and (**D0 Collaboration' 08**)

Observable	1σ C.L.	2σ C.L.
$\phi_{B_s}^{\text{NP}} [^\circ]$ (S1)	-19.9 ± 5.6	$[-30.45, -9.29]$
$\phi_{B_s}^{\text{NP}} [^\circ]$ (S2)	-68.2 ± 4.9	$[-78.45, -58.2]$
C_{B_s}	1.07 ± 0.29	$[0.62, 1.93]$

- C_{B_s} is consistent with its SM prediction, for given significance.
- $\phi_{B_s}^{\text{NP}}$ has two solutions, denoted as “S1” and “S2”.
- **Both of them deviate from its SM prediction by more than 3σ .**

B. $\Delta B = 1$ processes via $b \rightarrow s$: $B_d \rightarrow (\phi, \eta', \pi, \rho, \omega, f_0) K_S$
 Key observables: time-dependent CP asymmetries

$$C_{f_{CP}} = \frac{1 - |\lambda_{f_{CP}}|^2}{1 + |\lambda_{f_{CP}}|^2}, \quad S_{f_{CP}} = \frac{2\text{Im}[\lambda_{f_{CP}}]}{1 + |\lambda_{f_{CP}}|^2}. \quad (1)$$

The $\lambda_{f_{CP}}$ parameter ($A_{f_{CP}}$ is decay amplitude of $B_d \rightarrow f_{CP}$):

$$\lambda_{f_{CP}} \equiv \eta_{f_{CP}} e^{-2i\phi_{B_d}} \frac{\bar{A}_{f_{CP}}}{A_{f_{CP}}} \quad (2)$$

SM prediction: $\phi_{B_d} = \beta \equiv \arg[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$ and $\bar{A}_{f_{CP}} \simeq A_{f_{CP}} \Rightarrow$

$$C_{f_{CP}} \simeq 0, \quad -\eta_{f_{CP}} S_{f_{CP}} \simeq \sin 2\beta. \quad (3)$$

Table: World averages of the experimental results (HFAG'08).

f_{CP}	$-\eta_{CP} \mathcal{S}_{f_{CP}} (1\sigma \text{ C.L.})$	$\mathcal{C}_{f_{CP}} (1\sigma \text{ C.L.})$
ψK_S	$+0.672 \pm 0.024$	$+0.005 \pm 0.019$
ϕK_S	$+0.44^{+0.17}_{-0.18}$	-0.23 ± 0.15
$\eta' K_S$	$+0.59 \pm 0.07$	-0.05 ± 0.05
πK_S	$+0.57 \pm 0.17$	$+0.01 \pm 0.10$
ρK_S	$+0.63^{+0.17}_{-0.21}$	-0.01 ± 0.20
ωK_S	$+0.45 \pm 0.24$	-0.32 ± 0.17
$f_0 K_S$	$+0.62^{+0.11}_{-0.13}$	0.10 ± 0.13

- $\sin 2\beta$ obtained from the penguin-dominated modes are systematically below that obtained from $B_d \rightarrow \psi K_S$.
- $|\mathcal{C}_{(\phi,\omega)K_S}| \gg |\mathcal{C}_{\psi K_S}|$
- $B_d \rightarrow \psi K_S$ is dominated by the SM tree-level amplitude, $-\eta_{f_{CP}} \mathcal{S}_{f_{CP}} \neq -\eta_{\psi K_S} \mathcal{S}_{\psi K_S}$ or $\mathcal{C}_{f_{CP}} \neq \mathcal{C}_{\psi K_S}$ may imply interesting NP.

Non-universal $U(1)'$ Models and $b \rightarrow s$ Transitions


Non-universal $U(1)'$: (1) quarks, leptons charged in a family dependent way; (2) their couplings to Z' are phenomenologically characterized by the following structure:

$$B^{\psi_{L,R}} \equiv \frac{g_2 M_Z}{g_1 M_{Z'}} V_{\psi_{L,R}} \tilde{\epsilon}^{\psi_{L,R}} V_{\psi_{L,R}}^\dagger = \begin{pmatrix} B_{11}^{\psi_{L,R}} & 0 & B_{13}^{\psi_{L,R}} \\ 0 & B_{11}^{\psi_{L,R}} & B_{23}^{\psi_{L,R}} \\ B_{13}^{\psi_{L,R}*} & B_{23}^{\psi_{L,R}*} & B_{33}^{\psi_{L,R}} \end{pmatrix}$$

- $\psi_{L,R}$ – weak eigenstates of the SM fermions
 $\tilde{\epsilon}^{\psi_{L,R}}$ – $U(1)'$ gauge charge matrix of $\psi_{L,R}$, which is diagonal
 $V_{\psi_{L,R}}$ – the unitary matrices diagonalizing fermion mass matrices
- Non-trivial $B_{12}^{\psi_{L,R}}$ is excluded by $K - \bar{K}$ and $\mu - e$ constraints;
 $B_{23}^{\psi_{L,R}}$ can lead to sizable FCNC effects in $b \rightarrow s$ transitions;
 $B_{13}^{\psi_{L,R}} \sim B_{23}^{\psi_{L,R}}$ may also lead to non-trivial FCNC effects in $b \rightarrow d$ transitions.
- Such a structure can be generated in the case with
 $\tilde{\epsilon}_1^{\psi_{L,R}} = \tilde{\epsilon}_2^{\psi_{L,R}} \neq \tilde{\epsilon}_3^{\psi_{L,R}}$ and small fermion mixing angles in $V_{\psi_{L,R}}$.

Through current-current interactions, three classes of processes via $b \rightarrow s$ can be affected at tree-level by Z' -induced FCNC effects:

- $B_s - \bar{B}_s$ mixing;
- $b \rightarrow s\bar{q}q$, e.g., hadronic B_d meson decays;
- $b \rightarrow s\bar{l}l$, e.g., $B_s \rightarrow l^+l^-$



$$\begin{aligned}
\mathcal{H}_{\text{eff}}^{Z'}(B_s - \bar{B}_s) &= -\frac{G_F}{\sqrt{2}}(\Delta C_1^{B_s} Q_1^{B_s} + 2\Delta \tilde{C}_3^{B_s} \tilde{Q}_3^{B_s} + \Delta \tilde{C}_1^{B_s} \tilde{Q}_1^{B_s}) + h.c. \\
\mathcal{H}_{\text{eff}}^{Z'}(b \rightarrow s\bar{q}q) &= -\frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* (\Delta C_3 Q_3 + \Delta C_5 Q_5 + \Delta C_7 Q_7 + \Delta C_9 Q_9 \\
&\quad + \Delta \tilde{C}_3 \tilde{Q}_3 + \Delta \tilde{C}_5 \tilde{Q}_5 + \Delta \tilde{C}_7 \tilde{Q}_7 + \Delta \tilde{C}_9 \tilde{Q}_9) + h.c. \\
\mathcal{H}_{\text{eff}}^{Z'}(b \rightarrow s\bar{l}l) &= -\frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* (\Delta C_{9V} Q_{9V} + \Delta C_{10A} Q_{10A} \\
&\quad + \Delta \tilde{C}_{9V} \tilde{Q}_{9V} + \Delta \tilde{C}_{10A} \tilde{Q}_{10A}) + h.c.
\end{aligned}$$


- Q_i s: SM operators in OPE; \tilde{Q}_i s: new operators introduced by NP
- Q_i s and \tilde{Q}_i s: QCD penguin for $i = 3, \dots, 6$ and EW penguin for $i = 7, \dots, 10$
- Effectively, NP affects $b \rightarrow s$ FCNC processes: (1) by modifying Wilson coefficients of the SM operators, and (2) by introducing new FC operators

Correlated Analysis of $B_s - \bar{B}_s$ Mixing and B_d Decays

Strategy to study $B_s - \bar{B}_s$ mixing + Penguin-dominated B_d decays:


- Assume no $Z - Z'$ mixing, due to the constraints from weak neutral current data;
- Assume that NP enters mainly through EW penguins (A. Buras et. al.' 04), i.e., $\Delta C_{3,5}, \Delta \tilde{C}_{3,5} \ll \Delta C_7$ or $\Delta \tilde{C}_9 \Rightarrow$
 - Five relevant parameters: $|B_{bs}^{L,R}|$, $\phi_{bs}^{L,R}$ and B_{dd}^R ;
 - $|B_{bs}^L| < |B_{dd}^L| \ll |B_{dd}^R|$.
- Assume 15% and 25% non-perturbative uncertainties in the SM and NP calculations, respectively.

Figure: Distributions of $|B_{bs}^{L,R}|$ and $\phi_{bs}^{L,R}$ which are constrained by $B_s - \bar{B}_s$ mixing, in the general case.



- Blue: constrained by 1σ experimental data.
Purple: constrained by 2σ experimental data.
- Two representative limits:
 - Diagonal lines (left-bottom \rightarrow right-up): $B_{bs}^L = B_{bs}^R$ (LR limit)
 - Bottom line: $\epsilon^{\psi_R} \propto I \Rightarrow B_{bs}^R = 0$ (LL limit)
 - Three free parameters: $|B_{bs}^L|$, ϕ_{bs}^L and B_{dd}^R


Figure: $B_s - \bar{B}_s$ mixing, in the LR and LL limits.



$$C_{B_s} e^{2i\phi_{B_s}^{\text{NP}}} = 1 - 3.59 \times 10^5 (\Delta C_1^{B_s} + \Delta \tilde{C}_1^{B_s}) + 2.04 \times 10^6 \Delta \tilde{C}_3^{B_s} \quad (4)$$


- LR limit: $\Delta C_1^{B_s} = \Delta \tilde{C}_1^{B_s} = \Delta \tilde{C}_3^{B_s} = -(B_{bs}^L)^2$
- LL limit: $\Delta C_1^{B_s} = -(B_{bs}^L)^2, \Delta \tilde{C}_1^{B_s} = \Delta \tilde{C}_3^{B_s} = 0$
- Two parameters involved: $|B_{bs}^L|$ and ϕ_{bs}^L ; $|B_{bs}^L| \sim 10^{-3}$.

Figure: $B_d \rightarrow \pi K_S$, in the LR and LL limits.




- A deviation of $S_{\pi K_S}$ from its SM prediction can be understood as a modification of $qe^{i\phi} = \frac{P}{T+C}$ (A. Buras et. al.' 04). The experimental constraints on $qe^{i\phi}$ have been obtained from χ^2 fit of the $B \rightarrow \pi K + B \rightarrow \pi\pi$ data (R. Fleischer et. al.' 08)
- In our models: $qe^{i\phi} = 0.76(1 + 158.1\Delta C_7 - 102.4\Delta\tilde{C}_9)$
- LR (LL) limit: $\Delta C_7 = \Delta\tilde{C}_9 = \frac{4}{V_{tb}V_{ts}^*} B_{bs}^L B_{dd}^R$ ($\Delta C_7 = \frac{4}{V_{tb}V_{ts}^*} B_{bs}^L B_{dd}^R$)

Figure: $B_s - \bar{B}_s$ mixing (2σ C.L.) + χ^2 fit of ($B \rightarrow \pi K_S + B \rightarrow \pi\pi$) (1σ C.L.) + $\mathcal{C}_{(\phi, \eta', \rho, \omega, f_0)K_S}$, $\mathcal{S}_{(\phi, \eta', \rho, \omega, f_0)K_S}$ (1.7σ C.L.), in the LR and LL limit.



- Both solutions “S1” and “S2” in $B_s - \bar{B}_s$ mixing can be explained.
- $|B_{dd}^R| \lesssim 10^{-1}$. But, to get better fit, ...

Figure: $B_d \rightarrow \pi K_S$ (correlated analysis), in the LR and LL limits



- To get a relatively small χ^2 value, we need $|B_{dd}^R| \gtrsim 10^{-2}$.
- Similar effects can be seen by reducing the C.L. in the fitting of $\mathcal{C}_{(\phi, \eta', \rho, \omega, f_0)K_S}$ and $\mathcal{S}_{(\phi, \eta', \rho, \omega, f_0)K_S}$.

Consistency checks:

- $|B_{bs}^L| \sim 10^{-3}$, consistent with the assumption of small fermion mixing angles, since $B_{bs}^L \propto \frac{g_2 m_Z}{g_1 m_{Z'}} \times \text{fermion mixing angle}$.
- The experimental constraints from $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ can be easily satisfied, according to

$$\left\{ \left| 0.003 + \frac{B_{bs}^L B_{\mu\mu}^L}{V_{tb}^* V_{ts}} \right|^2 + \left| \frac{B_{bs}^L B_{\mu\mu}^R}{V_{tb}^* V_{ts}} \right|^2 \right\}^{1/2} \lesssim 10^{-2}$$

- $|B_{bd}^L| \sim |B_{bs}^L| \Rightarrow$ To be consistent with $B_d - \bar{B}_d$ mixing, a $\mathcal{O}(10\%)$ fine-tuning is necessary.
- $|B_{dd}^R| \gtrsim 10^{-2}$ is typically required for a good fit. $\Rightarrow \frac{g_1 m_{Z'}}{g_2 m_Z} \sim 10 - 100$ or TeV scale Z' for $g_1 \gtrsim g_2$, **a range approachable at the LHC!**
- Recall that the assumption of small QCD penguin corrections by the NP requires $|B_{bs}^L| < |B_{dd}^L| \ll |B_{dd}^R|$. This relation can be easily accommodated.

Conclusions

- The recent progresses in experiments on $b \rightarrow s$ FCNC effects provide a good chance to study possible NP;
- Within this class of family non-universal $U(1)'$ models, the anomalies in $B_s - \bar{B}_s$ mixing and the time-dependent CP asymmetries of the penguin-dominated $B_d \rightarrow (\pi, \phi, \eta', \rho, \omega, f_0)K_S$ can be consistently accommodated;
- No single measurement of $B_s - \bar{B}_s$ mixing phase or time-dependent CP asymmetries of penguin-dominated B_d decays has a 3 sigma significance. For a better understanding of the possible NP, we expect that more precise information can be provided by experimentalists in the near future.

Thank you!

A. Operators of Effective Hamiltonian

QCD-Penguins Operators:

$$Q_3 = (\bar{s}b)_{V-A} \sum_q (\bar{q}q)_{V-A}$$

$$Q_5 = (\bar{s}b)_{V-A} \sum_q (\bar{q}q)_{V+A}$$

$$\tilde{Q}_3 = (\bar{s}b)_{V+A} \sum_q (\bar{q}q)_{V+A}$$

$$\tilde{Q}_5 = (\bar{s}b)_{V+A} \sum_q (\bar{q}q)_{V-A}$$

$$Q_4 = (\bar{s}_\alpha b_\beta)_{V-A} \sum_q (\bar{q}_\beta q_\alpha)_{V-A}$$

$$Q_6 = (\bar{s}_\alpha b_\beta)_{V-A} \sum_q (\bar{q}_\beta q_\alpha)_{V+A}$$

$$\tilde{Q}_4 = (\bar{s}_\alpha b_\beta)_{V+A} \sum_q (\bar{q}_\beta q_\alpha)_{V+A}$$

$$\tilde{Q}_6 = (\bar{s}_\alpha b_\beta)_{V+A} \sum_q (\bar{q}_\beta q_\alpha)_{V-A}$$

Electroweak-Penguins Operators:

$$Q_7 = \frac{3}{2} (\bar{s}b)_{V-A} \sum_q e_q (\bar{q}q)_{V+A}$$

$$Q_9 = \frac{3}{2} (\bar{s}b)_{V-A} \sum_q e_q (\bar{q}q)_{V-A}$$

$$\tilde{Q}_7 = \frac{3}{2} (\bar{s}b)_{V+A} \sum_q e_q (\bar{q}q)_{V-A}$$

$$\tilde{Q}_9 = \frac{3}{2} (\bar{s}b)_{V+A} \sum_q e_q (\bar{q}q)_{V+A}$$

$$Q_8 = \frac{3}{2} (\bar{s}_\alpha b_\beta)_{V-A} \sum_q e_q (\bar{q}_\beta q_\alpha)_{V+A}$$

$$Q_{10} = \frac{3}{2} (\bar{s}_\alpha b_\beta)_{V-A} \sum_q e_q (\bar{q}_\beta q_\alpha)_{V-A}$$

$$\tilde{Q}_8 = \frac{3}{2} (\bar{s}_\alpha b_\beta)_{V+A} \sum_q e_q (\bar{q}_\beta q_\alpha)_{V-A}$$

$$\tilde{Q}_{10} = \frac{3}{2} (\bar{s}_\alpha b_\beta)_{V+A} \sum_q e_q (\bar{q}_\beta q_\alpha)_{V+A}$$

$B_s - \bar{B}_s$ Mixing Operators:

$$Q_1^{B_s} = (\bar{s}b)_{V-A}(\bar{s}b)_{V-A}$$

$$\tilde{Q}_1^{B_s} = (\bar{s}b)_{V+A}(\bar{s}b)_{V+A}$$

$$Q_3^{B_s} = \tilde{Q}_3^{B_s} = (\bar{s}b)_{V+A}(\bar{s}b)_{V-A}$$

$$Q_2^{B_s} = (\bar{s}_\alpha b_\beta)_{V-A}(\bar{s}_\beta b_\alpha)_{V-A}$$

$$\tilde{Q}_2^{B_s} = (\bar{s}_\alpha b_\beta)_{V+A}(\bar{s}_\beta b_\alpha)_{V+A}$$

$$Q_4^{B_s} = \tilde{Q}_4^{B_s} = (\bar{s}_\alpha b_\beta)_{V+A}(\bar{s}_\beta b_\alpha)_{V-A}$$

B. Corrections of Family Non-universal Z' to Wilson Coefficients

(1) LR limit: $B_{bs}^L = B_{bs}^R$

$$\Delta C_1^{B_s} = \Delta \tilde{C}_1^{B_s} = \Delta \tilde{C}_3^{B_s} = -(B_{bs}^L)^2,$$

$$\Delta C_3 = \Delta \tilde{C}_5 = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L B_{dd}^L,$$

$$\Delta \tilde{C}_3 = \Delta C_5 = -\frac{2}{3V_{tb} V_{ts}^*} B_{bs}^L (B_{uu}^R + 2B_{dd}^R),$$

$$\Delta C_7 = \Delta \tilde{C}_9 = -\frac{4}{3V_{tb} V_{ts}^*} B_{bs}^L (B_{uu}^R - B_{dd}^R),$$

$$\Delta C_{9V} = \Delta \tilde{C}_{9V} = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L (B_{ll}^L + B_{ll}^R),$$

$$\Delta C_{10A} = \Delta \tilde{C}_{10A} = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L (-B_{ll}^L + B_{ll}^R).$$

(2) LL limit: $\epsilon^{\psi_R} \propto I$

$$\Delta C_1^{B_s} = -(B_{bs}^L)^2,$$

$$\Delta C_3 = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L B_{dd}^L,$$

$$\Delta C_5 = -\frac{2}{3 V_{tb} V_{ts}^*} B_{bs}^L (B_{uu}^R + 2B_{dd}^R),$$

$$\Delta C_7 = -\frac{4}{3 V_{tb} V_{ts}^*} B_{bs}^L (B_{uu}^R - B_{dd}^R),$$



$$\Delta C_{9V} = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L (B_{ll}^L + B_{ll}^R),$$

$$\Delta C_{10A} = -\frac{2}{V_{tb} V_{ts}^*} B_{bs}^L (-B_{ll}^L + B_{ll}^R).$$

(3) RR limit: $\epsilon^{\psi L} \propto I$

$$\begin{aligned}\Delta \tilde{C}_1^{B_s} &= -(B_{bs}^R)^2, \\ \Delta \tilde{C}_3 &= -\frac{2}{3V_{tb}V_{ts}^*} B_{bs}^R (B_{uu}^R + 2B_{dd}^R), \\ \Delta \tilde{C}_5 &= -\frac{2}{V_{tb}V_{ts}^*} B_{bs}^R B_{dd}^L, \\ \Delta \tilde{C}_9 &= -\frac{4}{3V_{tb}V_{ts}^*} B_{bs}^R (B_{uu}^R - B_{dd}^R), \\ \Delta \tilde{C}_{9V} &= -\frac{2}{V_{tb}V_{ts}^*} B_{bs}^R (B_{ll}^L + B_{ll}^R), \\ \Delta \tilde{C}_{10A} &= -\frac{2}{V_{tb}V_{ts}^*} B_{bs}^R (-B_{ll}^L + B_{ll}^R).\end{aligned}$$

C. Correlated Analyses in LR limit



D. Constraints on NP from $B_s - \bar{B}_s$ mixing (by CKMfitter)

Figure: Constraints on NP from $B_s - \bar{B}_s$ mixing (by CKMfitter), here

$$\Delta_S = C_{B_s} e^{2i\phi_{B_s}^{\text{NP}}}.$$

