

PREDICTIONS FOR HEAVY QUARKONIUM AT WEAK COUPLING

Antonio Pineda

Universitat Autònoma de Barcelona (IFAE)

QWG: International Workshop on Heavy Quarkonium, May 18 - 21, 2010

Motivation:

- ▶ 1st principle computation of heavy quarkonium properties from QCD
- ▶ Determination of Standard Model parameters: m_Q , α_s , ...

We have an effective field theory, **Potential Non-Relativistic QCD**, which describes the heavy quarkonium dynamics in the weak and strong coupling situation. $m \gg mv \gg mv^2$

$$\left. \begin{array}{l} \left(i\partial_0 - \frac{\mathbf{p}^2}{2m} - V_s^{(0)}(r) \right) \Phi(\mathbf{r}) = 0 \\ + \text{corrections to the potential} \\ + \text{interaction with other low} \\ \quad \text{energy degrees of freedom} \end{array} \right\} \text{potential NRQCD} \quad E \sim mv^2$$

In the weak coupling regime the starting point is $V_s^{(0)} = -C_f \frac{\alpha_s}{r}$.

Golden Mode: Bottomonium ground state (?)

In the strong coupling regime case

$$V_s^{(0)}(\mathbf{r}) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle W_{\square} \rangle \quad \text{Wilson, Susskind}$$

Motivation:

- ▶ 1st principle computation of heavy quarkonium properties from QCD
- ▶ Determination of Standard Model parameters: m_Q, α_s, \dots

We have an effective field theory, **Potential Non-Relativistic QCD**, which describes the heavy quarkonium dynamics in the weak and strong coupling situation. $m \gg mv \gg mv^2$

$$\left. \begin{array}{l} \left(i\partial_0 - \frac{\mathbf{p}^2}{2m} - V_s^{(0)}(r) \right) \Phi(\mathbf{r}) = 0 \\ + \text{corrections to the potential} \\ + \text{interaction with other low} \\ \quad \text{energy degrees of freedom} \end{array} \right\} \text{potential NRQCD} \quad E \sim mv^2$$

In the weak coupling regime the starting point is $V_s^{(0)} = -C_f \frac{\alpha_s}{r}$.

Golden Mode: Bottomonium ground state (?)

In the strong coupling regime case

$$V_s^{(0)}(\mathbf{r}) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle W_{\square} \rangle \quad \text{Wilson, Susskind}$$

Motivation:

- ▶ 1st principle computation of heavy quarkonium properties from QCD
- ▶ Determination of Standard Model parameters: m_Q, α_s, \dots

We have an effective field theory, **Potential Non-Relativistic QCD**, which describes the heavy quarkonium dynamics in the weak and strong coupling situation. $m \gg mv \gg mv^2$

$$\left. \begin{array}{l} \left(i\partial_0 - \frac{\mathbf{p}^2}{2m} - V_s^{(0)}(r) \right) \Phi(\mathbf{r}) = 0 \\ + \text{corrections to the potential} \\ + \text{interaction with other low} \\ \quad \text{energy degrees of freedom} \end{array} \right\} \text{potential NRQCD} \quad E \sim mv^2$$

In the weak coupling regime the starting point is $V_s^{(0)} = -C_f \frac{\alpha_s}{r}$.

Golden Mode: Bottomonium ground state (?)

In the strong coupling regime case

$$V_s^{(0)}(\mathbf{r}) = \lim_{T \rightarrow \infty} \frac{i}{T} \log \langle W_{\square} \rangle \quad \text{Wilson, Susskind}$$

Phenomenological analysis:



- ▶ **Which states belong to the weak/strong coupling regime?**

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, pineda; Brambilla, Garcia, Soto, Vairo; ...

Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**
To which extent the static potential can be described with perturbation theory
Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, pineda; Brambilla, Garcia, Soto, Vairo; ...



Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ **Spectroscopy at weak coupling**

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...

Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ **Spectroscopy at weak coupling**

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...

Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**



To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ **Spectroscopy at weak coupling**

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...



Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**
To which extent the static potential can be described with perturbation theory
Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...
- ▶ **Spectroscopy at weak coupling**
 $M_{\Upsilon(1S)} \rightarrow m_b(m_b) \sim 4.2$
Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...
 η_b
Sumino, Recksiegel; Kniehl, Penin, Pineda, Smirnov, Steinhauser (with Renormalization group).
Problems with the recent experimental determination:
theory ~ 40 MeV versus experiment ~ 70 MeV

Phenomenological analysis:

- ▶ Which states belong to the weak/strong coupling regime?

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ Spectroscopy at weak coupling

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...


η_b

Sumino, Recksiegel; Kniehl, Penin, Pineda, Smirnov, Steinhauser (with Renormalization group).

Problems with the recent experimental determination:
theory ~ 40 MeV versus experiment ~ 70 MeV

Kniehl, Penin, Smirnov, Steinhauser, Pineda;
 Penin, Smirnov, Steinhauser, Pineda

$$\begin{aligned} \delta E &\sim m\alpha^4 + m\alpha^5 \ln \alpha + m\alpha^6 \ln^2 \alpha + \dots \\ &+ m\alpha^5 + m\alpha^6 \ln \alpha + m\alpha^7 \ln^2 \alpha + m\alpha^8 \ln^3 \alpha + \dots \end{aligned}$$



Phenomenological analysis:

- ▶ Which states belong to the weak/strong coupling regime?

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ Spectroscopy at weak coupling

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...

η_b

Sumino, Recksiegel; Kniehl, Penin, Pineda, Smirnov, Steinhauser (with Renormalization group).

Problems with the recent experimental determination:

theory ~ 40 MeV versus experiment ~ 70 MeV

Other states

Phenomenological analysis:

- ▶ Which states belong to the weak/strong coupling regime?

To which extent the static potential can be described with perturbation theory

Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...

- ▶ Spectroscopy at weak coupling

$$M_{\Upsilon(1S)} \longrightarrow m_b(m_b) \sim 4.2$$

Beneke, Signer; Brambilla, Sumino, Vairo; Pineda; Lee; ...

η_b

Sumino, Recksiegel; Kniehl, Penin, Pineda, Smirnov, Steinhauser (with Renormalization group).

Problems with the recent experimental determination:

theory ~ 40 MeV versus experiment ~ 70 MeV

Other states

Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**
To which extent the static potential can be described with perturbation theory
Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...
- ▶ **Spectroscopy at weak coupling**
- ▶ **Coupling with EM:**
hard photons, inclusive decays of the bottomonium ground state and bottomonium sum rules at weak coupling, $t\bar{t}$.
.....

Phenomenological analysis:

- ▶ **Which states belong to the weak/strong coupling regime?**
To which extent the static potential can be described with perturbation theory
Sumino; Pineda; Sumino, Recksiegel; Lee; Bali, Pineda; Brambilla, Garcia, Soto, Vairo; ...
- ▶ **Spectroscopy at weak coupling**
- ▶ **Coupling with EM:**
hard photons, inclusive decays of the bottomonium ground state and bottomonium sum rules at weak coupling, $t\bar{t}$.
.....

Relation of the vacuum polarization and $\Gamma(V \rightarrow e^+ e^-)$

$$J^\mu = \bar{Q}\gamma^\mu Q = c_1 \psi^\dagger \sigma \chi + \dots, \quad c_1 = 1 + a_1 \alpha_s + a_2 \alpha_s^2 + \dots$$

$$(q_\mu q_\nu - g_{\mu\nu})\Pi(q^2) = i \int d^4 x e^{iqx} \langle \text{vac} | J_\mu(x) J_\nu(0) | \text{vac} \rangle$$

$$\Pi(q^2) \sim c_1^2 \langle \mathbf{r} = \mathbf{0} | \frac{1}{E - H} | \mathbf{r} = \mathbf{0} \rangle$$

$$G(\mathbf{0}, \mathbf{0}, E) = \sum_{m=0}^{\infty} \frac{|\phi_m(\mathbf{0})|^2}{E_m - E + i\epsilon} + \frac{1}{\pi} \int_0^{\infty} dE' \frac{|\phi_{E'}(\mathbf{0})|^2}{E' - E + i\epsilon}$$

$$\Gamma(V \rightarrow e^+ e^-) \sim \frac{1}{m^2} c_v^2 |\phi_n^v(\mathbf{0})|^2 \quad \Gamma(P \rightarrow \gamma\gamma) \sim \frac{1}{m^2} c_s^2 |\phi_n^s(\mathbf{0})|^2$$

$$|\phi_n^{v/s}(\mathbf{0})|^2 = |\phi_n^C(\mathbf{0})|^2 \left(1 + \delta\phi_n^{v/s}\right) = \underset{E=E_n}{\text{Res}} G(\mathbf{0}, \mathbf{0}; E),$$

where the Coulomb wave function is given by

$$|\phi_n^C(\mathbf{0})|^2 = \frac{1}{\pi} \left(\frac{m_Q C_F \alpha_s}{2n} \right)^3.$$

Note that $|\phi_n^{v/s}(\mathbf{0})|^2$ are **SCHEME** and **SCALE** dependent.

Inclusive electromagnetic decays: bottomonium




Figure: Prediction for the $\Upsilon(1S)$ decay rate to e^+e^- . We work in the RS' scheme. Pineda, Signer

The effect of the resummation of logarithms is important if compared with just keeping the single logarithm.




Figure: Prediction for the $\eta_b(1S)$ decay rate to two photons. We work in the RS' scheme.

Decay Ratio at NNLL

Penin, Smirnov, Steinhauser, Pineda

$$\frac{\Gamma(V_Q(nS) \rightarrow e^+ e^-)}{\Gamma(P_Q(nS) \rightarrow \gamma\gamma)} \sim 1 + \alpha \ln \alpha + \alpha^2 \ln^2 \alpha + \dots$$

$$+ \alpha + \alpha^2 \ln \alpha + \alpha^3 \ln^2 \alpha + \dots$$

$$+ \alpha^2 + \alpha^3 \ln \alpha + \alpha^4 \ln^2 \alpha + \dots$$


$$\left| \phi_n^{v/s}(\mathbf{0}) \right|^2 = \left| \phi_n^C(\mathbf{0}) \right|^2 \left(1 + \delta\phi_n^{v/s} \right) = \underset{E=E_n}{\text{Res}G(\mathbf{0}, \mathbf{0}; E)},$$

Note that $\delta\phi_n^{v/s}$ are **DIVERGENT**: **SCHEME** and **SCALE** dependent.


$$\frac{\Gamma(V \rightarrow e^+ e^-)}{\Gamma(P \rightarrow \gamma\gamma)} \sim \frac{C_V}{C_S}(\mu) \left(1 + \delta\phi_n^V(\mu) - \delta\phi_n^S(\mu) \right)$$

$$\sim 1 + \mathcal{O}(\alpha_s) + \mathcal{O}(\alpha_s^2) + \mathcal{O}(v^2) + \dots$$

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.659 \pm 0.089(\text{th.})_{-0.018}^{+0.019}(\delta\alpha_s) \pm 0.015(\text{exp.}) \text{ KeV},$$



The spin ratio as the function of the renormalization scale ν for the (would be) toponium ground state. The yellow band reflects the errors due to $\alpha_s(M_Z)$.



The spin ratio as the function of the renormalization scale ν for the bottomonium ground state. The yellow band reflects the errors due to $\alpha_s(M_Z)$.




Figure: The decay ratio. The horizontal band represents the experimental error of the ratio. The NNLL band reflects the errors due to $\alpha_s(M_Z) = 0.118 \pm 0.003$.

Proposal: Reorganization of perturbation theory (Kiyo, Pineda, Signer: PRELIMINARY)

Improved perturbation theory "acceleration of perturbation theory"

$$H^{(0)} = \frac{\mathbf{p}^2}{m} + V^{(0)} \longrightarrow E_n^{(0)}, \phi_n^{(0)}(\mathbf{r})$$

Keep the static potential exactly

$$V_s^{(0)} = -C_F \frac{\alpha_s(1/r)}{r} \left(1 + a_1 \frac{\alpha_s(1/r)}{4\pi} + a_2 \frac{\alpha_s^2(1/r)}{16\pi^2} + \dots \right)$$

Relativistic corrections:

$$\Delta H = \frac{V^{(1)}}{m} + \frac{V^{(2)}}{m^2} + \dots$$

$$\Gamma(V \rightarrow e^+ e^-) \sim \frac{1}{m^2} c_v^2 |\phi_n^v(\mathbf{0})|^2 + \dots \quad \Gamma(P \rightarrow \gamma\gamma) \sim \frac{1}{m^2} c_s^2 |\phi_n^s(\mathbf{0})|^2 + \dots$$

$$E_{HF} \sim c_{HF} |\phi_n^v(\mathbf{0})|^2 + \dots$$

$$\left| \phi_n^{v/s}(\mathbf{0}) \right|^2 = \left| \phi_n^{(0)}(\mathbf{0}) \right|^2 \left(1 + \delta\phi_n^{v/s} \right) = \underset{E=E_n}{\text{Res}G(\mathbf{0}, \mathbf{0}; E)},$$

HF and decays sensitive to the behavior of the wave function at the origin:



$$\left| \phi_n^C(\mathbf{0}) \right|^2 \longrightarrow \left| \phi_n^{(0)}(\mathbf{0}) \right|^2$$

- ▶ $\mathcal{O}(v^2)$ relativistic corrections beyond the Coulomb approximation.

$$\begin{aligned} \frac{\Gamma(V \rightarrow e^+ e^-)}{\Gamma(P \rightarrow \gamma\gamma)} &\sim \frac{c_v}{c_s}(\mu) \left(1 + \delta\phi_n^v(\mu) - \delta\phi_n^s(\mu) \right) \\ &\sim 1 + \mathcal{O}(\alpha_s) + \mathcal{O}(\alpha_s^2) + \mathcal{O}(v^2) + \dots \end{aligned}$$

Use of "Improved" potential \rightarrow numerical analysis (relevant for future analysis with nonperturbative potentials!!)

Regularization in position space \rightarrow change to $\overline{\text{MS}}$ (known at one loop)

RG plays an important role.

TOP: PRELIMINARY




Figure: Decay ratio.

TOP: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s)$ in the static potential.

Sizable but small correction: $\sim 1.3\%$

TOP: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^2)$ in the static potential.

Small correction: $\lesssim 1\%$

TOP: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^3)$ in the static potential.

Very small correction. Convergent series

TOP: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^4)$ in the static potential.

Very small correction. Convergent series.

BOTTOM: PRELIMINARY




Figure: Decay ratio.

Magnitude of the splitting larger than for top.

BOTTOM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s)$ in the static potential.

Sizeable correction: $\sim 10\%$

BOTTOM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^2)$ in the static potential.

Sizeable but convergent correction: $\sim 6.7\%$

BOTTOM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^3)$ in the static potential.

Sizeable but convergent correction: $\sim 3.3\%$

BOTTOM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^4)$ in the static potential.

Small correction but not very much convergent. $\mathcal{O}(\alpha_s^4)$ incomplete.

BOTTOM: PRELIMINARY

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

CHARM: PRELIMINARY




Figure: Decay ratio.

Magnitude of the corrections are large.

CHARM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s)$ in the static potential.

Sizable correction: $\sim 20\%$

CHARM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^2)$ in the static potential.

Sizable correction: $\sim 27\%$

CHARM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^3)$ in the static potential.

Sizable but convergent: $\sim 13.3\%$

CHARM: PRELIMINARY




Figure: Decay ratio. $\mathcal{O}(\alpha_s^4)$ in the static potential.

Not very convergent: $\sim 10\%$. $\mathcal{O}(\alpha_s^4)$ incomplete.
 Pattern to bring closer agreement with experiment.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime.
- ▶ Strong coupling regime.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ **Weak coupling regime**

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.
Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).

Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

CONCLUSIONS

pNRQCD: Effective field theory from QCD that describes Heavy Quarkonium.

- ▶ Weak coupling regime

Problems with hyperfine splitting and decays?

Reorganization of perturbative series may lead to a faster convergence.

Sizable effect.

Applied to decay ratio. Updated prediction (PRELIMINARY) for

$$\Gamma(\eta_b(1S) \rightarrow \gamma\gamma) = 0.544 \pm 0.146(\text{th})_{-0.004}^{+0.002}(\alpha_s) \pm 0.007(\text{exp})\text{keV},$$

Future:

Heavy Quarkonium hyperfine splitting, Decays, sum rules, Spectrum,

Application with nonperturbative potential (charm?). Assign errors to (unquenched) lattice potentials.

Wave function at the origin: relation with lattice/experimental. Determination of NRQCD matrix element (scheme and scale dependent).


Other considerations:

The hard and RG contribution are not included in non-relativistic lattice determinations of heavy quarkonium properties. Those effects are sizable.

We need not to do perturbation theory/matching in the lattice. Our approach is analytic as far as the divergences is concerned.

Bottomonium hyperfine splitting? MUCH MORE PRELIMINARY


$$E_{HF} \sim c_{HF} |\phi_n^V(\mathbf{0})|^2 + \dots$$



LO Coulomb: $c_{HF} \sim \alpha_s$ $|\phi_n^V(\mathbf{0})|^2 = |\phi_n^C(\mathbf{0})|^2 \sim m\alpha_s^3$

Bottomonium hyperfine splitting? MUCH MORE PRELIMINARY


$$E_{HF} \sim c_{HF} |\phi_n^V(\mathbf{0})|^2 + \dots$$



LO Improved: $c_{HF} \sim \alpha_s$ $|\phi_n^V(\mathbf{0})|^2 \sim 0.35$
 Result roughly equivalent to non-relativistic lattice.

Bottomonium hyperfine splitting? MUCH MORE PRELIMINARY

$$E_{HF} \sim c_{HF} |\phi_n^V(\mathbf{0})|^2 + \dots$$



LI Improved: $c_{HF} \sim \alpha_s + \alpha_s^2 \ln + \dots$ $|\phi_n^V(\mathbf{0})|^2 \sim 0.35$

This gap is not included in non-relativistic lattice.

$$\widehat{G}(E_n) \equiv \sum'_m \frac{|\psi_m^{(0)}(0)|^2}{E_m^{(0)} - E_n^{(0)}} = \lim_{E \rightarrow E_n^{(0)}} \left(G(E) - \frac{|\psi_n^{(0)}(0)|^2}{E_n^{(0)} - E} \right). \quad (1)$$

The prime indicates that the sum does not include the state n and

$$G(E) = G(0, 0; E) \equiv \lim_{r \rightarrow 0} G(r, r; E) = \lim_{r \rightarrow 0} \langle \mathbf{r} | \frac{1}{H^{(0)} - E - i0} | \mathbf{r} \rangle \quad (2)$$

$$G^{(r)}(E) = \frac{m_r}{2\pi} \left[A^{(r)}(r_0; \mu) + B_{V_s^{(0)}}^{(r)}(E; \mu) \right], \quad (3)$$

$$A^{(r)}(r_0; \mu) = \frac{u_0(r_0)}{r_0} = \frac{1}{r_0} - 2m_r C_F \alpha_s \ln(\mu e^{\gamma_E} r_0) + \mathcal{O}(\alpha_s^2), \quad (4)$$

$$\widehat{G}^{(r)}(E_n) = \frac{m_r}{2\pi} \left[A^{(r)}(r_0; \mu) + \widehat{B}_{V_s^{(0)}}^{(r)}(E_n; \mu) \right]. \quad (5)$$

$$\delta\rho_n^{\overline{\text{MS}}}(\mu) = -\frac{8m_r C_F}{3m_1 m_2} D_{S^2, s}^{(2)}(\mu) \left(\widehat{B}_{V_s^{(0)}}^{(r)}(E_n^{(0)}; \mu) + \frac{1}{3} m_r C_F \alpha_s + \mathcal{O}(\alpha_s^2) \right). \quad (6)$$