

$\mathcal{N} = (0, 2)$ Heterotic $\mathbb{C}P^{N-1}$ Sigma Models and a Nonrenormalization Theorem

Xiaoyi Cui

Department of Physics, University of Minnesota

Joint work with Mikhail Shifman

XC and M. Shifman, PRD **82**, 105022(2010), arXiv:1009.4421.

XC and M. Shifman, arXiv:1105.5107.

August 28, 2011

Heterotic nonlinear sigma models

4D gauge theories \Leftrightarrow 2D NLSM

$\mathcal{N} = 2$, $U(N)^N$ theories \Leftrightarrow $\mathcal{N} = (2, 2)$, $\mathbb{C}\mathbb{P}^{N-1}$

$\mathcal{N} = 1$ mass deformation \Leftrightarrow $\mathcal{N} = (0, 2)$ heterotic models

Objective: to understand the perturbative aspects of the deformed $\mathbb{C}\mathbb{P}^{N-1}$ models.

Heterotic nonlinear sigma models

$$\mathcal{L}_{(2,2)} = G \left\{ \partial^\mu \phi \partial_\mu \phi^\dagger + i \bar{\psi} \not{D} \psi + \text{four fermion interactions} \right\},$$

$$\mathcal{L}_{(0,2)} = \mathcal{L}_{(2,2)} + \zeta_R^\dagger i \partial_{LL} \zeta_R + \left[\gamma \zeta_R G (i \partial_{LL} \phi^\dagger) \psi_R + \text{H.c.} \right] \\ + \text{four fermion interactions.}$$

- ▶ theory with two couplings.

$$\beta_{g^2} = -\frac{Ng^4}{2\pi} + \dots, \quad \beta_\gamma = \frac{N\gamma}{2\pi}(\gamma^2 - g^2) + \dots$$

- ▶ define $\rho = \frac{\gamma^2}{g^2}$, we have

$$\beta_\rho = \frac{Ng^2}{2\pi} \rho(2\rho - 1) + \dots$$

- ▶ The factorization sustains to all loop level.

[XC and M. Shifman,

$\mathcal{N} = (0, 2)$ supersymmetry

- ▶ $\mathcal{N} = (0, 2)$ superspace:

$$\begin{array}{cccc} x_L & & x_R & \\ \theta_L & \theta_L^\dagger & \theta_R & \theta_R^\dagger \end{array} \Rightarrow \begin{array}{cc} x_L & x_R \\ \theta_R & \theta_R^\dagger \end{array}$$

- ▶ super-derivatives

$$D, \bar{D} \Rightarrow \{D, \bar{D}\} = 2i\partial_{LL}$$

∂_{RR} commutes with everything!

- ▶ chiral superfields

bosonic ϕ ψ_L

fermionic ψ_R F

A simplified model with flat target space

In a bid to understand the deformation and to get some higher loop correction, we consider the following model, which corresponds to the limiting case $\frac{\gamma^2}{g^2} \rightarrow \infty$ of the previous model.

$$\mathcal{L}_{\text{linear}} = \frac{1}{2} \int d^2\theta_R iA^\dagger \overleftrightarrow{\partial}_{RR} A + B^\dagger B + \mathcal{B}^\dagger \mathcal{B} - \left(\gamma \mathcal{B} B A^\dagger + \text{H.c.} \right).$$

- ▶ background symmetry

$$A \rightarrow A + \alpha + \alpha^\dagger A^2 \Rightarrow A \rightarrow A + \alpha.$$

- ▶ fermion flavor symmetry:

$$\left(\frac{\mathcal{B}}{1+A^\dagger A} \right) \Rightarrow \left(\frac{\mathcal{B}}{B} \right).$$


Supergraph as a probe

Feynman rules to remember:

$$T[A_z, A_{z'}^\dagger] = \begin{array}{c} p \\ \text{---} \longrightarrow \text{---} \\ z' \qquad \qquad \qquad z \end{array} = -\frac{i}{p^2} \delta^1(\theta_R - \theta'_R)$$

$$T[B_z, B_{z'}^\dagger] = \begin{array}{c} p \\ \text{---} \longrightarrow \text{---} \\ z' \qquad \qquad \qquad z \end{array} = -\frac{i}{p_L} \delta^1(\theta_R - \theta'_R)$$

► Propagators:



$$= \frac{i\gamma}{2}$$

► vertices:

► (anti)chiral projectors:


$$A \text{ --- } \begin{array}{c} p \\ \longrightarrow \\ \bullet \\ z \end{array} = \bar{D}_R(p, \theta_R, \theta_R^\dagger) A_z$$

$$A^\dagger \text{ --- } \begin{array}{c} p \\ \longrightarrow \\ \bullet \\ z \end{array} = A_z^\dagger \bar{D}_R(p, \theta_R, \theta_R^\dagger)$$


Supergraph as a probe

Feynman diagrams to calculate:

- ▶ one-loop diagrams:



- ▶ two-loop diagrams:



Finally, two loop contribution to $\beta(\gamma)$ vanishes! [XC and M. Shifman, 2011]

Background field method as a proof

Key idea: shift symmetry of θ_R^\dagger preserved classically and quantum mechanically $\Rightarrow \int d\theta_R d\theta_R^\dagger$ constant = vanishing loop correction!
Need: nontrivial background!

Upshot:

- ▶ something \bar{Q}_R -closed, ie,

$$[\bar{Q}_R, X] = 0.$$

- ▶ F term:

$$\int d^2x [Q_R, X]|_{\theta_R=\theta_R^\dagger=0} \neq 0,$$

we end up at the usual nonrenormalization theorem.

- ▶ D term:

$$\int d^2x \{Q_R, [\bar{Q}_R, X]\}|_{\theta_R=\theta_R^\dagger=0} = 0!$$

Background field method as a proof

The way out: target space symmetry helps!

- ▶ consider an equivariant version of the previous story,

$$[\bar{Q}_R, \bar{X}] = \bar{0}, \quad \int d^2x \{Q_R, [\bar{Q}_R, X]\}|_{\theta_R=\theta_R^\dagger=0} \neq 0.$$

- ▶ Target space symmetry is given by

$$A_{bk} \rightarrow A_{bk} + f(x_R), \quad A_{qu} \rightarrow A_{qu}.$$

Quantum correction is the same for all possible backgrounds in the equivalent class, but the background could be nontrivial!

quantum



background



$f(t-z)$

0

Nonperturbative regime

Following Seiberg's argument [Seiberg, 93]

Good news:

- ▶ can promote γ to a superfield \Rightarrow chirality is preserved!
- ▶ can assign R-charges!

Bad news:

- ▶ no F term, no holomorphic function!

So you will get,

$$f \left(\gamma B B A^\dagger, \frac{\gamma B B}{A}, |B|^2, |B|^2, |\gamma|^2 \right).$$

Now apply target space symmetry, the holomorphic piece will save our day. Similarly one could analyze Z_A . [XC and M. Shifman, 2011]

Conclusion

Lessons we've learned.

- ▶ Flat target space, so quantum correction is uniformal, in the sense of respecting target space symmetry.
- ▶ $\mathcal{N} = (0, 2)$ makes sure that the translation is powerful.

⇒ generalization to other linear models will be straightforward.

What about nonlinear models?

- ▶ For heterotic \mathbb{CP}^1 model, Z_A and Z_γ are free from $|\gamma|^{2n}$ -correction at n loop order.
- ▶ We shall explore the mixing contribution of g and γ at higher loop order.
- ▶ For generic non-linear $\mathcal{N} = (0, 2)$ models, can use topologically non-trivial backgrounds to obtain NRT. [XC and M.

Shifman, in progress]

Thank you!