Supersymmetry 2011 (SUSY11)



Contribution ID: 234

Type: not specified

Dark Light Higgs

Thursday, 1 September 2011 14:30 (20 minutes)

We discovered a benchmark scenario in the NMSSM which possesses novel Higgs and dark matter (DM) properties. There naturally co-exist three light singlet-like particles in this scenario: a scalar, a pseudoscalar, and their superpartner, all with masses of order 0.1-10 GeV. New non-standard decay channel is opened for the Standard Model (SM)-like Higgs boson, while its pair-decays to the light scalars or pseudoscalars are generically suppressed. This will dramatically change our search strategies for the SM-like and light Higgs bosons at colliders. In addition, this scenario is characterized by a light siglino-like DM candidate (~ 1-10 GeV). For a certain parameter window annihilation into the light pseudoscalar and exchange of the light scalar with nucleons allow the singlino to achieve the correct relic density and a (spin-independent) direct detection cross section as large as 10⁴-40} cm2 simultaneously, which is accessible to the current direct detections (e.g., CoGeNT and DAMA) or the ones in the near future.

Primary author: Dr LIU, Tao (Department of Physics, UC@Santa Barbara)

Co-authors: Prof. WAGNER, Carlos (University of Chicago); Dr ZHANG, Hao (Peking Univ.); Prof. WANG, Liantao (University of Chicago); Dr DRAPER, Patrick (UC, Santa Cruz)

Presenter: Dr LIU, Tao (Department of Physics, UC@Santa Barbara)

Session Classification: Parallel Session 9

Track Classification: SUSY: phenomenology