

# Leptogenesis in neutrinophilic Higgs doublet models

Osamu Seto (Hokkai-Gakuen Univ.)


With Naoyuki Haba (Osaka Univ.)

Refs: Prog. Theor. Phys. 125, 1155 (2011)

arXiv 1106.5354 [hep-ph]

# § Introduction

- Baryon asymmetry
- Why baryon number in our Universe is not same as anti-baryon number?



- Baryogenesis via leptogenesis

**Thermal leptogenesis**

[NASA]

# Baryon asymmetry by thermal leptogenesis

- The simplest model
- Adding right-handed neutrinos

$$\mathcal{L}_N^{\text{SM}} = -y_{ij}^\nu \overline{l_{L,i}} \Phi N_j - \frac{1}{2} \sum_i M_i \overline{N_i^c} N_i \quad m_{ij} = \sum_k \frac{y_{ik}^\nu v y_{kj}^{\nu T} v}{M_k}$$

- Scenario
- Hot Universe after inflation
- Right-handed neutrinos are thermally produced and reach thermal equilibrium
- Its out of equilibrium decay into leptons and anti-leptons
- Sphaleron transfer


# Baryon asymmetry by thermal leptogenesis

- Resultant baryon asymmetry

$$\frac{n_b}{s} \simeq C \kappa \frac{\varepsilon}{g_*}$$

- CP asymmetry  $\varepsilon \equiv \frac{\Gamma(N_1 \rightarrow \Phi + \bar{l}_j) - \Gamma(N_1 \rightarrow \Phi^* + l_j)}{\Gamma(N_1 \rightarrow \Phi + \bar{l}_j) + \Gamma(N_1 \rightarrow \Phi^* + l_j)}$
- Efficiency (dilution, washout) factor  $\kappa$
- Sphaleron transfer  $C$
- Degrees of freedom in thermal bath

# CP asymmetry




- In hierarchical right-handed neutrino mass

$$\varepsilon_1 \simeq -\frac{3}{8\pi} \frac{1}{\left(h_\nu h_\nu^\dagger\right)_{11}} \sum_{i=2,3} \text{Im} \left[ \left(h_\nu h_\nu^\dagger\right)_{1i}^2 \right] \frac{M_1}{M_i}$$


$$M_1 \gtrsim 10^9 \left( \frac{\eta_B}{5 \times 10^{-11}} \right) \left( \frac{.06 eV}{m_3} \right) \left( \frac{2 \times 10^{-4}}{n_{\nu R}/s \delta} \right) \text{ GeV}$$

- The lower bound on RH neutrino mass

# Washout



$\Delta L = 2$  scatterings




etc...

$\Delta L = 1$  scatterings

# Washout

- Efficiency (dilution, washout) factor  $\kappa$



# § Neutrinophilic Higgs doublet models [Ma, Gabriel and Nandi,...]

- Yukawa couplings

$$\mathcal{L}_{yukawa} = y^u \bar{Q}_L \Phi U_R + y^d \bar{Q}_L \tilde{\Phi} D_R + y^l \bar{L} \Phi E_R + y^\nu \bar{L} \Phi_\nu N + \frac{1}{2} M \bar{N}^c N + \text{h.c.}$$

- Higgs potential

$$V^{\text{THDM}} = m_\Phi^2 \Phi^\dagger \Phi + m_{\Phi_\nu}^2 \Phi_\nu^\dagger \Phi_\nu - m_3^2 (\Phi^\dagger \Phi_\nu + \Phi_\nu^\dagger \Phi) + \frac{\lambda_1}{2} (\Phi^\dagger \Phi)^2 + \frac{\lambda_2}{2} (\Phi_\nu^\dagger \Phi_\nu)^2 + \lambda_3 (\Phi^\dagger \Phi) (\Phi_\nu^\dagger \Phi_\nu) + \lambda_4 (\Phi^\dagger \Phi_\nu) (\Phi_\nu^\dagger \Phi) + \frac{\lambda_5}{2} [(\Phi^\dagger \Phi_\nu)^2 + (\Phi_\nu^\dagger \Phi)^2].$$



# § Neutrinophilic Higgs doublet models [Ma, Gabriel and Nandi,...]

- Yukawa couplings

$$\mathcal{L}_{yukawa} = y^u \bar{Q}_L \Phi U_R + y^d \bar{Q}_L \tilde{\Phi} D_R + y^l \bar{L} \Phi E_R + y^\nu \bar{L} \Phi_\nu N + \frac{1}{2} M \bar{N}^c N + \text{h.c.}$$

fields	$Z_2$ -parity
SM Higgs doublet, $\Phi$	+
new Higgs doublet, $\Phi_\nu$	-
right-handed neutrinos, $N$	-
others	+


soft breaking



$$m_3^2 (\Phi^\dagger \Phi_\nu + \Phi_\nu^\dagger \Phi) + \frac{\lambda_1}{2} (\Phi^\dagger \Phi)^2 + \frac{\lambda_2}{2} (\Phi_\nu^\dagger \Phi_\nu)^2$$

$$\lambda_4 (\Phi^\dagger \Phi_\nu)(\Phi_\nu^\dagger \Phi) + \frac{\lambda_5}{2} [(\Phi^\dagger \Phi_\nu)^2 + (\Phi_\nu^\dagger \Phi)^2].$$

# Higgs doublet



$v_\nu$  and Nandi, ...]

$$L + y^\nu \bar{L} \Phi_\nu N + \frac{1}{2} M \bar{N}^c N + \text{h.c.}$$

fields	$Z_2$ -parity
SM Higgs doublet, $\Phi$	+
new Higgs doublet, $\Phi_\nu$	-
right-handed neutrinos, $N$	-
others	+

soft breaking

$$m_3^2 (\Phi^\dagger \Phi_\nu + \Phi_\nu^\dagger \Phi) + \frac{\lambda_1}{2} (\Phi^\dagger \Phi)^2 + \frac{\lambda_2}{2} (\Phi_\nu^\dagger \Phi_\nu)^2$$

$$\lambda_4 (\Phi^\dagger \Phi_\nu)(\Phi_\nu^\dagger \Phi) + \frac{\lambda_5}{2} [(\Phi^\dagger \Phi_\nu)^2 + (\Phi_\nu^\dagger \Phi)^2].$$

# § § Concept of neutrinophilic Higgs doublet models

- Smallness of neutrino mass  $\frac{y_{ik}^\nu v y_{kj}^{\nu T} v}{M_k}$



tiny Yukawa, or large M

- If neutrino mass is given by  $\frac{y_{ik}^\nu v_\nu y_{kj}^{\nu T} v_\nu}{M_k}$ , the smallness is at least partially due to smallness of Higgs VEV

- $v_\nu$  ↓  $y^\nu$  ↑ and/or  $M_k$  ↓


# § Leptogenesis with neutrinoophilic Higgs

- Resultant baryon asymmetry

$$\frac{n_b}{s} \simeq C \kappa \frac{\varepsilon}{g_*}$$

- CP asymmetry : **subject to change**
- Efficiency (wash out) factor : **subject to change**
- Sphaleron transfer : **similar**
- Degrees of freedom in thermal bath: **similar**

# § § CP asymmetry (standard)




- In hierarchical right-handed neutrino mass

$$\varepsilon_1 \simeq -\frac{3}{8\pi} \frac{1}{\left(h_\nu h_\nu^\dagger\right)_{11}} \sum_{i=2,3} \text{Im} \left[ \left(h_\nu h_\nu^\dagger\right)_{1i}^2 \right] \frac{M_1}{M_i}$$

$$M_1 \gtrsim 10^9 \left( \frac{\eta_B}{5 \times 10^{-11}} \right) \left( \frac{.06 eV}{m_3} \right) \left( \frac{2 \times 10^{-4}}{n_{\nu R}/s \delta} \right) \text{ GeV}$$

- The lower bound on RH neutrino mass

# § § CP asymmetry (ν-philic)



- In hierarchical right-handed neutrino mass


$$\varepsilon \simeq -\frac{3}{8\pi} \frac{1}{(y^{\nu\dagger}y^\nu)_{11}} \left( \text{Im}(y^{\nu\dagger}y^\nu)_{12}^2 \frac{M_1}{M_2} + \text{Im}(y^{\nu\dagger}y^\nu)_{13}^2 \frac{M_1}{M_3} \right)$$

$$\simeq -\frac{3}{16\pi} 10^{-6} \left( \frac{0.1\text{GeV}}{v_\nu} \right)^2 \left( \frac{M_1}{100\text{GeV}} \right) \left( \frac{m_\nu}{0.05\text{eV}} \right) \sin \theta$$

- **Relaxed** lower bound on RH neutrino mass

# § § Washout


- $\Delta L=2$  scattering could be effective.



- Condition

$$\sum_i \left( \sum_j \frac{y_{ij}^\nu y_{ji}^{\nu\dagger} v_\nu^2}{M_j} \right)^2 < 32\pi^3 \zeta(3) \sqrt{\frac{\pi^2 g_*}{90}} \frac{v_\nu^4}{TM_P}$$

# § § Result






# § Leptogenesis in a neutrinophilic SUSY model

# § § Leptogenesis in SUSY model

- Gravitino problem



[Kawasaki et al]

Bound on N mass  
for leptogenesis

$$M_1 \gtrsim 10^9 \left( \frac{\eta_B}{5 \times 10^{-11}} \right) \left( \frac{.06 eV}{m_3} \right) \left( \frac{2 \times 10^{-4}}{n_{\nu_R}/s \delta} \right) \text{ GeV}$$

# § § SUSY $\nu$ -philic model

- Superpotential


$$\begin{aligned}
 W = & y^u \bar{Q} H_u U_R + y^d \bar{Q} H_d D_R + y^l \bar{L} H_d E_R \\
 & + y^\nu \bar{L} H_\nu N + \frac{1}{2} M N^2 \\
 & + \mu H_u H_d + \mu' H_\nu H'_\nu + \rho H_u H'_\nu + \rho' H_\nu H_d
 \end{aligned}$$

- Parity assignment


 soft breaking

fields	$Z_2$ -parity
MSSM Higgs doublets, $H_u, H_d$	+
new Higgs doublets, $H_\nu, H'_\nu$	-
right-handed neutrinos, $N$	-
others	+

# § § SUSY Result



# § Summary

We have studied thermal leptogenesis in models with a neutrinophilic Higgs field.

- **Large CP asymmetry even for a light  $N$**
- **Low scale leptogenesis**
- **$\Delta L=2$  washout also enhanced**

**A solution to realize thermal leptogenesis in supergravity without gravitino problem**

