



Contribution ID: 162

Type: not specified

## A multi-step Fitting Approach to the allowed Parameter Space of the Constrained Minimal Supersymmetric Model and Comparison with the newest LHC Results

*Tuesday, August 30, 2011 3:00 PM (20 minutes)*

We study the strong correlations between the parameters of the popular Constrained Minimal Supersymmetric Model (CMSSM) with four independent parameters. Knowing these correlations allows to develop a multi-step fitting procedure, which efficiently determines the allowed parameter space by simply minimizing the  $\chi^2$  in multiple steps, i.e. first the parameters with the highest correlation are determined. The data included in the fit are the relic density, the b-physics observables, the anomalous magnetic moment of the muon, upper limits on direct neutralino-nucleon scattering cross sections and lower limits on Higgs and sparticle masses from LEP and newest LHC data.

With this new fitting method and including non-Gaussian systematic errors we find a significantly larger allowed parameter region than previous methods based on random or Markov Chain sampling of the parameter space.

**Primary authors:** Ms BESKIDT, Conny (IEKP, KIT); Prof. KAZAKOV, Dmitri (Dubna, JINR & Moscow, ITEP); Dr ZIEBARTH, Eva (IEKP, KIT); Dr ZHUKOV, Valery (IEKP, KIT); Prof. DE BOER, Wim (IEKP, KIT)

**Presenter:** Ms BESKIDT, Conny (IEKP, KIT)

**Session Classification:** Parallel Session 6

**Track Classification:** SUSY: models