

# Probing a minimal $U(1)_X$ model at future $e^-e^+$ collider via the fermion pair production channel

Based on : 2104.10902

In collaboration with P. S. Bhupal Dev  
Yutaka Hosotani  
Sanjoy Mandal



Arindam Das


Hokkaido University  
WIN2021 June, 2021

# Introduction

Over the decades experiments have found each and every missing pieces

Verified the facts that they belong to this family

Finally at the Large Hadron collider Higgs has been observed  
 → Its properties must be verified



Strongly established with interesting shortcomings  
 Few of the very interesting anomalies :

Tiny neutrino mass and flavor mixings  
 Relic abundance of dark matter...

Neutrino oscillation experiment : SNO, Super - K, etc .

- Nature : Majorana/ Dirac
- Ordering : Normal/Inverted
- Nature of the mixing between the mass and the flavor eigenstates

Unkown ←

SM can not explain them

# Particle Content

Dobrescu, Fox; Cox, Han, Yanagida; AD, Okada, Raut; AD, Dev, Okada;  
Chiang, Cottin, AD, Mandal; AD, Takahashi, Oda, Okada

	SU(3) <sub>c</sub>	SU(2) <sub>L</sub>	U(1) <sub>Y</sub>		U(1) <sub>X</sub>
$q_L^i$	<b>3</b>	<b>2</b>	+1/6	$x_q$	$= \frac{1}{6}x_H + \frac{1}{3}x_\Phi$
$u_R^i$	<b>3</b>	<b>1</b>	+2/3	$x_u$	$= \frac{2}{3}x_H + \frac{1}{3}x_\Phi$
$d_R^i$	<b>3</b>	<b>1</b>	-1/3	$x_d$	$= -\frac{1}{3}x_H + \frac{1}{3}x_\Phi$
$\ell_L^i$	<b>1</b>	<b>2</b>	-1/2	$x_\ell$	$= -\frac{1}{2}x_H - x_\Phi$
$e_R^i$	<b>1</b>	<b>1</b>	-1	$x_e$	$= -x_H - x_\Phi$
$H$	<b>1</b>	<b>2</b>	+1/2	$x'_H$	$= \frac{1}{2}x_H$
$N_R^i$	<b>1</b>	<b>1</b>	0	$x_\nu$	$= -x_\Phi$
$\Phi$	<b>1</b>	<b>1</b>	0	$x'_\Phi$	$= 2x_\Phi$

$$m_{Z'} = 2 g' v_\Phi$$

$x_H, x_\Phi$  will appear the coupling with  $Z'$

3 generations of SM singlet right handed neutrinos (anomaly free)

Charges before the anomaly cancellations

Charges after imposing the anomaly cancellations

$U(1)_X$  breaking

$$\mathcal{L}_Y \supset - \sum_{i,j=1}^3 Y_D^{ij} \bar{\ell}_L^i H N_R^j - \frac{1}{2} \sum_{i=k}^3 Y_N^k \Phi \overline{N_R^k} N_R^k + \text{h.c.},$$

$$m_D^{ij} = \frac{Y_D^{ij}}{\sqrt{2}} v_h$$

$$m_{N^i} = \frac{Y_N^i}{\sqrt{2}} v_\Phi$$

$$m_\nu = \begin{pmatrix} 0 & M_D \\ M_D^T & M_N \end{pmatrix} \quad m_\nu \simeq -M_D M_N^{-1} M_D^T$$

Seesaw mechanism

Higgs potential

$$V = m_h^2(H^\dagger H) + \lambda(H^\dagger H)^2 + m_\Phi^2(\Phi^\dagger \Phi) + \lambda_\Phi(\Phi^\dagger \Phi)^2 + \lambda'(H^\dagger H)(\Phi^\dagger \Phi)$$

U(1)<sub>X</sub> breaking

Electroweak breaking

$$\langle \Phi \rangle = \frac{v_\Phi + \phi}{\sqrt{2}} \quad \langle H \rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} v+h \\ 0 \end{pmatrix} \quad v \simeq 246 \text{ GeV}, v_\Phi \gg v_h$$

Mass of the neutral gauge boson **Z'**

$$M_{Z'} = g' \sqrt{4v_\Phi^2 + \frac{1}{4}x_H^2 v_h^2} \simeq 2g'v_\Phi.$$

Neutrino mass

$$\mathcal{L}^{\text{mass}} = -Y_\nu^{\alpha\beta} \bar{\ell}_L^\alpha H N_R^\beta - Y_N^\alpha \Phi \bar{N}_R^{\alpha c} N_R^\alpha + \text{h.c.}$$

$$m_{N_\alpha} = \frac{Y_N^\alpha}{\sqrt{2}} v_\Phi, \quad m_D^{\alpha\beta} = \frac{Y_\nu^{\alpha\beta}}{\sqrt{2}} v. \quad m_\nu^{\text{mass}} = \begin{pmatrix} 0 & m_D \\ m_D^T & m_N \end{pmatrix} \quad m_\nu \simeq -m_D m_N^{-1} m_D^T$$

**seesaw**

## Z' interactions

Interaction between the quarks and Z'  $\mathcal{L}^q = -g'(\bar{q}\gamma_\mu q_{x_L}^q P_L q + \bar{q}\gamma_\mu q_{x_R}^q P_R q)Z'_\mu$

Interaction between the leptons and Z'  $\mathcal{L}^\ell = -g'(\bar{\ell}\gamma_\mu q_{x_L}^\ell P_L \ell + \bar{e}\gamma_\mu q_{x_R}^\ell P_R e)Z'_\mu$

## Partial decay width

Charged fermions  $\Gamma(Z' \rightarrow 2f) = N_c \frac{M_{Z'}}{24\pi} \left( g_L^f [g', x_H, x_\Phi]^2 + g_R^f [g', x_H, x_\Phi]^2 \right)$

light neutrinos  $\Gamma(Z' \rightarrow 2\nu) = \frac{M_{Z'}}{24\pi} g_L^\nu [g', x_H, x_\Phi]^2$

heavy neutrinos  $\Gamma(Z' \rightarrow 2N) = \frac{M_{Z'}}{24\pi} g_R^N [g', x_\Phi]^2 \left( 1 - 4 \frac{m_N^2}{M_{Z'}^2} \right)^{\frac{3}{2}}$

# Implications of the choices of $x_H$ keeping $x_\Phi = 1$

No interaction with  $e_R$

No interaction with  $d_R$

	SU(3) <sub>c</sub>	SU(2) <sub>L</sub>	U(1) <sub>Y</sub>	U(1) <sub>X</sub>	-2	-1	-0.5	0	0.5	1	2
					U(1) <sub>R</sub>			B-L			
$q_L^i$	<b>3</b>	<b>2</b>	$\frac{1}{6}$	$x'_q = \frac{1}{6}x_H + \frac{1}{3}x_\Phi$	0	$\frac{1}{6}$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{5}{12}$	$\frac{1}{2}$	$\frac{1}{3}$
$u_R^i$	<b>3</b>	<b>1</b>	$\frac{2}{3}$	$x'_u = \frac{2}{3}x_H + \frac{1}{3}x_\Phi$	-1	$-\frac{1}{3}$	0	$\frac{1}{3}$	$\frac{1}{2}$	1	$\frac{5}{3}$
$d_R^i$	<b>3</b>	<b>1</b>	$-\frac{1}{3}$	$x'_d = -\frac{1}{3}x_H + \frac{1}{3}x_\Phi$	1	$\frac{2}{3}$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{1}{6}$	0	$-\frac{1}{3}$
$\ell_L^i$	<b>1</b>	<b>2</b>	$-\frac{1}{2}$	$x'_\ell = -\frac{1}{2}x_H - x_\Phi$	0	$-\frac{1}{2}$	$-\frac{3}{4}$	-1	$\frac{5}{4}$	$-\frac{3}{2}$	-2
$e_R^i$	<b>1</b>	<b>1</b>	-1	$x'_e = -x_H - x_\Phi$	1	0	$-\frac{1}{2}$	-1	$-\frac{3}{2}$	-2	-3
$N_R^i$	<b>1</b>	<b>1</b>	0	$x'_\nu = -x_\Phi$	-1	-1	-1	-1	-1	-1	-1
$H$	<b>1</b>	<b>2</b>	$-\frac{1}{2}$	$-\frac{x_H}{2} = -\frac{x_H}{2}$	1	$\frac{1}{2}$	$\frac{1}{2}$	0	$\frac{1}{4}$	$\frac{1}{4}$	1
$\Phi$	<b>1</b>	<b>1</b>	0	$2x_\Phi = 2x_\Phi$	2	2	2	2	2	2	2

6


No interaction with left handed fermions

No interaction with  $u_R$


# Phenomenological aspects of the model

**New particles**  $Z'$  boson Heavy Majorana Neutrino  $U(1)_X$  Higgs boson  
**Phenomenology**  $Z'$  boson production and decay Heavy neutrino production  
 Dark Matter collider  $U(1)_X$  Higgs phenomenology : Vacuum Stability  
 Leptogenesis and many more

## Fermionic pair production from the $Z'$



$V_i = \{\gamma, Z, Z'\}$



Limits on the model parameters

Considering the limit  $M_{Z'} \gg \sqrt{s}$  and applying effective theory we find the limits on  $\frac{M_{Z'}}{g'}$  using **LEP – II (1302.3415)** and **(prospective) ILC (1908.11299)** :


$$\frac{\pm 4\pi}{(1 + \delta_{ef})(\Lambda_{AB}^{f\pm})^2} (\bar{e}\gamma_\mu P_A e) (\bar{f}\gamma_\mu P_B f)$$

$Z'$  exchange matrix element for our process

$$\frac{(g')^2}{M_{Z'}^2 - s} [\bar{e}\gamma_\mu (x_{e'} P_L + x_{e'} P_R) e] [\bar{f}\gamma_\mu (x_{f_L} P_L + x_{f_R} P_R) f]$$

Matching the above equations we obtain

$$M_{Z'}^2 - s \geq \frac{g'^2}{4\pi} |x_{e_A} x_{f_B}| (\Lambda_{AB}^{f\pm})^2$$







Indicates a large VEV scale can be probed from LEP – II to ILC1000 via ILC250 and ILC500

Shows limits on  $M_{Z'}$  vs  $g'$  for LEP – II, ILC250, ILC500 and ILC1000

Limits on  $M_{Z'}$  and  $g'$  can also be obtained from dilepton and dijet searches at the LHC




$$g' = \sqrt{g_{\text{Model}}^2 \left( \frac{\sigma_{\text{ATLAS}}^{\text{Obs.}}}{\sigma_{\text{Model}}} \right)}$$





For heavier  $Z'$ , the limits from  $e^-e^+$  colliders are stronger than the current LHC results




$x_H > 0$



For heavier  $Z'$ , the limits from  $e^-e^+$  colliders are stronger than the current LHC results




$$e^-e^+ \rightarrow \mu^+\mu^-$$

$M_{Z'} = 7.5 \text{ TeV}$




$$e^-e^+ \rightarrow e^+e^-$$

Deviations in total cross sections from SM is more than 100% for  $x_H \geq 1$  for  $\sqrt{s} = 3 \text{ TeV}$ . For  $\sqrt{s} < 3 \text{ TeV}$  the deviation is also sizable.



# Integrated Forward – Backward Asymmetry ( $e^-e^+ \rightarrow \mu^-\mu^+$ ) : $\mathcal{A}_{FB}$

$M_{Z'} = 7.5$  TeV








Integrated

$$\mathcal{A}_{FB}(P_{e^-}, P_{e^+}) = \frac{\sigma_F(P_{e^-}, P_{e^+}) - \sigma_B(P_{e^-}, P_{e^+})}{\sigma_F(P_{e^-}, P_{e^+}) + \sigma_B(P_{e^-}, P_{e^+})}$$

Deviation from the SM

$$\Delta_{\mathcal{A}_{FB}} = \frac{\mathcal{A}_{FB}^{U(1)X}}{\mathcal{A}_{FB}^{SM}} - 1.$$

$x_H = 2$  : 3.8 % for  $P_{e^-} = -0.8$  at 500 GeV  
 $x_H = 1$  : 79 % for  $P_{e^-} = -0.8$  at 1 TeV  
 $x_H = -1$  : 20 % for  $P_{e^-} = 0.3$  at 3 TeV



Statistical error

$$\Delta_{\mathcal{A}_{FB}} = 2 \frac{\sqrt{n_1 n_2} (\sqrt{n_1} + \sqrt{n_2})}{(n_1 + n_2)^2} = \frac{2\sqrt{n_1 n_2}}{(n_1 + n_2) (\sqrt{n_1} - \sqrt{n_2})} \mathcal{A}_{FB}$$

$$(n_1, n_2) = (N_F, N_B) \quad N_{F(B)} = L_{\text{int}} \sigma_{F(B)}(P_{e^-}, P_{e^+})$$

# Differential and integrated Left – Right Asymmetry ( $e^-e^+ \rightarrow \mu^-\mu^+$ ) : $\mathcal{A}_{LR}$ $M_{Z'} = 7.5$ TeV








## Differential

$$\mathcal{A}_{LR}(\cos\theta) = \frac{\frac{d\sigma_{LR}}{d\cos\theta}(\cos\theta) - \frac{d\sigma_{RL}}{d\cos\theta}(\cos\theta)}{\frac{d\sigma_{LR}}{d\cos\theta}(\cos\theta) + \frac{d\sigma_{RL}}{d\cos\theta}(\cos\theta)}$$

## Deviation from the SM



$$\Delta_{\mathcal{A}_{LR}}(\cos\theta) = \frac{\mathcal{A}_{LR}^{U(1)X}(\cos\theta)}{\mathcal{A}_{LR}^{\text{SM}}(\cos\theta)} - 1$$

$x_H = 2$  : 10 % for at 250 GeV  
 $x_H = 1$  : 20 % for at 500 GeV  
 $x_H = -2$  : 8 % for at 500 GeV



## Integral

$$\mathcal{A}_{LR} = \frac{\sigma^{LR} - \sigma^{RL}}{\sigma^{LR} + \sigma^{RL}}$$



# Differential Left – Right, Forward – Backward Asymmetry ( $e^-e^+ \rightarrow \mu^-\mu^+$ ) : $\mathcal{A}_{LR,FB}$

$M_{Z'} = 7.5 \text{ TeV}$






## Differential

$$\mathcal{A}_{LR,FB}(\cos\theta) = \frac{[\sigma_{LR}(\cos\theta) - \sigma_{RL}(\cos\theta)] - [\sigma_{LR}(-\cos\theta) - \sigma_{RL}(-\cos\theta)]}{[\sigma_{LR}(\cos\theta) + \sigma_{RL}(\cos\theta)] + [\sigma_{LR}(-\cos\theta) + \sigma_{RL}(-\cos\theta)]}$$

## Deviation from the SM

$$\Delta_{\mathcal{A}_{LR,FB}}(\cos\theta) = \frac{\mathcal{A}_{LR,FB}^{U(1)X}(\cos\theta)}{\mathcal{A}_{LR,FB}^{SM}(\cos\theta)} - 1$$

$x_H = 2$  : 8.2 % for at 250 GeV  
 $x_H = 1$  : 15 % for at 500 GeV  
 $x_H = -2$  : 7.5 % for at 500 GeV



## Statistical error

$$\Delta \mathcal{A}_{LR,FB} = 2 \frac{(n_3 + n_2)(\sqrt{n_1} + \sqrt{n_4}) + (n_1 + n_4)(\sqrt{n_3} + \sqrt{n_2})}{(n_1 + n_4)^2 - (n_3 + n_2)^2} \mathcal{A}_{LR,FB}$$

## Conclusions :

We are looking for a scenario where which can explain a variety of beyond the SM sceanrios .

The proposal for the generation of the tiny neutrino mass, from the seesaw mechanism, under investigation at the energy frontier .

We study  $\mathcal{A}_{\text{FB}}$ ,  $\mathcal{A}_{\text{LR}}$ ,  $\mathcal{A}_{\text{LR, FB}}$  . The asymmetries are sizable at the 250 GeV and 500 GeV  $e^-e^+$  colliders or higher in the near future .

Such a model can be studied at muon colliders with high CM energy . This allows us to probe heavier  $Z'$  .

For more detail including the Bhabha scattering at  $e^-e^+$  colliders see 2104.10902

The motovation of this work is to find a new particle and/or a new force carrier as a part of the of the new physics searches including a variety of BSM aspects . 15

# Thank You



आदित्यवर्णं तमसः परस्तात्

Effulgent as the sun beyond darkness

(YV 31.18)