

Top Quark Physics at LHeC and FCC-eh

Christian Schwanenberger
DESY



for the LHeC/FCC-eh Study Group

**Snowmass EF03 Meeting
Heavy Flavour and Top**

ONLINE

5 November 2020



Outline

Introduction
CC Top Production
NC Top Production
BSM Top Production
Conclusions

Outline

Introduction

CC Top Production

NC Top Production

BSM Top Production

Conclusions

Linac–Ring Collider, LHeC and FCC–eh



LHeC

LHeC CDRs, arXiv:1206.2913, arXiv:2007.14491

operated **synchronously**

- with HL-LHC:
p beam: 7 TeV, $\sqrt{s}=1.3$ TeV
- with HE-LHC:
p beam: 13.5 TeV, $\sqrt{s}=1.8$ TeV
- or later with LE-FCC-hh:
p beam: 19 TeV, $\sqrt{s}=2.1$ TeV
- or later with FCC-hh:
p beam: 50 TeV, $\sqrt{s}=3.5$ TeV


Energy Recovering Linac

e^\pm beam: 60 GeV


$L_{\text{int}} = 1-2 \text{ ab}^{-1}$ (1-2k \times HERA!)

FCC CDR, Eur. Phys. J. C 79, no. 6, 474 (2019)

FCC-eh




LHeC and FCC-eh Detector Layout



LHeC


FCC-eh



DELPHES

SM Top Quark Production

CC DIS top production



$$\sigma = 1.89 \text{ pb @}$$

LHeC

$$\sigma = 4.46 \text{ pb @}$$


HE-LHC

$$\sigma = 15.3 \text{ pb @}$$

FCC-eh

$$E_e = 60 \text{ GeV}$$

NC top photoproduction



$$\sigma = 0.05 \text{ pb @}$$

LHeC

$$\sigma = 0.?? \text{ pb @}$$

HE-LHC

$$\sigma = 1.14 \text{ pb @}$$

FCC-eh

→ future ep collider is ideal to study EWK interactions of the top quark

Outline

Introduction

CC Top Production

NC Top Production

BSM Top Production

Conclusions


Signal and Backgrounds

Dutta, Goyal, Kumar, Mellado, Eur. Phys. J. C75 (2015) no.12, 577

signal

$$\cancel{E}_T \geq 25 \text{ GeV}$$

hadronic



$$\Delta\Phi_{\cancel{E},j} \geq 0.4$$

$$\Delta\Phi_{\cancel{E},b} \geq 0.4$$

$$|m_{j_1 j_2} - m_W| \leq 22 \text{ GeV}$$


$$p_{T,j,b} \geq 20 \text{ GeV}$$

$$|\eta_j| \leq 5, |\eta_b| \leq 2.5$$

$$\Delta R_{j,b/j} \geq 0.4$$

$N_t = 220k, s/b = 1.2$


background



e beam: 60 GeV
1000 fb⁻¹

leptonic

$$\cancel{E}_T \geq 25 \text{ GeV}$$



$$\Delta\Phi_{\cancel{E},j} \geq 0.4$$

$$\Delta\Phi_{\cancel{E},b} \geq 0.4$$


$$\Delta\Phi_{\cancel{E},l} \geq 0.4$$

$$p_{T,j,b,l} \geq 20 \text{ GeV}$$

$$|\eta_j| \leq 5, |\eta_{b,l}| \leq 2.5$$


$$\Delta R_{j,b/j} \geq 0.4$$

$N_t = 110k, s/b = 11$



→ top quark factory (with low backgrounds)

Direct Measurement of $|V_{tb}|$




$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$

Limits on Anomalous Wtb Couplings

= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$




$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$


Limits on Anomalous Wtb Couplings

= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$



$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$



+ other variables sensitive on W helicity


Limits on Anomalous Wtb Couplings

= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L - f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L - f_T^R P_R) t W_\mu^- + h.c.$$

Dutta, Goyal, Kumar, Mellado, arXiv:1307.1688
Kumar, Ruan, to be publ.



95% C.L.



DELPHES

LHeC


hadronic channel:



including detector simulation (Delphes)

Direct Measurement of $|V_{tb}|$

- ¹ including top-quark mass uncertainty
- ² σ_{theo} : NLO PDF4LHC11 (NPPS205 (2010) 10, CPC191 (2015) 74)
- ³ including beam energy uncertainty



$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$

ATLAS+CMS Preliminary
LHCtopWG

$|f_{LV}V_{tb}| = \sqrt{\frac{\sigma_{\text{meas}}}{\sigma_{\text{theo}}}}$ from single top quark production September 2019

σ_{theo} : NLO+NNLL MSTW2008nnlo
 PRD 83 (2011) 091503, PRD 82 (2010) 054018, PRD 81 (2010) 054028 total theo

$\Delta\sigma_{\text{theo}}$: scale \oplus PDF
 $m_{\text{top}} = 172.5 \text{ GeV}$

t-channel:

ATLAS+CMS combination 7+8 TeV^{1,3}
 JHEP 05 (2019) 088

CMS 13 TeV²
 arXiv:1812.10514 (35.9 fb⁻¹)

ATLAS 13 TeV²
 JHEP 04 (2017) 086 (3.2 fb⁻¹)

tW:

ATLAS+CMS combination 7+8 TeV^{1,3}
 JHEP 05 (2019) 088

ATLAS 13 TeV²
 JHEP 01 (2018) 63 (3.2 fb⁻¹)

CMS 13 TeV
 JHEP 10 (2018) 117 (35.9 fb⁻¹)


s-channel:

ATLAS+CMS combination 8 TeV^{1,3}
 JHEP 05 (2019) 088

all channels:


ATLAS+CMS combination 7+8 TeV^{1,3}
 JHEP 05 (2019) 088

LHeC 100 fb⁻¹



Top Quark Polarisation

Atag, Sahin,
PRD 73, 074001 (2006)



$\cos\theta$: angle between charged lepton and spin quantisation axis in top rest frame

$$\frac{1}{\Gamma_T} \frac{d\Gamma}{d\cos\theta} = \frac{1}{2} (1 + A_{\uparrow\downarrow} \alpha \cos\theta) \quad A_{\uparrow\downarrow} = \frac{N_{\uparrow} - N_{\downarrow}}{N_{\uparrow} + N_{\downarrow}}$$

using simply e-beam axis:
polarisation: $P_t = 96\%$

TESLA+HERAp:


$\sqrt{s} = 1.6 \text{ TeV}$

$L_{\text{int}} = 20 \text{ fb}^{-1}$



$19.7 \text{ fb}^{-1}: A_{\uparrow\downarrow} = 0.26 \pm 0.11$

JHEP 04 (2016) 073




Measurement of $|V_{td}|$

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ \mathbf{V_{td}} & V_{ts} & V_{tb} \end{pmatrix}$$

arXiv:1709.07887
LHC, 3000 fb⁻¹@14TeV

HL-LHC




DELPHES

arXiv:1501.05013

H. Sun PoS DIS 2018, 167 (2018) SM

→ extend HL-LHC limits




7TeV⊕60GeV@LHeC
P_e=0.8
5% systematic
2σ (95%C.L.) limit

2σ

arXiv:1709.07887

LHC

→ $|V_{td}| < 0.06$



50TeV⊕60GeV@FCCeh
P_e=0.8
5% systematic
2σ (95%C.L.) limit

LHC


HL-LHC

→ $|V_{td}| < 0.037$



Measurement of $|V_{ts}|$

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & \mathbf{V_{ts}} & V_{tb} \end{pmatrix}$$




DELPHES

arXiv:1501.05013

H. Sun PoS DIS 2018, 167 (2018) SM

→ probing SM prediction directly for the first time




2σ

LHC


→ $|V_{ts}| < 0.06$

FCC CDR, Eur. Phys. J. C 79, no. 6, 474 (2019)




→ $|V_{ts}| < 0.037$

CP Nature of Top-Higgs Coupling




$$\mathcal{L} = -\frac{m_t}{v} \bar{t} [\kappa \cos \zeta_t + i\gamma_5 \sin \zeta_t] t h$$




LHeC


rapidity difference (H, \bar{t})




top polarisation




angular asymmetries (b₂, l⁻)



CP Nature of Top-Higgs Coupling






$$\mathcal{L} = -\frac{m_t}{v} \bar{t} [\kappa \cos \zeta_t + i\gamma_5 \sin \zeta_t] t h$$



rapidity difference (H, \bar{t})

top polarisation

fiducial incl. cross-section



Exclusion Contours (fiducial cross section)


$$\mathcal{L} = -\frac{m_t}{v} \bar{t} [\kappa \cos \zeta_t + i\gamma_5 \sin \zeta_t] t h$$

LHeC

CP-even
(flipped sign)

CP-odd

CP-even
(SM)



→ powerful probe
of ttH coupling

10% uncertainty on
background yields

$$\kappa = 1.00 \pm 0.17$$


Outline

Introduction
CC Top Production
NC Top Production
BSM Top Production
Conclusions


Analysis of the $t\bar{t}\gamma$ Vertex

signal


background




→ measure top quark charge



LHC




OR



?

→ difficult at the LHC

Search for Anomalous $t\bar{t}\gamma$ Couplings



$\tilde{\kappa}, \kappa$

LHeC


8% and 16% accuracy
 10% 18%
 → systematically limited



27% accuracy
 (4.59fb⁻¹, 7 TeV)

$$\mathcal{L}_{t\bar{t}\gamma} = e\bar{t} \left(Q_t \gamma^\mu A_\mu + \frac{1}{4m_t} \sigma^{\mu\nu} F_{\mu\nu} (\kappa + i\tilde{\kappa}\gamma_5) \right) t$$

electric dipole moment: $\tilde{\kappa}$




magnetic dipole moment: κ

Bouzas, Larios,
 Physical Review D 88, 094007 (2013)

Search for Anomalous $t\bar{t}Z$ Couplings

Bouzas, Larios,
Physical Review D 88, 094007 (2013)





$t\bar{t}$ photoproduction

property	precision
EDM: $\tilde{\kappa} / \tilde{\kappa}_Z$	0.20-0.28/0.6-0.8
MDM: κ / κ_Z	0.05-0.09/0.9-1.3


LHeC

10% and 18% accuracy



Search for Anomalous $t\bar{t}Z$ Couplings

Bouzas, Larios,
Physical Review D 88, 094007 (2013)





$t\bar{t}$ photoproduction

$t\bar{t}\gamma$

property	precision
EDM: $\tilde{\kappa} / \tilde{\kappa}_Z$	0.20-0.28, 0.6-0.8
MDM: κ / κ_Z	0.05-0.09, 0.9-1.3

LHeC

10% and 18% accuracy



Outline


Introduction
CC Top Production
NC Top Production
BSM Top Production
Conclusions

Search for Anomalous FCNC $t\bar{u}\gamma$, $t\bar{u}Z$ Couplings


signal

I. Cakir, Yilmaz, Denizli, Senol, Karadeniz, O. Cakir, Adv. High Energy Phys. 2017, 1572053 (2017)

$$L = -g_e \sum_{q=u,c} Q_q \frac{\lambda^q}{\Lambda} \bar{t} \sigma^{\mu\nu} (f_q + h_q \gamma_5) q A_{\mu\nu} + h.c.$$



background




Search for Anomalous FCNC $t\bar{u}\gamma$ Coupling

I. Cakir, Yilmaz, Denizli, Senol,
Karadeniz, O. Cakir, Adv. High Energy
Phys. 2017, 1572053 (2017)


signal

$$L = -g_e \sum_{q=u,c} Q_q \frac{\lambda_q}{\Lambda} \bar{t} \sigma^{\mu\nu} (f_q + h_q \gamma_5) q A_{\mu\nu} + h.c.$$



$130 < M_{Wb} < 190$ GeV

$50 < M_{jj} < 100$ GeV




Search for Anomalous FCNC $tq\gamma$ Coupling



I. Cakir, Yilmaz, Denizli, Senol, Karadeniz, O. Cakir, Adv. High Energy Phys. 2017, 1572053 (2017)

signal

$$L = -g_e \sum_{q=u,c} Q_q \frac{\lambda_q}{\Lambda} \bar{t} \sigma^{\mu\nu} (f_q + h_q \gamma_5) q A_{\mu\nu} + h.c.$$



→ test exotic models leading to FCNC




Search for Anomalous FCNC $t \rightarrow q \gamma$ Coupling

signal

I. Cakir, Yilmaz, Denizli, Senol, Karadeniz, O. Cakir, Adv. High Energy Phys. 2017, 1572053 (2017)



$$L = -g_e \sum_{q=u,c} Q_q \frac{\lambda_q}{\Lambda} \bar{t} \sigma^{\mu\nu} (f_q + h_q \gamma_5) q A_{\mu\nu} + h.c.$$



$130 < M_{Wb} < 190 \text{ GeV}$

$50 < M_{jj} < 100 \text{ GeV}$

→ test exotic models leading to FCNC




Search for Anomalous FCNC $t \rightarrow q$ Coupling



signal

I. Cakir, Yilmaz, Denizli, Senol, Karadeniz, O. Cakir, Adv. High Energy Phys. 2017, 1572053 (2017)

$$L = -g_e \sum_{q=u,c} Q_q \frac{\lambda_q}{\Lambda} \bar{t} \sigma^{\mu\nu} (f_q + h_q \gamma_5) q A_{\mu\nu} + h.c.$$




→ test exotic models leading to FCNC



Search for Anomalous FCNC tHu Coupling

signal

Sun, Wang,
arXiv:1602.04670 [hep-ph]



$e^- p \rightarrow \nu_e \bar{t} \rightarrow \nu_e h \bar{q} \rightarrow \nu_e b \bar{b} \bar{q}, \quad q = u, c$


$$\mathcal{L} = \kappa_{tuh} \bar{t} u h + \kappa_{tch} \bar{t} c h + \text{h.c.}$$

Search for Anomalous FCNC tHu Coupling




signal

Sun, Wang,
arXiv:1602.04670 [hep-ph]

background



$$\mathcal{L} = \kappa_{tuh} \bar{t} u h + \kappa_{tch} \bar{t} c h + h.c.$$




Search for Anomalous FCNC tHu Coupling




signal

Sun, Wang,
arXiv:1602.04670 [hep-ph]

background



$$\mathcal{L} = \kappa_{tuh} \bar{t} u h + \kappa_{tch} \bar{t} c h + \text{h.c.}$$



- parametrised assumed resolutions for electrons/photons, muons, jets and unclustered energy using ATLAS values
- b-tag rate of 60%, c-jet fake rate of 10%, light-jet fake rate of 1%
- selections optimized for LHeC and FCC-ep scenarios ($s/\sqrt{(S+B)}$)
- cut-based and MVA-based analyses


Upper Limit on $\text{Br}(t \rightarrow uH)$ in MVA analysis

Sun, Wang,
arXiv:1602.04670 [hep-ph]


parametrisation

HL-LHC

HL-LHC



LHC, 3000 fb^{-1} @14TeV
 < 0.0015
 (1 ab^{-1})
 2σ



LHC, 3000 fb^{-1} @14TeV
 < 0.0002 (2 ab^{-1})
 2σ
 → improves HL-LHC sensitivity





FCNC Top Quark Couplings

FCC CDR, Eur. Phys. J. C 79, no. 6, 474 (2019)



~2030's ≥2050's

LHeC

FCC-ep




→ complementarity of colliders



CLIC FCC-ee

~2040's

≥2050's FCC-pp



Outline

Introduction
CC Top Production
NC Top Production
BSM Top Production
Conclusions





Summary

- future ep collider has a rich analysis programme, in particular for electroweak interactions of top quark
 - single top quark factory: $|V_{tb}|$ ($\sim 1\%$)
 - top quark couplings to bosons (Wtb , $t\bar{t}\gamma$, $t\bar{t}Z$, $t\bar{t}H$, FCNC)
 - analyse top quark properties with high precision: polarisation, charge, PDFs of tops, ...
 - many stringent searches for new physics: anomalous couplings, FCNC, CP violation in $t\bar{t}H$, heavy top, SUSY stops...
- DIS is **competitive** and **complementary** in performing **high precision** measurements of top properties
- large sensitivities to discover new physics!
- FCC-eh (2050+) sensitivities ~ 10 times better than LHeC (2030s) – maybe room for both if LINAC is reused



Backup

FCNC Top Quark Couplings

LHeC




→ complementarity of colliders



CLIC



HL-LHC



LHeC and FCC-eh Detector Layout

FCC-eh

LHeC



Central Tracker
 circular-elliptical beam pipe
 4 layers Si-pixel
 5 layers Si-strixel
 (see Table of Detector Dimensions/
 Parameters in backup)


Length of Inner Solenoid ~12m

DELPHES


Top Quark Parton Density Function

LHeC CDR, J.Phys. G39, 075001 (2012)

parton momentum fraction



six-flavour
variable number
scheme for top
quark





→ LHeC offers new field of research for top quark PDF

Top Quark Structure Function

Boroun, Phys. Lett. B744, 142 (2015)

$L_{int} = 10 \text{ fb}^{-1}$

variable flavour
number scheme
for top quark



$$\tau_t = \left(1 + \frac{4m_t^2}{Q^2}\right)^{1+\lambda} \frac{Q^2}{Q_0^2} \left(\frac{x_B}{x_0}\right)^\lambda$$


$$x = x_B \left(1 + \frac{4m_t^2}{Q^2}\right)$$

→ LHeC opens up a new field of top quark PDFs


Searches for RPV SUSY and stops

$$W_{\mathbb{R}p} = \underbrace{\lambda_{ijk} \hat{L}_i \hat{L}_j \hat{E}_k^C}_{\text{L-number violating terms}} + \underbrace{\lambda'_{ijk} \hat{L}_i \hat{Q}_j \hat{D}_k^C}_{\text{L-number violating terms}} + \underbrace{\epsilon_i \hat{L}_i \hat{H}_u}_{\text{bilinear terms}} + \underbrace{\lambda''_{ijk} \hat{U}_i^C \hat{D}_j^C \hat{D}_k^C}_{\text{B-number violating terms}}$$

$\Delta L = 1$, 9 λ couplings, 27 λ' couplings



similar to leptoquark searches with generation mixing



→ very promising with high luminosity

→ RPV can be probed at unprecedented levels


Upper Limit on $\text{Br}(t \rightarrow uH)$ in MVA analysis

Sun, Wang,
arXiv:1602.04670 [hep-ph]


parametrisation

HL-LHC

HL-LHC



LHC, 3000 fb^{-1} @14TeV
 < 0.0015
 (1 ab^{-1})
 2σ




LHC, 3000 fb^{-1} @14TeV
 < 0.0002 (2 ab^{-1})
 2σ
 → improves HL-LHC sensitivity

FCNC Branching Ratios at Colliders

ATLAS+CMS Preliminary
LHCtopWG


September 2018



FCNC Branching Ratios at Colliders

ATLAS+CMS Preliminary
LHCtopWG

September 2018




$E_e = 60 \text{ GeV}$
 1000 fb^{-1}

FCNC Branching Ratios at Colliders

ATLAS+CMS Preliminary
LHCtopWG

September 2018




$E_e = 60 \text{ GeV}$
 1000 fb^{-1}
 2000 fb^{-1}

FCNC Branching Ratios at Colliders

ATLAS+CMS Preliminary
LHCtopWG

September 2018



● improve limits on BR($t \rightarrow \gamma u$), BR($t \rightarrow Zu$), BR($t \rightarrow Hu$) considerably

→ test SUSY, little Higgs, technicolor...


$E_e = 60 \text{ GeV}$
 1000 fb^{-1}
 2000 fb^{-1}



Energy Recovering Linac


Energy Recovering Linac (ERL):

$E_e = 60 \text{ GeV}$



8.9 km

- **power limit: 100 MW**
- **luminosity: $10^{34} \text{ cm}^{-2} \text{ s}^{-1}$**
- **factor of 15/120 (LHeC/FCC-eh)**
- **extension of Q^2 , $1/x$ reach**



M. Klein, F. Zimmermann

Initial, tentative, rough scaling estimate of basic cost (tunnel, linac (XFEL), magnets)


Powerful ERL for Experiments (PERLE)

in Orsay

- 2 Linacs (Four 5-Cell 801.58 MHz SC cavities)
- 3 turns (160 MeV/turn)
- Max. beam energy 500 MeV at 20mA → 10 MW

- BINP
- CERN
- Daresbury/Liverpool
- Jlab
- Orsay

- CDR 1705.08783 [J. Phys G]
- TDR in 2019



First 802 MHz cavity successfully built (Jlab)



cf Walid Kaabi at Amsterdam FCC

- ERL demonstrator
- O(10 MeV) physics




LHeC Detector Layout

[arXiv:1802.04317]

- Cross section with MadGraph5
 - tree-level Feynman diagrams using p_T of scattered quark as scale for ep processes
 - Fragmentation & hadronisation uses ep-customised Pythia.

- DELPHES Fast Detector Simulation
 - ‘Standard’ GPD LHC-detectors
 - Optimising vertex resolution a la ATLAS IBL of $\sim 5 \mu\text{m}$
 - ATLAS b-tagging efficiencies
 - Using state-of-the art hadronic and el.mag. Resolutions
 - Considering displaced vertices and impact parameter distributions



Length x Diameter: LHeC (13.3 x 9 m²) HE-LHC (15.6 x 10.4) FCCeh (19 x 12)
 ATLAS (45 x 25) CMS (21 x 15): [LHeC < CMS, FCC-eh ~ CMS size]


Object	Acceptance
Electrons	$ \eta < 4.7$
Muons	$ \eta < 4.7$
Jets	$ \eta < 5$
b-tagging	$ \eta < 3.5$

Slide: M. Schott

CC Single Top Quark Cross Section

Dutta, Goyal, Kumar, Mellado,
arXiv:1307.1688 [hep-ph]

100 fb⁻¹:
2 · 10⁵ events
3 · 10⁵ events




→ LHeC offers excellent prospects for top quark physics

CC Single Top Quark Cross Section

Dutta, Goyal, Kumar, Mellado,
arXiv:1307.1688 [hep-ph]



100 fb⁻¹:
2–6 · 10⁵ events
3–10 · 10⁵ events



→ LHeC offers excellent prospects for top quark physics

CC Single Top Quark Cross Section

Dutta, Goyal, Kumar, Mellado,
arXiv:1307.1688 [hep-ph]



Backgrounds: Hadronic Channel

No.	Background Process	$p_{T,j,b} \geq 20 \text{ GeV}$ $ \eta_j \leq 5, \eta_b \leq 2.5$ $\Delta R_{j,b/j} \geq 0.4$ $\cancel{E}_T \geq 25$	$\Delta\Phi_{\cancel{E},j} \geq 0.4$ $\Delta\Phi_{\cancel{E},b} \geq 0.4$	$ m_{j_1j_2} - m_W \leq 22 \text{ GeV}$	$\sigma_{\text{eff.}}$
1	$e^- p \rightarrow \nu_e W^- \bar{b}$ without anti-top line	7.5×10^{-3}	6.8×10^{-3}	4.5×10^{-3}	2.7×10^{-3}
2	$e^- p \rightarrow \nu_e jjj$	4.2×10^0	3.6×10^0	2.4×10^0	7.2×10^{-2}
3	$e^- p \rightarrow \nu_e cjj$ & $e^- p \rightarrow \nu_e \bar{c}jj$	1.5×10^0	1.2×10^0	8.6×10^{-1}	8.6×10^{-2}
4	$e^- p \rightarrow \nu_e c\bar{c}j$	5.8×10^{-2}	5.0×10^{-2}	3.2×10^{-2}	6.7×10^{-3}
5	$e^- p \rightarrow \nu_e b\bar{b}j$	2.5×10^{-2}	2.2×10^{-2}	5.6×10^{-3}	1.3×10^{-3}
6	$e^- p \rightarrow \bar{c}\nu_e$ ($\bar{c} \rightarrow W^- \bar{s}$)	2.5×10^{-2}	2.2×10^{-2}	1.5×10^{-2}	1.5×10^{-4}


Event Selection	$p_{T,j,b} \geq 20 \text{ GeV}$ $ \eta_j \leq 5, \eta_b \leq 2.5$ $\Delta R_{j,b/j} \geq 0.4$ $\cancel{E}_T \geq 25$	$\Delta\Phi_{\cancel{E},j} \geq 0.4$ $\Delta\Phi_{\cancel{E},b} \geq 0.4$	$ m_{j_1j_2} - m_W \leq 22 \text{ GeV}$	Fiducial Efficiency	$S/\sqrt{S+B}$
SM	3.2×10^4	2.3×10^4	2.2×10^4	66.7 %	–
$SM + \sum_i \text{Bkg}_i$	6.5×10^4	5.0×10^4	4.0×10^4	61.5 %	
$ V_{tb} \Delta f_1^L = .5$	7.3×10^4	5.0×10^4	5.0×10^4	68.0 %	1.92
$f_1^R = .5$	4.6×10^4	3.2×10^4	3.2×10^4	69.7 %	1.43
$f_2^L = .5$	4.9×10^4	3.6×10^4	3.6×10^4	73.2 %	1.55
$f_2^L = -.5$	3.4×10^4	2.3×10^4	2.3×10^4	69.6 %	1.40
$f_2^R = .5$	5.7×10^4	4.1×10^4	4.1×10^4	72.3 %	1.69

Backgrounds: Leptonic Channel

No.	Background Process	$p_{T_{j,b,l}} \geq 20 \text{ GeV}, \Delta R_{j,b/j} \geq 0.4, \cancel{E}_T \geq 25$ $ \eta_j \geq 5, \eta_{b,l} \geq 2.5$	$\Delta\Phi_{\cancel{E},j} \geq 0.4$ $\Delta\Phi_{\cancel{E},b} \geq 0.4$ $\Delta\Phi_{\cancel{E},l} \geq 0.4$	$\sigma_{\text{eff.}}$
1	$e^- p \rightarrow l^- \bar{\nu}_l \nu_e j$	1.5×10^{-1}	1.4×10^{-1}	1.4×10^{-3}
2	$e^- p \rightarrow l^- \bar{\nu}_l \nu_e c$ & $e^- p \rightarrow l^- \bar{\nu}_l \nu_e \bar{c}$	6.6×10^{-3}	6.1×10^{-3}	6.1×10^{-4}
3	$e^- p \rightarrow l^- \bar{\nu}_l \nu_e b$ & $e^- p \rightarrow l^- \bar{\nu}_l \nu_e \bar{b}$ Without top line	3.6×10^{-3}	3.2×10^{-3}	1.9×10^{-3}
4	$e^- p \rightarrow e^- l^- \bar{\nu}_l c$	1.5×10^{-2}	6.9×10^{-3}	6.9×10^{-4}
5	$e^- p \rightarrow e^- l^- \bar{\nu}_l j$	1.2×10^{-1}	5.5×10^{-2}	5.5×10^{-4}

Event Selection	$p_{T_{j,b}} \geq 20 \text{ GeV}$ $ \eta_j \leq 5, \eta_b \leq 2.5$ $\Delta R_{j,b/j} \geq 0.4$ $\cancel{E}_T \geq 25$	$\Delta\Phi_{\cancel{E},j} \geq 0.4$ $\Delta\Phi_{\cancel{E},b} \geq 0.4$ $\Delta\Phi_{\cancel{E},l} \geq 0.4$	Fiducial Efficiency	$S/\sqrt{S+B}$
SM	1.2×10^4	1.1×10^4	92.0 %	–
SM + $\sum_i \text{Bkg}_i$	1.3×10^4	1.2×10^4	92.0 %	–
$ V_{tb} \Delta f_1^L = .5$	4.5×10^4	2.5×10^4	92.6 %	1.55
$f_1^R = .5$	2.8×10^4	1.6×10^4	94.1 %	1.23
$f_2^L = .5$	3.1×10^4	1.7×10^4	89.5 %	1.27
$f_2^L = -.5$	1.8×10^4	1.0×10^4	90.9 %	0.95
$f_2^R = .5$	3.6×10^4	2.0×10^4	90.9 %	1.38


Direct Measurement of $|V_{tb}|$



$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$

$\Delta|V_{tb}| \cdot 1000$

$\Delta\beta$: luminosity uncertainty



Dutta, Goyal, Kumar, Mellado,
arXiv:1307.1688 [hep-ph]

100 fb⁻¹: $\Delta|V_{tb}| = 0.005$

Search for Anomalous Wtb Couplings

= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$

Search for Anomalous Wtb Couplings

= 1 in SM


$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^-$$
$$-\frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$

Search for Anomalous Wtb Couplings




= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$

Dutta, Goyal, Kumar, Mellado, arXiv:1307.1688



$L_{int} = 100 \text{ fb}^{-1}$




+ other variables sensitive on W helicity

Search for Anomalous Wtb Couplings

Dutta, Goyal, Kumar,
Mellado, arXiv:1307.1688

= 1 in SM






$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^-$$

$$-\frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$

68% C.L.

property	precision
f_V^L	0.001-0.01
f_V^R, f_T^L, f_T^R	0.01-0.1







Search for Anomalous Wtb Couplings

= 1 in SM

$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$



68% C.L.



Search for Anomalous Wtb Couplings

= 1 in SM



$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$




Search for Anomalous Wtb Couplings

= 1 in SM


$$L = -\frac{g}{\sqrt{2}} \bar{b} \gamma^\mu V_{tb} (f_V^L P_L + f_V^R P_R) t W_\mu^- - \frac{g}{\sqrt{2}} \bar{b} \frac{i\sigma^{\mu\nu} q_\nu}{M_W} (f_T^L P_L + f_T^R P_R) t W_\mu^- + h.c.$$



f_{TR}



Gluon Parton Density Function





→ measure gluon density at high x

Top Quark Structure Function

Boroun, Phys. Lett. B744, 142 (2015)

variable flavour
number scheme
for top quark

$$\tilde{\sigma}^{t\bar{t}}(\tau_t) \rightarrow F_2^t(\tau_t) [1 - R^t(\tau_t)]$$



→ approximately: $1/\tau_t$

→ independent of x and τ_t

→ longitudinal top structure function component could be good to probe top quark density in proton at $Q^2 \simeq 4m_t^2$

NC Top Quark Production

Bouzas, Larios,
Physical Review D 88, 094007 (2013)




top pair production

single top production

DIS

photoproduction

photoproduction



e-beam 60 GeV, 100 fb⁻¹:

0.023 pb

$N_{t\bar{t}}=2,300$


0.70 pb

$N_{t\bar{t}}=70,000$

0.031 pb


$N_t=3,100$

Search for Anomalous $t\bar{t}\gamma$ Couplings




$$\mathcal{L}_{t\bar{t}\gamma} = e\bar{t} \left(Q_t \gamma^\mu A_\mu + \frac{1}{4m_t} \sigma^{\mu\nu} F_{\mu\nu} (\kappa + i\tilde{\kappa}\gamma_5) \right) t$$

electric dipole moment: $\tilde{\kappa}$



LHeC:
10% and 18% accuracy

 27% accuracy
(4.59fb⁻¹, 7 TeV)


magnetic dipole moment: κ

Bouzas, Larios,
Physical Review D 88, 094007 (2013)

Top Quark Dimension 6 Operators

$$\Lambda^2 \mathcal{L}_{4f} = C_1(\bar{\nu}_L \gamma^\mu t_L \bar{b}_L \gamma_\mu e_L + h.c.) + [C_2 \bar{\nu}_L e_R \bar{b}_R t_L + C_3 \bar{b}_L e_R \bar{\nu}_L t_R + C_4 \bar{\nu}_L e_R \bar{b}_L t_R + h.c.]$$

$\Lambda=1\text{TeV}$



property	precision
C_1	0.50-0.85
C_2^r	2.2-5.0
C_3^r	1.4-2.9
C_4^r	2.2-4.9


Sarmiento-Alvarado,
Bouzas, Larios,
arXiv:1412.6679

$$\mathcal{P}_e = 0$$

$$\mathcal{P}_e = 0.4$$



$$\mathcal{P}_e = 0.7$$

cross section




FCNC Top Couplings at Colliders


Top Quark Working Group
 Collaboration,
 arXiv:1311.2028 [hep-ph]



Reconstructed Higgs mass




Reconstructed top quark mass




Direct Measurement of $|V_{tb}|$

- ¹ including top-quark mass uncertainty
- ² σ_{theo} : NLO PDF4LHC11
- ³ NPPS205 (2010) 10, CPC191 (2015) 74 including beam energy uncertainty



$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & \mathbf{V_{tb}} \end{pmatrix}$$



Upper Limit on $\text{Br}(t \rightarrow uH)$ in MVA analysis

Sun, Wang,
arXiv:1602.04670 [hep-ph]

parametrisation

