

Aspects of MC event generation for e^+p collisions

Stefan Höche



Fermi National Accelerator Laboratory

Joint EF07/EF06 meeting on Jets at EIC

Online, 11/09/2019

Precision QCD calculations

- ▶ Inclusive DIS at NLO QCD
[Bardeen,Buras,Duke,Muta] PRD18(1978)3998
[Altarelli,Ellis,Martinelli] NPB143(1978)521
[Humpert,van Neerven] NPB184(1981)225
- ▶ ... at N²LO QCD [Zijlstra,vanNeerven]
NPB383(1992)525, PLB297(1992)377
[Moch,Vermaseren] hep-ph/9912355
- ▶ ... at N³LO QCD [Moch,Vermaseren,Vogt]
hep-ph/0504242, arXiv:0812.4168
- ▶ Di-jet production at NLO QCD
[Mirkes,Zeppenfeld] hep-ph/9511448
[Graudenz] hep-ph/9710244
[Nagy,Trocsanyi] hep-ph/0104315
- ▶ ... at N²LO QCD
[Abelof,Boughezal,Liu,Petriello] arXiv:1607.04921
[Currie,Gehrmann,Niehues] arXiv:1606.03991
[Currie,Gehrmann,Huss,Niehues] arXiv:1703.05977
- ▶ DIS at N³LO QCD, fully exclusive
[Currie,Gehrmann,Glover,Huss,Niehues,Vogt]
arXiv:1803.09973




Fixed-order calculations vs event generators

Anatomy of MC simulations

- ▶ **Hard interaction**
LO, NLO QCD/EW¹, NNLO QCD²
Generic matrix-element generators
- ▶ **Radiative corrections**
Parton Showers, YFS resummation
- ▶ **Hadronization & Decays**
Cluster / String model
Phase space or EFTs + YFS

Comparison to fixed order (FO)

- ▶ **Hard interaction**
Lower precision than FO
- ▶ **Radiative corrections**
Resummed & matched to FO
- ▶ **Hadronization & Decays**
Not accessible at FO



¹via interfaces to 1-loop generators

²for selected processes

Peculiarities of DIS


- ▶ Leading order $e^\pm p$ - scattering in collinear factorization

- ▶ No jets, sole kinematical variables are


$$Q^2 = q^2 = (k' - k)^2 \text{ and } x = \frac{Q^2}{2q \cdot p}$$

- ▶ Hadronic cm energy

$$W = Q\sqrt{(1-x)/x}$$




- ▶ QCD dynamics at higher orders



- ▶ Multiple scales, e.g. $E_{T,B}^2$
 - ▶ $e^\pm q \rightarrow e^\pm q$ if $E_{T,B}^2 \lesssim Q^2$
 - ▶ $\gamma^* g \rightarrow jets$ if $Q^2 \lesssim E_{T,B}^2$

- ▶ What makes DIS different from $e^+e^- \rightarrow jj$ and $pp \rightarrow e^+e^-$ is that the virtuality of the exchanged photon tends to be close to zero
- ▶ Also the case in low-mass Drell-Yan $pp \rightarrow e^+e^-$, but recent experimental studies usually focus on $m_{l\bar{l}} \approx m_Z$

Merging fixed-order calculations and parton-showers



Merging fixed-order calculations and parton-showers


- ▶ QCD dynamics of the multi-jet final state must be reflected accurately when identifying parton-shower branching history

[Carli,Gehrmann,SH] arXiv:0912.3715

- ▶ $e^\pm q \rightarrow e^\pm q$ if $E_{T,B}^2 \lesssim Q^2$
- ▶ $\gamma^* g \rightarrow \text{jets}$ if $Q^2 \lesssim E_{T,B}^2$
- ▶ $qg \rightarrow \text{jets}$ if $Q^2 \ll E_{T,B}^2$

- ▶ Similar to taking direct and fragmentation component into account in hard photon production at hadron colliders



[Schumann,Siegert,SH] arXiv:0912.3501




Comparison to HERA data

[Carli,Gehrmann,SH] arXiv:0912.3715

Variation of maximum matrix-element multiplicity, N_{\max}



Matching fixed-order NLO calculations to parton-showers



Matching fixed-order NLO calculations to parton-showers

Two possible ways to match NLO calculations and parton showers

MC@NLO

[Frixione,Webber] hep-ph/0204244

- ▶ Use parton-shower splitting kernel as infrared subtraction term
- ▶ Multiply LO event weight by Born-local K-factor including integrated subtraction term and virtual corrections
- ▶ Add hard remainder function consisting of subtracted real-emission correction

POWHEG


[Nason] hep-ph/0409146

- ▶ Use matrix-element corrections to replace parton-shower splitting kernel by full real-emission matrix element in first shower branching
- ▶ Multiply LO event weight by Born-local NLO K-factor (integrated over real corrections that can be mapped to Born according to parton-shower kinematics)


Both cases: Beware of sub-leading color terms and spin correlations!

Matching vs Merging

[Krauss, Schönherr, Siegert, SH] arXiv:1008.5399




[Carli, Gehrmann, SH] arXiv:0912.3715



Matching at NNLO accuracy

[Kuttimalai,Li,SH] arXiv:1809.04192




- ▶ New Sherpa module for computation of inclusive DIS at NNLO QCD
- ▶ Projection-to-Born method for fully differential fixed order predictions
[Zijlstra,vanNeerven] NPB383(1992)525, PLB297(1992)377 [Moch,Vermaseren,Vogt] hep-ph/0504242
[Bern,Dixon,Kosower] hep-ph/9708239, [Berger et al.] arXiv:0803.4180
- ▶ UN²LOPS matching to parton shower for particle-level simulations
[Lönnblad,Prestel] arXiv:1211.7278, [Li,Prestel,SH] arXiv:1405.3607
- ▶ Scale choice appropriate for simultaneous description of inclusive DIS and inclusive jet / di-jet / tri-jet production $\rightarrow \mu_{R/F}^2 = (Q^2 + (H_T/2)^2)/2$



- ▶ Good agreement with H1 measurements in both high- Q^2 and low- Q^2 region [Andreev et al.] arXiv:1406.4709, arXiv:1611.03421




NNLO particle-level simulation vs. H1 high- Q^2 data

[Kuttimalai,Li,SH] arXiv:1809.04192




NNLO particle-level simulation vs. H1 low- Q^2 data

[Kuttimalai,Li,SH] arXiv:1809.04192




Hadronization corrections at high Q^2

[Kuttimalai,Li,SH] arXiv:1809.04192



Hadronization corrections at low Q^2

[Kuttimalai,Li,SH] arXiv:1809.04192



Availability of DIS simulations

▶ Herwig

- ▶ Matching fully automated [Gieseke,Plätzer] arXiv:1109.6256
- ▶ External 1-loop providers & builtin loop library
- ▶ Merging in modified unitarized approach [Plätzer] arXiv:1211.5467, [Bellm,Gieseke,Plätzer] arXiv:1705.06700
- ▶ QED & mixed higher-order corrections work in progress

▶ Pythia



- ▶ New parton shower Dire [Prestel,SH] arXiv:1506.05057
- ▶ Unitarized merging under development [Prestel,Lönnblad] arXiv:1211.4827
- ▶ Matching via interface to POWHEG / MC@NLO

▶ Sherpa

- ▶ Matching fully automated [Krauss,Schönherr,Siegert,SH] arXiv:1008.5399, arXiv:1111.1220
- ▶ External 1-loop providers & builtin loop library
- ▶ Merging in non-unitarized approach [Krauss,Schönherr,Siegert,SH] arXiv:1207.5030
- ▶ NNLO matching [Kuttimalai,Li,SH] arXiv:1809.04192



Performance examples – Herwig

► NLO Merged calculation vs data from hep-ex/9907027



Performance examples – Pythia

- ▶ Parton-shower calculation vs data from hep-ex/0512014




Performance examples – Sherpa

[Carli,Gehrmann,SH] arXiv:0912.3715

three-jet center-of-mass frame:



$1 + 2 \rightarrow 3 + 4 + 5$

$E_3 > E_4 > E_5$



$\cos \theta_3 \quad Q^2 > 150 \text{ GeV}^2$



$\Psi_3 \quad Q^2 > 150 \text{ GeV}^2$



Spin correlations in decays and showers

- ▶ Spin-correlation algorithms known since '80s [Collins] NPB304(1988)794, [Knowles] CPC58(1990)271, [Richardson] hep-ph/0110108
- ▶ Decay correlations in \sim all generators [Gigg,Richardson] hep-ph/0703199 [Artoisenet, Frederix,Mattelaer,Rietkerk] arXiv:1212.3460 [Kuttimalai,Schumann,Siegert,SH] arXiv:1412.6478
- ▶ Spin-dependent parton showers Herwig [Richardson,Webster] arXiv:1807.01955 Vincia [Fischer,Lifson,Skands] arXiv:1708.01736

[Richardson,Webster] arXiv:1807.01955



Summary

- ▶ DIS simulations available in all three event generation frameworks
- ▶ NLO matching & merging standard, NNLO matching available
- ▶ Peculiarities of DIS require careful selection of clustering history
- ▶ Very good description of wide range of experimental data
- ▶ Hadronization corrections fairly model independent
- ▶ Spin-correlated parton showers now becoming available