

Calibration and Standardization of Large Surveys and Missions in Astronomy and Astrophysics



Contribution ID : 15

Type : **not specified**

DA-Type White Dwarfs: Soft X-ray Standards for the Calibration of X-ray Instruments

Monday, April 16, 2012 5:10 PM (0:50)

Abstract content

Thermal soft X-ray emission is detected from many hot hydrogen-rich white dwarfs (spectral type DA) with an effective temperature in excess of 20 000 K. Most of the objects with effective temperatures < 40 000 K have virtually pure hydrogen atmospheres while the majority of the hotter ones emit X-ray fluxes lower than predicted by hydrogen model atmospheres and therefore must contain heavier elements as absorbers.

Although such objects have a relatively soft X-ray spectrum, they are invaluable for the calibration of X-ray instruments. The hydrogen-rich DA-type white dwarfs HZ 43 A and Sirius B with effective temperatures of 51 100 K and 24 900 K, respectively, were used to establish soft X-ray standards: A cross-calibration between the Chandra LETG+HRC-S, the EUVE spectrometer, and the ROSAT PSPC was successfully performed.

The DA-type white dwarf HZ43 A provides, thus, an ideal calibration target for space-borne X-ray missions that cover the soft X-ray range.

Summary

Primary author(s) : Dr. RAUCH, Thomas (Institute for Astronomy and Astrophysics, Kepler Center for Astro and Particle Physics, Eberhard Karls University, Tübingen, Germany)

Presenter(s) : Dr. RAUCH, Thomas (Institute for Astronomy and Astrophysics, Kepler Center for Astro and Particle Physics, Eberhard Karls University, Tübingen, Germany)

Session Classification : Posters