

Snowmass 2021 White Paper: Charged lepton flavor violation in the tau sector

Swagato Banerjee^{1†}, Vincenzo Cirigliano^{2,3‡}, Mogens Dam⁴, Abhay Deshpande^{5,6,7}, Luca Fiorini⁸, Kaori Fuyuto², Ciprian Gal^{5,9}, Tomáš Husek¹⁰, Emanuele Mereghetti², Kevin Monsálvez-Pozo⁸, Haiping Peng¹¹, Francesco Polci¹², Jorge Portolés⁸, Armine Rostomyan¹³, Michel Hernández Villanueva¹³, Bin Yan², Jinlong Zhang¹⁴, Xiaorong Zhou¹¹

¹ *University of Louisville, Louisville KY 40292*

² *Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545*

³ *Institute for Nuclear Theory, University of Washington, Seattle WA 98195-1550*

⁴ *Niels Bohr Institute, Copenhagen University, Copenhagen, Denmark*

⁵ *Center for Frontiers in Nuclear Science, Stony Brook University, NY 11764*

⁶ *Stony Brook University, Stony Brook, NY 11794-3800*

⁷ *Brookhaven National Laboratory, Upton, NY 11973-5000*

⁸ *Instituto de Física Corpuscular (IFIC), Universidad de Valencia – CSIC, Valencia, Spain*

⁹ *Mississippi State University, MS 39762*

¹⁰ *Department of Astronomy and Theoretical Physics, Lund University, Lund, Sweden*

¹¹ *University of Science and Technology of China, Hefei 230026, China*

¹² *LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*

¹³ *Deutsches Elektronen-Synchrotron, Hamburg 22607, Germany*


¹⁴ *Shandong University, Qingdao, Shandong, China*



Snowmass Rare Processes and Precision Measurements Frontier Spring Meeting

University of Cincinnati, Cincinnati, Ohio, USA (May 16 - May 19, 2022)

Search for lepton number/flavor violation in τ decays



$$\mathcal{B}(\tau^\pm \rightarrow \mu^\pm \gamma) [\text{Lee-Shrock, Phys. Rev. D 16, 1444 (1977)}]$$
$$= \frac{3\alpha}{128\pi} \left(\frac{\Delta m_{23}^2}{M_W^2} \right)^2 \sin^2 2\theta_{\text{mix}} \mathcal{B}(\tau \rightarrow \mu \bar{\nu}_\mu \nu_\tau)$$

With $\Delta \sim 10^{-3} \text{ eV}^2$, $M_W \sim \mathcal{O}(10^{11}) \text{ eV}$
 $\approx \mathcal{O}(10^{-54})$ ($\theta_{\text{mix}} : \text{max}$)

many orders below experimental sensitivity!



LNv/LFV is NOT forbidden by any continuous symmetry
 \Rightarrow most New Physics (NP) models naturally include such processes

**Any observation of LNv/LFV
 \Rightarrow unambiguous signature of NP**

- Some models predict LFV up to existing experimental bounds
- eg. SUSY models: non-diagonal slepton mass matrix \Rightarrow LFV
- Normal (Inverted) hierarchy for slepton $\Rightarrow \tau \rightarrow \mu \gamma$ ($\tau \rightarrow e \gamma$)

New Physics expectations

- Mass dependent couplings enhance tau LFV w.r.t. lighter leptons



A. Crivellin et. al.
Phys. Rev. D 99, 035004 (2019)

Tau sector complementary to muons in certain parameter spaces of New Physics

Current limit on $\mathcal{B}(\mu \rightarrow e\gamma)$ does not forbid $\mathcal{B}(\tau \rightarrow \ell\gamma)$ above 10^{-9}


B. Grinstein, et. al. Nucl.Phys.B763, 35 (2007)

- Neutrinoless 2 and 3 body τ decays have different sensitivity


	$\mathcal{B}(\tau \rightarrow \ell\gamma)$	$\mathcal{B}(\tau \rightarrow \ell\ell\ell)$
mSUGRA+seesaw (EPJC14(2000)319, PRD66(2002)115013)	10^{-8}	10^{-9}
SUSY SO(10) (NPB649(2003)189, PRD68(2003)033012)	10^{-8}	10^{-10}
SUSY Higgs (PLB549(2002)159, PLB566(2003)217)	10^{-10}	10^{-8}
Non-Universal Z' (PLB547(2002)252)	10^{-9}	10^{-8}
SM+Heavy Majorana ν_R (PRD66(2002)034008)	10^{-9}	10^{-10}

New Physics illustrations


Tree level :



Compositeness




Leptoquarks




Heavy gauge bosons


Loop induced :



Supersymmetry




Heavy neutrinos




Extended Higgs models

Current and future experiments

Belle II at SuperKEKB




ATLAS, CMS, LHCb at LHC




STCF proposal at China/Novosibirsk




EIC at Brookhaven



FCC-ee proposal – CERN



Tentative timeline



About fifty τ decay modes and decays with τ in the final state

- **Lepton flavor violation (charge conjugate modes implied)**
 - $\tau \rightarrow e/\mu \gamma$ (Belle II, STCF, FCC-ee)
 - $\tau \rightarrow e/\mu$ (scalar/pseudoscalar/vector mesons) (Belle II)
 - $\tau \rightarrow e e e$ (Belle II)
 - $\tau \rightarrow \mu \mu \mu$ (Belle II, ATLAS, CMS, LHCb, STCF, FCC-ee)
 - $\tau \rightarrow e \mu \mu, \mu e e$ (Belle II)
 - $\tau \rightarrow e/\mu h h$ (non-resonant final states with $h=\pi/K$) (Belle II, STCF)
 - $H \rightarrow e \tau, \mu \tau$ (ATLAS, CMS)
 - $Z(Z') \rightarrow e \tau, \mu \tau$ (ATLAS, CMS)
 - $e \rightarrow \tau$ transitions (EIC)
- **Lepton number violation**
 - $\tau^- \rightarrow e^+ h^- h^-$ (non-resonant final states with $h=\pi/K$) (Belle II)
 - $\tau^- \rightarrow \mu^+ h^- h^-$ (non-resonant final states with $h=\pi/K$) (Belle II)
- **Baryon number violation**
 - $\tau^- \rightarrow \Lambda \pi^-, \bar{\Lambda} \pi^-$ (Belle II)
 - $\tau^- \rightarrow \bar{p} \mu^+ \mu^-, p \mu^- \mu^-$ (Belle II, LHCb)

Sensitivity estimates

$$B_{UL}^{90} = N_{UL}^{90} / (N_{\tau} \times \varepsilon)$$

● ε : high statistics *signal MC* simulated for different Data-taking periods

$\varepsilon = \text{Trigger} \cdot \text{Reco} \cdot \text{Topology} \cdot \text{PID} \cdot \text{Cuts} \cdot \text{Signal-Box}$

90% 70% 70% 50% 50% 50%


Cumulative:

90% 63% 44% 22% 11% ~5%

	\sqrt{s}	Luminosity (L)	$N_{\tau} = 2L\sigma$
Belle II	10.58 GeV	50 ab ⁻¹	9.2 x10 ¹⁰
HL-LHC	14 TeV	3 ab ⁻¹	$\mathcal{O}(10^{15})$
STCF	2-7 GeV	1 ab ⁻¹	7.0 x10 ⁹
FCC-ee	91.2 GeV	150 ab ⁻¹	3.4 x 10 ¹¹

(Efficiency much lower)

Current status of LFV limits



$\tau \rightarrow \mu\mu\mu$ at electron-positron annihilations


- Known initial conditions (beam energy constraint)
- Clean environment (less backgrounds)

Two independent variables:

$$M_\tau = \sqrt{E_{\mu\mu\mu}^2 - P_{\mu\mu\mu}^2}$$

$$\Delta E = E_{\mu\mu\mu}^{CMS} - E_{\text{beam}}^{CMS}$$

- ➔ ΔE close to 0 for signal
- ➔ Mass of tau daughters close to τ mass




Higher signal efficiency is foreseen at Belle II than at Belle or BaBar

- higher trigger efficiencies
- improved vertexing detectors
- upgraded tracking /calorimetry
- momentum dependent particle identification optimizations

Expected Belle II sensitivity: $\mathcal{B}(\tau \rightarrow \mu\mu\mu) < 3.6 \times 10^{-10}$ with 50ab^{-1}

Projected limits at Belle II


	Background limited search	Background free search
N_{UL}^{90}	$\sqrt{\mathcal{L}}$	2.44 [Feldman – Cousins for $N_{obs} = 0$]
B_{UL}^{90}	$\propto 1/\sqrt{\mathcal{L}}$	$\propto 1/\mathcal{L}$



Belle II to probe LFV in several channels $\simeq \mathcal{O}(10^{-10})$ to $\mathcal{O}(10^{-9})$ with 50 ab^{-1}

Beam polarization upgrade at SuperKEKB/Belle II

- Further improvements are expected with polarized beams
- With beam polarization, helicity distributions can suppress backgrounds
- Optimization study shows at least 10% improvement in $\tau \rightarrow \ell \gamma$ sensitivity



<https://arxiv.org/pdf/0810.1312.pdf>



Intriguing aspect of having the polarization is the possibility to determine the helicity structure of the LFV coupling in $\tau \rightarrow \mu \mu \mu$ from Dalitz plots.

Super Tau-Charm Facility



X. Zhou

“Physics Potential of a Super tau-Charm Facility” (RF/SNOWMASS21-RF7_RF1_STCF-013.pdf)

- Peaking luminosity $> 0.5 \times 10^{35} \text{ cm}^{-2}\text{s}^{-1}$ at 4 GeV
- Energy range $E_{\text{cm}} = 2\text{-}7 \text{ GeV}$
- **Potential** to increase luminosity and realize beam polarization
- A nature extension and a viable option for China accelerator project in the post **BEPCII/BESIII** era



At 4.26 GeV, number of tau pairs per year: $N_{\tau\tau} \sim 1.0 \text{ ab}^{-1} \times 3.5 \text{ nb} = 3.5 \times 10^9$




➤ STCF with 1 ab^{-1} :

$$\mathcal{B}_{UL}^{90}(\tau \rightarrow \gamma \mu) < \frac{N_{UL}^{90}}{2\epsilon N_{\tau\tau}} \sim 1.2 \times 10^{-8}$$

➤ STCF with 1 ab^{-1} :

$$\mathcal{B}_{UL}^{90}(\tau \rightarrow \mu \mu \mu) < \frac{N_{UL}^{90}}{2\epsilon N_{\tau\tau}} \sim 1.4 \times 10^{-9}$$

$$\mathcal{B}(\tau \rightarrow \mu\gamma)$$



♦ **Main background:** Radiative events (IRS+FSR), $e^+e^- \rightarrow \tau^+\tau^-\gamma$

□ $\tau \rightarrow \mu\gamma$ decay faked by combination of γ from IRS/FSR and μ from $\tau \rightarrow \mu\nu\bar{\nu}$

Smear with assumed FCC-ee detector resolutions (ILC-like detector):

- Muon momentum [GeV]
 $\sigma(p_T)/p_T = 2 \times 10^{-5} \times p_T \oplus 1 \times 10^{-3}$
- Photon ECAL energy [GeV]
 $\sigma(E)/E = 0.165/\sqrt{E} \oplus 0.010/E \oplus 0.011$
- Photon ECAL spatial [mm]
 $\sigma(x) = \sigma(y) = (6/E \oplus 2) \text{ mm}$

$$\sigma(m_{\gamma\mu}) = 26 \text{ MeV}; \quad \sigma(E_{\gamma\mu}) = 850 \text{ MeV}$$

□ From study (assuming 25% signal & background efficiency), projected BR sensitivity


$$2 \times 10^{-9}$$

$$\mathcal{B}(\tau \rightarrow \mu\mu\mu)$$

□ Expect this search to have *very low* background, even with FCC-ee like statistics


□ Should be able to have sensitivity down to BRs of $\lesssim 10^{-10}$

$\tau \rightarrow \mu\mu\mu$ at LHC



ATL-PHYS-PUB-2018-032

Scenario	$\mathcal{A} \times \epsilon$ [%]	$N_{\text{bkg}}^{\text{exp}}$	90% CL UL on $\text{BR}(\tau \rightarrow 3\mu)$ [10^{-9}]
High background	0.88	507.05	6.40
Medium background	0.88	152.12	2.31
Low background	0.88	50.71	1.03



CMS-TDR-016




	Category 1	Category 2
Number of background events	2.4×10^6	2.6×10^6
Number of signal events	4580	3640
Trimuon mass resolution	18 MeV	31 MeV
$B(\tau \rightarrow 3\mu)$ limit per event category	4.3×10^{-9}	7.0×10^{-9}
$B(\tau \rightarrow 3\mu)$ 90% C.L. limit	3.7×10^{-9}	

LHCb-PUB-2018-009

The cross-section is five orders of magnitude larger than at Belle II. This compensates for the higher background levels and lower integrated luminosity. As pointed out in [76], during the HL-LHC era, the LHCb Upgrade II detector will allow to collect 300 fb^{-1} . With this large data sample, LHCb will be able to probe the branching ratio down to $O(10^{-9})$, and either independently confirm any Belle II discovery or significantly improve the limit.

$e \rightarrow \tau$ transitions at EIC

\sqrt{s} between 20 GeV (5 GeV electron on 41 GeV protons)
and 140 GeV (18 GeV electron on 275 GeV protons)





Deshpande, Huang, Kumar Zhang, Zhao

Potential to improve current sensitivity by two orders of magnitude

Tau to electron transitions

Model-independent probes of new physics at scale (Λ) encoded as Wilson coefficients (C_n) via EFT approach




e-Print: [2203.14919](https://arxiv.org/abs/2203.14919) [hep-ph]

Tau to muon transitions

Model-independent probes of new physics at scale (Λ) encoded as Wilson coefficients (C_n) via EFT approach

WC	Operator	WC	Operator
$C_{LQ}^{(1)}$	$(\bar{L}_p \gamma_\mu L_r) (\bar{Q}_s \gamma^\mu Q_t)$	$C_{e\varphi}$	$(\varphi^\dagger \varphi) (\bar{L}_p e_r \varphi)$
$C_{LQ}^{(3)}$	$(\bar{L}_p \gamma_\mu \sigma^I L_r) (\bar{Q}_s \gamma^\mu \sigma^I Q_t)$	$C_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (e_p \gamma^\mu e_r)$
C_{eu}	$(\bar{e}_p \gamma_\mu e_r) (\bar{u}_s \gamma^\mu u_t)$	$C_{\varphi L}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{L}_p \gamma^\mu L_r)$
C_{ed}	$(\bar{e}_p \gamma_\mu e_r) (\bar{d}_s \gamma^\mu d_t)$	$C_{\varphi L}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_{I\mu} \varphi) (\bar{L}_p \sigma_I \gamma^\mu L_r)$
C_{Lu}	$(\bar{L}_p \gamma_\mu L_r) (\bar{u}_s \gamma^\mu u_t)$	C_{eW}	$(\bar{L}_p \sigma^{\mu\nu} e_r) \sigma_I \varphi W_{\mu\nu}^I$
C_{Ld}	$(\bar{L}_p \gamma_\mu L_r) (\bar{d}_s \gamma^\mu d_t)$	C_{eB}	$(\bar{L}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$
C_{Qe}	$(\bar{Q}_p \gamma_\mu Q_r) (\bar{e}_s \gamma^\mu e_t)$		
C_{LedQ}	$(\bar{L}_p^j e_r) (\bar{d}_s Q_t^j)$		
$C_{LeQu}^{(1)}$	$(\bar{L}_p^j e_r) \varepsilon_{jk} (\bar{Q}_s^k u_t)$		
$C_{LeQu}^{(3)}$	$(\bar{L}_p^j \sigma_{\mu\nu} e_r) \varepsilon_{jk} (\bar{Q}_s^k \sigma^{\mu\nu} u_t)$		



e-Print: [2203.14919](https://arxiv.org/abs/2203.14919) [hep-ph]

Summary and outlook

$\tau^- \rightarrow$	Observed Limits			Expected Limits		
	Experiment	Luminosity	UL (obs)	Experiment	Luminosity	UL (exp)
$\mu^- \gamma$	Belle [93]	988 fb ⁻¹	4.2×10 ⁻⁸	Belle II [54]	50 ab ⁻¹	6.9×10 ⁻⁹
	BaBar [83]	516 fb ⁻¹	4.4×10 ⁻⁸			
				STCF [74]	1 ab ⁻¹	1.8×10 ⁻⁸
$\mu^- \mu^+ \mu^-$				FCC-ee [87, 91]	150 ab ⁻¹	$\mathcal{O}(10^{-9})$
	Belle [102]	782 fb ⁻¹	2.1×10 ⁻⁸	Belle II [54]	50 ab ⁻¹	3.6×10 ⁻¹⁰
	BaBar [103]	468 fb ⁻¹	3.3×10 ⁻⁸			
	LHCb [61]	3 fb ⁻¹	4.6×10 ⁻⁸	LHCb [76]	300 fb ⁻¹	$\mathcal{O}(10^{-9})$
	CMS [67]	33 fb ⁻¹	8.0×10 ⁻⁸	CMS [77]	3 ab ⁻¹	3.7×10 ⁻⁹
	ATLAS [68]	20 fb ⁻¹	3.8×10 ⁻⁷	ATLAS [78]	3 ab ⁻¹	1.0×10 ⁻⁹
				STCF [74]	1 ab ⁻¹	1.4×10 ⁻⁹
				FCC-ee [87, 91]	150 ab ⁻¹	$\mathcal{O}(10^{-10})$

- **Observation of LFV in the charged lepton sector would completely change our understanding of physics and herald a new period of discoveries in particle physics. Synergies between different experiments compliment discovery potential/confirmation.**
- **Now is a very interesting era in the searches for LFV in decays of the tau lepton, as the current limits will improve by an order of magnitude down to a few parts in 10⁻¹⁰ to 10⁻⁹ at the Belle II experiment. Polarized beams can further improve the sensitivity.**
- **Similar sensitivities will be probed at ATLAS, CMS & LHCb with high luminosity upgrade.**
- **Proposed experiments at STCF, EIC & FCC-ee will continue searches for LFV in the tau sector, also with the possibility of beam polarization.**