



Latest results from JSNS² and status of JSNS²-II

**Jungsic Park (Kyungpook National University)
On behalf of the JSNS² / JSNS²-II collaboration**

NuFACT 2022, July 31 - August 6, 2022


Talks in the parallel sessions

- (WG 6) Tuesday, 04:00 PM: “Gain calibration using dark hits in off-time region of regular data at JSNS² experiment”
- (WG 1) Thursday, 05:28 PM: “Status of the KDAR neutrino search with JSNS² experiment”

JSNS² / JSNS²-II Collaboration

(J-PARC Sterile Neutrino Search at J-PARC Spallation Neutron Source)

Collaboration meeting @ J-PARC (2020/Feb)



- JSNS² collaboration (61 collaborators)
- 10 Korean institutions (24 members)
 - 6 Japanese institutions (29 members)
 - 4 US institutions (7 members)
 - 1 UK institution (1 member)



Chonnam National
Jeonbuk National
Dongshin
GIST
Kyungpook
Kyung Hee
Seoyeong
Soongsil
Sungkyunkwan
Seoul National of sci
and tech



JAEA
KEK
Kitasato
Kyoto
Osaka
Tohoku



BNL
Florida
Michigan
Utah



Sussex



Indication of a sterile neutrino ($\Delta m^2 \sim 1 eV^2$)

(Direct test of the LSND)

Experiments (Neutrino source, signal, significance, energy, baseline)



- **LSND** (μ Decay-At-Rest, $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$, 3.8σ , 40 MeV, 30 m)
- **MiniBooNE** (π Decay-In-Flight, $\nu_\mu \rightarrow \nu_e$, $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$, 4.8σ (combined), 800 MeV, 600 m)
- **BEST** (e capture, $\nu_e \rightarrow \nu_x$, $\sim 4 \sigma$, < 3 MeV, 10 m)
- **Reactors** (Beta decay, $\bar{\nu}_e \rightarrow \bar{\nu}_x$, significance varies, 1-8 MeV, 10 - 100 m)

JSNS² uses the **same** neutrino source (μ), target (H), and detection principle (IBD) **as the LSND**

- Even if the excess is not due to the oscillation, JSNS² can catch this directly.

Two advantages: **short-pulsed beam** and **use of the gadolinium(Gd)-loaded liquid scintillator**

J-PARC Facility



Low duty factor beam
(short-pulses + low repetition rate)
Gives an excellent signal to noise ratio

1 MW (design)


0.6 MW (2021/Jan - Apr/5)

0.7 MW (2021/Apr/5 - June/22)

0.7 MW (2022/Jan/28 - Apr/6)

0.8 MW (2022/Apr/7 -)

JSNS² detector and data taking



17 tons target, Gd-LS + 10% DIN
120, 10-inch PMTs

Commissioning (2020)

- Calibration
- Beam data with 25 us window
- Eur. Phys. J. C (2022) 82:331

First long physics run (2021)

- Smooth data taking (0.5 years)
- Beam power: 0.6 - 0.7 MW
- Analyses: in progress

Second long physics run (2022)

- Jan/28 - May/31
- Beam power: 0.7 - 0.8 MW

Operation

1st long physics run


- 0.6 MW (2021/Jan - Apr/5)
- 0.7 MW (2021/Apr/5 - June/22)

2nd long physics run

- 0.7 MW (2022/Jan/28 - Apr/6)
- 0.8 MW (2022/Apr/7 - May/31)

There is an accelerator maintenance period every year

2.77×10^{22} POT has been delivered by May/20
(23 % of the approved POT of JSNS²)




Sterile neutrino

(Production and detection)

- A double coincidence between
- The positron annihilation and
 - Gammas from neutron captured on gadolinium (Gd)

	Timing	Energy
prompt	$1.5 \leq T_p \leq 10 \mu s$	$20 \leq E \leq 60 \text{ MeV}$
delayed	$\Delta T_{p-d} < 100 \mu s$	$7 \leq E \leq 12 \text{ MeV}$






Expected visible energy and sensitivity

(From JSNS² TDR, arXiv:1705.08629)




- $\bar{\nu}_e$ follows decay-at-rest $\bar{\nu}_\mu$ energy distribution
- Prompt candidate: $\sim 3.9 \times 10^{-4}$ per spill
- Delayed candidate: $\sim 4.4 \times 10^{-3}$ per spill
- Spectral fit is sensitive to the difference of energy spectrum

Signal	$\sin^2 2\theta = 3.0 \times 10^{-3}$ $\Delta m^2 = 2.5 eV^2$ (Best fit values of MLF)	87
	$\sin^2 2\theta = 3.0 \times 10^{-3}$ $\Delta m^2 = 1.2 eV^2$ (Best fit values of LSND)	62
background	$\bar{\nu}_e$ from μ^-	43
	$^{12}\text{C}(\nu_e, e^-)^{12}\text{N}_{g.s.}$	3
	beam-associated fast n	≤ 2
	Cosmic-induced fast n	negligible
	Total accidental events	20



Commissioning run (Eur. Phys. J. C (2022) 82:331)

- June/5-15, 2020
- Integrated POT: 8.9×10^{20}
- Beam trigger with $25 \mu s$ width



Observed correlated event candidates

- 59 ± 8 events / 8 M spills
- **Cosmic-induced fast neutrons** are the dominant background
 - **Correlated background:** 55.9 ± 4.3
 - **Pulse shape discrimination (PSD) would reject them.**
 - Two independent groups are working on it.

Analysis using 2021 data set

Pulse Shape Discrimination (PSD)

(2-dimensional likelihood method)


- It has been developed **based on the real data of JSNS²**.
- **10% DIN** (2000 L) has been added for improving the PSD power.
- Based on the charge ratio of each FADC bin (208 bins in total) divided by the peak of each waveform.
- **Using control samples of Michel electrons (ME) and Fast neutrons (FN)**

• PSD evaluation score: $\sum_{PMT} Q_{PMT}^{total} [\ln(L_{ME}) - \ln(L_{FN})]$

where $L = \prod_{bin} [P_{bin}(Q_{ratio})]$, $Q_{ratio} = Q_{bin}/Q_{peak}$

- Note that there are two types of ME for considering afterpulse.



Example of waveforms



Pulse Shape Discrimination (PSD)

(2-dimensional likelihood method)



ME efficiency: $91.4 \pm 0.5 \%$
FN rejection: $95.4 \pm 0.3 \%$





Pulse Shape Discrimination (PSD)

(Convolutional Neural Network, CNN)

- Treated time-series data from a PMT with image data
- Data: Training (37.5%), validation (12,5%) and evaluation (50 %)
- Two independent efforts show **consistent FN-rejection result**




Jungsic Park (KNU)




Roadmap

(sterile neutrino search)

Estimation of the
background



Blind analysis



Sensitivity

JSNS² is here

- Reduction
- Event selection
- **Compare single rate with TDR**
- Coincidence
 - Energy and timing
 - Delta vertex
 - Fiducial
- More sophisticated method

- Energy side-bands (and timing side-band)
- **Consistency check**
- self-trigger / beam trigger
- PSD
- More sophisticated method
- Toward a signal region


- Uncertainties
 - Flux
 - Energy scale
 - Others

Cosmic muon identification

- Michel electron induced by cosmic muon and muon itself are one of the backgrounds.
- Tag muons passing through the detector & stopping at the detector using veto information.

JSNS² equips 24 Veto PMTs

- 12 PMTs on top / 12 PMTs on bottom
- 6 PMTs among 12 face radial direction
- Other half face vertical direction
- Muon candidate rate [Hz]: 1487.8 ± 0.6
- Michel candidate rate [Hz]: 110.1 ± 0.2 (> 10 MeV)




Single rate of the sterile prompt

(Background estimation effort)

- External particle rejection with Veto PMTs
- **ME rejection: 10 μs vetoing after a stopping-muon**
- Fiducial cut
- Decomposition have been performed
 - To estimate each component
 - correlated / accidental
 - **Spectrum is obtained from each control sample.**
- **Reference (JSNS² TDR): 3.8×10^{-4} per spill**

Prompt candidate after ME rejection




component	rate [Hz]
cosmic fast neutron (green)	3.9±0.7
michel electron (cyan)	13.3±0.3
cosmic gamma (violet)	9.3±0.6
stopping muon (gray)	0.01±0.02
total (black)	26.8±0.1
rate per spill	(2.144±0.010)x10 ⁻⁴

Toward the sterile neutrino search

(For the blind analysis)

- **Side-bands are defined by energies**
- The rates in the side-band regions can be predicted by other data
 - self-trigger: cosmic-induced neutron
 - beam-trigger: accidental
- After application of PSD, all of side-band regions will be accidental dominated.
- JSNS² has been studying each side-band
- All side-bands should be understood thoroughly before opening the signal region





- WG 1, Thursday, 05:28 PM:
- “Status of the KDAR neutrino search with JSNS² experiment”

Kaon Decay-At-Rest (KDAR) Neutrino (Toward first KDAR precise measurement)


Kaon Decay-At-Rest neutrino measurement (KDAR neutrino: 236 MeV mono-energetic)

- Neutrino interaction models are a crucial part of neutrino physics, but poorly known at low energies.
- **The JSNS² detector has the unique ability to measure the mono-energetic KDAR neutrino.**
- Note that horn focused beam can not make a decay-at-rest neutrino.



Probing the nucleus with KDAR neutrinos

- KDAR neutrinos: a known-energy, weak-interaction-only probe of the nucleus, right at the transition between neutrino-nucleus and neutrino-nucleon scattering



$$K^+ \rightarrow \mu^+ \nu_\mu \text{ [BR = 63.6 \%]}$$
$$E_\nu = 236 \text{ MeV if } K^+ \text{ is at rest}$$

Why it is important

- The transition region between neutrino-nucleus and neutrino-nucleon scattering is very hard to model
- Models and generators strongly disagree.



Relevant for:

- Long baseline exps and modeling of neutrino-nucleus interactions in the 100s-of-MeV region.
- Understanding MiniBooNE low-energy excess.
- Supernova neutrinos.
- Solar dark matter annihilation signatures.
- Future oscillation searches with KDAR (muon disappearance and electron appearance)

Only previous KDAR measurement (MiniBooNE)
[3.9 σ first observation of KDAR neutrinos]

Jungsic Park (KNU)

NuFACT 2022




KDAR signal measurement in JSNS²


A double coincidence between

- The initial neutrino interaction products and the subsequent muon decay.



KDAR



Prompt



Delayed




KDAR event selection

Used data: 2.256×10^8 spill


- 11.9% of the approved POT

- Identify muon and remove proceeding $10 \mu s$ events
- Beam-timing cut (150 ns each)
- Prompt candidate: 20 - 140 MeV (μ^-)
- Delayed candidate: 20 - 60 MeV (e^-)
- Delta T: $< 10 \mu s$
- Delta vertex: < 300 mm
- Applied the Fiducial cut
- Background template: timing side-band
- Magnitude: area normalization b/w 140 - 250 MeV

Template of background energy



Visible energy spectrum w/ backgrounds




First clear KDAR signal

(Toward first precise KDAR measurement)

- KDAR peak is clearly seen
- High purity (95%) KDAR signal
 - Background: 5.2 %
- Note that the systematics on the energy scale are not included yet.


BKG ID	Correlated/ Accidental	BKG (# of events)	
1	Correlated	36.6 ± 34.8	$5.0 \pm 5.1\%$
2	Accidental	1.5 ± 0.1	$0.2 \pm 0.01\%$
KDAR Signal : 730 events		38.1 ± 38.4	$5.2 \pm 5.3\%$




**Toward the second phase
(JSNS²-II)**

JSNS²-II

(arXiv:2012.10807)



- Second phase of the JSNS² experiment with two detectors
 - near (17 tons, 120 10-inch PMTs, 24 m),
 - far (32 tons, 220 10-inch PMTs, 48 m) detector
- Improve the sensitivity especially in the low Δm^2 region
- J-PARC/KEK grants the stage 2 (2/2) approval
- The stainless steel tank has been constructed
- A lot of progress about PMTs



demonstration of the PMT attachment



catcher + veto + spill tank



PMTs




- 190 PMTs arrived at J-PARC on June 27 in 2022 (from Paris)
- JSNS²-II appreciates to Double-Chooz experiment
- Dismantling the DC PMT jigs structure has been finished.
- Dead PMTs checks have been done
- Will be installed in the detector soon.



Sensitivity for the JSNS²-II


(Based on the simulation)

- Each background simulation was done based on the JSNS² data
- Covering LSND by 3 sigma



JSNS²-II: Schedule


(Even under the COVID-19, the overall schedule was well followed from 2021's plan)



Summary

- **JSNS² is working toward first precise KDAR measurement**
 - **Clearly see the high purity KDAR signal**
- There have been 1st (2021) and 2nd (2022, ongoing) long physics runs in JSNS².
- Analyses are ongoing with the data.
 - Has been developing two separate PSD tools
 - Sterile neutrino search is on-going according to the roadmap
- Based on the JSNS² data, JSNS²-II has been granted.
- Even under the COVID-19, the overall schedule was well followed from 2021's plan.
- JSNS²-II expects to start data taking at around the end of 2023.

Thank you for your attention



acknowledgements:

- MEXT, JSPS (Japan)
- Korea Ministry of Science, NRF (Korea)
- DOE, Heising-Simons Foundation (US)
- Royal Society (UK)




KDAR summary slide

- JSNS2 clearly observes a high purity (95%) KDAR signal!
- This signal will provide the first precision measurement of KDAR neutrinos!
- Working towards finalizing visible energy

$(T_\mu + \sum T_p)$ spectrum measurement and detailed model comparison.





- Will elucidate the difficult to model neutrino-nucleon to neutrino-nucleus transition region, highly relevant across many aspects of neutrino physics



Backup





Dedicated sterile trigger

(Conditional and sequential of beam and self trigger)



Electronics upgrade (Under developing)

- The goal is to provide excellent efficiency and resolution over a wide energy range
- FPGA based trigger
- Hosted by an ATCA shelf
- A combined test has been done



JSNS² vs. LSND

	LSND	JSNS ²	Advantage of JSNS ²
Detector Mass	167 Tons	17 Tons	
Baseline	30 m	24 m	
Beam Kinetic Energy	0.8 GeV	3.0 GeV	Allows for KDAR measurement / 10 times higher pion production
Beam Power	0.8 MW	1.0 MW (designed)	More intense beam
Beam Pulse	600 us, 120 Hz	100 ns (x2), 25 Hz	300 times less steady-state background for BID
Capture Nucleus	H (2.2 MeV)	Gd (~8 MeV)	Shorter capture time, higher signal to ratio