

Charged-meson-induced new physics in beam-focused neutrino experiments

Thursday, August 4, 2022 11:20 AM (30 minutes)

We point out that the production of new bosons by charged meson decays can greatly enhance the sensitivity of beam-focused accelerator-based experiments to new physics signals. This enhancement arises since the charged mesons are focused and their three-body decays do not suffer from helicity suppression in the same way as their usual two-body decays. As a realistic application, we attempt to explain the MiniBooNE low energy excess utilizing this overlooked mechanism, uniquely realizing dark-sector interpretations as plausible solutions to the excess. For illustration purposes, we consider two well-motivated classes of dark-sector models, models of vector-portal dark matter and models of long-lived (pseudo)scalar. We argue that the model parameter values to accommodate the excess are consistent with existing limits and that they can be tested at current and future accelerator-based neutrino experiments.

Attendance type

In-person presentation

Primary author: KIM, Doojin (Texas A&M University)

Co-authors: DUTTA, Bhaskar (Texas A&M University); VAN DE WATER, Richard (LANL); THOMPSON, Adrian (Texas A&M University); THORNTON, Remington

Presenter: KIM, Doojin (Texas A&M University)

Session Classification: WG5: Beyond PMNS

Track Classification: WG5: Beyond PMNS