

# Long-Baseline Neutrino Oscillation Physics Sensitivities of the Hyper-Kamiokande Experiment

*Friday, August 5, 2022 2:56 PM (18 minutes)*

The Hyper-Kamiokande experiment has a rich long-baseline neutrino program, as well as a variety of other physics goals. The long-baseline program will utilize a world-class neutrino beam produced at the high-intensity J-PARC accelerator constrained by a suite of near detectors. The Hyper-Kamiokande detector is a ~185 kton fiducial volume water Cherenkov detector, located 295 km from the beam neutrino source. Hyper-Kamiokande long-baseline measurements will be sensitive to the leptonic CP violating phase,  $\delta_{CP}$ , as well as the atmospheric oscillation parameters,  $\sin^2\theta_{23}$  and  $\Delta m^2_{32}$ . Combining the long-baseline and atmospheric neutrino measurements at the Hyper-Kamiokande detector will also allow for enhanced sensitivities, and in particular will help to resolve the neutrino mass ordering.

## Attendance type

In-person presentation

**Primary author:** FRIEND, Megan (KEK)

**Presenter:** FRIEND, Megan (KEK)

**Session Classification:** WG1: Neutrino Oscillations

**Track Classification:** WG1: Neutrino Oscillation Physics