

## Status of the KDAR neutrino search with JSNS2 experiment

*Thursday, 4 August 2022 17:16 (12 minutes)*

Kaon Decay-At-Rest (KDAR) provides a neutrino signal with well-known neutrino energy, which is an important probe for measuring the neutrino cross-section in an energy range that is otherwise difficult to access experimentally. The J-PARC Sterile Neutrino Search at the J-PARC Spallation Neutron Source (JSNS2) experiment is in a unique place for measuring monoenergetic neutrinos at 236 MeV from charged Kaon decay-at-rest (KDAR). JSNS2 is located at the J-PARC's Material and Life Science Facility (MLF) where the world's most intense source of KDAR was created by a 3 GeV proton beam incident on a liquid mercury target. In this presentation, We will present the first result of the search for the KDAR neutrinos conducted with the JSNS2 experiment with the data during the JSNS2's first long-term physics run in 2021, consisting of more than 115 days of data and  $1.45 \times 10^{22}$  POT.

### Attendance type

Virtual presentation

**Primary author:** JEON, Hyoungku (SungKyunKwan University)

**Presenter:** JEON, Hyoungku (SungKyunKwan University)

**Session Classification:** WG1: Neutrino Oscillations

**Track Classification:** WG1: Neutrino Oscillation Physics