

# Tension between the T2K and NOvA appearance data and hints to new physics

*Friday, 5 August 2022 17:16 (22 minutes)*

The tension between the T2K and NOvA long-baseline experiments arises mostly due to the mismatch in the  $\nu_\mu \rightarrow \nu_e$  and  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$  appearance data. Assuming vacuum oscillation as the reference point, with maximal  $\theta_{23}$  and  $\delta_{CP} = 0$ , we compute the  $\nu_e/\bar{\nu}_e$  appearance events for each of the experiments. T2K observes a large excess in the  $\nu_e$  appearance event sample compared to the expected  $\nu_e$  events at the reference point, whereas NOvA observes a moderate excess. The large excess in T2K dictates that  $\delta_{CP}$  be anchored at  $-90^\circ$  and that  $\theta_{23} > \pi/4$  with a preference for normal hierarchy (NH). The moderate excess at NOvA leads to two degenerate solutions: (a) NH,  $0 < \delta_{CP} < 180^\circ$ , and  $\theta_{23} > \pi/4$ ; (b) inverted hierarchy (IH) with  $-180^\circ < \delta_{CP} < 0$ , and  $\theta_{23} > \pi/4$ . This is the main cause of tension between the two experiments. We show that beyond the standard model (BSM) physics scenarios such as non-unitary neutrino mixing, Lorentz invariance violation, and non-standard neutrino interactions, may resolve the tension.

## Attendance type

Virtual presentation

**Primary authors:** RAHAMAN, Ushak (University of Johannesburg); RAZZAQUE, Soebur (University of Johannesburg); UMASANKAR, Sankagiri (Indian Institute of Technology Bombay)

**Presenter:** RAHAMAN, Ushak (University of Johannesburg)

**Session Classification:** WG1: Neutrino Oscillations

**Track Classification:** WG1: Neutrino Oscillation Physics