

# Electromagnetic Energy Reconstruction in ProtoDUNE

Aleena Rafique

On behalf of the DUNE Collaboration  
ICHEP 2022




## 1. DUNE/ProtoDUNE Experiment

### DUNE:


- 1300 km baseline
- 70 kton Liquid Argon Time Projection Chamber (LArTPC) Far Detector (FD) at SURF, South Dakota, 1.5 km underground [1]
- Multiple technologies for the Near Detector (ND) at Fermilab
- Will measure neutrino oscillation probability to determine mass ordering and CP violation phase; potential for BSM physics and supernova neutrinos

### ProtoDUNE-single phase:


- ~7x6x7 m<sup>3</sup> in charged test beam at CERN
- A crucial part of the DUNE effort towards the construction of the first DUNE far detector module
- ProtoDUNE-SP I operated from September 2018 to July 2020



## 2. Event Displays




## 3. Michel Electron: Selection



## 4. Reconstruction

All hits confined inside a cone at the end point of the parent muon are taken to be Michel hits. Reconstructed Michel energy spectrum is derived using stopping muon calibration [2].

- Accuracy of the reconstructed Michel energy spectrum is > 98%




## 5. Resolution

► Michel resolution is parametrized by:


$$\frac{\sigma(E)}{E} = p_0 \oplus \frac{p_1}{\sqrt{E}} \oplus \frac{p_2}{E}$$

- The energy resolution improves after adding the missing energy contributions
- The constant term captures the resolution losses due to the missing energy




## 6. Beam Electron: Selection

► Beam electron selection flow chart



## 7. Reconstruction

- Selected runs with different beam momentum values
- Evaluated the reconstructed electron shower energies
- Linear correlation between electron beam momentum and shower energy shows great response of TPC to electron beams




## 8. Resolution

► Beam electron resolution is parametrized by:


$$\sigma_E/E = \sqrt{a^2 + (b/\sqrt{E})^2 + (c/E)^2}$$

- True energy resolution for 1 GeV (1.2%, gray plot) comparable to values found in other simulations for liquid argon (e.g. 1.68% in [3])



## 9. Neutral pion: Selection

- EM CNN score > 0.5: Remove track-like daughter particles
- # hits (total) > 80: Remove low completeness showers
- 3 cm < dist. < 90 cm: Remove charged pions and muons
- IP < 20 cm: Remove low completeness showers




Purity: ~ 80%, Efficiency: ~ 37%

## 10. Reconstruction

► π<sup>0</sup> mass distribution after correcting for the negative bias (-15%) in the reconstructed shower energy

► The energy spectrum of reconstructed neutral pion after selections and kinematic fitting is also plotted



## 11. Resolution

- The resolution for the neutral pion energy as a function of true energy after correction and kinematic fitting is presented
- The analysis will be repeated after including more data

