


Renormalization of quasi transverse-momentum-dependent parton distribution function (quasi TMD PDF) on the lattice

Speaker: Kuan Zhang

Xiangdong Ji, Yi-Bo Yang, Fei Yao, Jian-Hui Zhang


Lattice 2023 2023.07.31-2023.08.04

Motivation




- TMDPDF describes the hadron structure in the transverse direction, and is much less known.
- How to determine it from first principle?

Large Momentum Effect Theory (LaMET)



Lorentz boost and perturbative matching

Light-cone PDF : Cannot be calculated on the lattice




Quasi-PDF : Directly calculable on the lattice

Outline

- 1、 Renormalization of quasi PDF operator in RI/MOM scheme
- 2、 Renormalization of quasi TMD-PDF operator
 - 2.1 RI/MOM scheme
 - 2.2 Wilson loop + Short distance hadron matrix element
- 3、 Numerical details
- 4、 Summary



1、Quasi-PDF operator

$$O_{\gamma_t}(z) = \bar{q}(0)\gamma_t U_z(0,z)q(z)$$



$$\langle \pi | O_{\gamma_t}(z) | \pi \rangle^{\text{RI/MOM}} \equiv \frac{\langle \pi | O_{\gamma_t}(z) | \pi \rangle}{\langle q | O_{\gamma_t}(z) | q \rangle}$$


RI/MOM
renormalization



Coulomb gauge quasi PDF (preliminary)

Xiang Gao, Wei-Yang Liu, Yong Zhao, arXiv: 2306.14960


$$O_c(z) = \bar{\psi}_c(0)\Gamma\psi_c(z\hat{n}_z)$$



2、Quasi TMD-PDF on lattice

- Matrix element
$$\tilde{h}_{\chi,\gamma_t}(b, z, L, P_z; 1/a) = \langle \chi(P_z) | O_{\gamma_t}(b, z, L) | \chi(P_z) \rangle,$$
$$O_\Gamma(b, z, L) \equiv \bar{\psi}(\vec{0}_\perp, 0) \Gamma \mathcal{W}(b, z, L) \psi(\vec{b}_\perp, z)$$
$$h_{\chi,\gamma_t}(b, z, P_z; 1/a) = \lim_{L \rightarrow \infty} \frac{\tilde{h}_{\chi,\gamma_t}(b, z, L, P_z; 1/a)}{\sqrt{Z_E(b, 2L + z; 1/a)}}$$

- Staple-shaped link




X. Ji, Y. Liu, Y. S. Liu, J.-H. Zhang, and Y. Zhao, Rev. Mod.Phys. 93, 035005 (2021).

X. Ji, Y. Liu, and Y.-S. Liu, Nucl. Phys. B955, 115054 (2020).


M. A. Ebert, I.W. Stewart, and Y. Zhao, J. High Energy Phys. 09 (2019) 037.

Lattice setup


- $\langle \pi(0) | o_{\gamma_t}(b, z) | \pi(0) \rangle$ pion ME (matrix element) in rest frame wall source
- $\langle q(p) | o_{\gamma_t}(b, z) | q(p) \rangle$ $(p_x, p_y, p_z, p_t) = \frac{2\pi}{La} (0, 5, 0, 5)$ $|p| \sim 3 GeV$
- 1-step HYP smearing on staple link
- Clover & overlap valence quark




2.1 Staple link ME renormalized in RI/MOM scheme



$$\begin{aligned}
 h_{\chi,\Gamma}^{\text{MOM}}(b,z,P_z;p) &= \sum_{\Gamma'} [Z^{\text{MOM}}(b,z,p;1/a)]_{\Gamma\Gamma'}^{-1} h_{\chi,\Gamma'}(b,z,P_z;1/a) \\
 &= \lim_{L \rightarrow \infty} \sum_{\Gamma'} \tilde{Z}^{\text{MOM}}(b,z,L,p;1/a)]_{\Gamma\Gamma'}^{-1} \tilde{h}_{\chi,\Gamma'}(b,z,L,P_z;1/a),
 \end{aligned}$$



2.2 Staple link ME renormalized in SDR scheme



$$h_{\chi, \gamma_t}^{\text{SDR}}(b, z, P_z; \frac{1}{b_0}) = \frac{h_{\chi, \gamma_t}(b, z, P_z; 1/a)}{h_{\pi, \gamma_t}(b_0, z_0 = 0, 0, 1/a)}$$



$$\begin{aligned} h_{\chi, \gamma_t}^{\overline{\text{MS}}}(b_0, z_0, 0; \mu) &= 1 + \frac{\alpha_s C_F}{2\pi} \left\{ \frac{1}{2} + 3\gamma_E - 3 \ln 2 \right. \\ &\quad \left. + \frac{3}{2} \ln [\mu^2(b_0^2 + z_0^2)] - 2 \frac{z_0}{b_0} \arctan \frac{z_0}{b_0} \right\} \\ &\quad + \mathcal{O}(\alpha_s^2) \end{aligned}$$

$$h_{\chi, \gamma_t}^{\overline{\text{MS}}}(b, z, P_z; \mu) = h_{\gamma_t}^{\overline{\text{MS}}}(b_0, 0, 0; \mu) h_{\chi, \gamma_t}^{\text{SDR}}(b, z, P_z; \frac{1}{b_0})$$

The UV divergence is of a multiplicative structure
 M. A. Ebert, I.W. Stewart, and Y. Zhao, J. High Energy Phys. 03 (2020) 099
 J. R. Green, K. Jansen, and F. Steffens, Phys. Rev. D 101, 074509 (2020).



The mixing for nonchiral lattice fermion actions appears to be negligibly small
 Y. Ji, J.-H. Zhang, S. Zhao, and R. Zhu, Phys. Rev. D 104, 094510 (2021).
 Q.-A. Zhang, et.al, LPC, Phys. Rev. Lett. 125, 192001(2020)
 Constantia Alexandrou, et.al, arXiv:2305.11824

2.2 Staple link ME renormalized in SDR scheme





$$m_{\text{eff}}(b, r; a) = \ln \frac{Z_E(b, r; a)}{Z_E(b, r+1; a)}$$

3.1 Extrapolation of Wilson loop






3.1 Extrapolation of Wilson loop





$$h_{\chi,\gamma_t}(b, z, P_z; 1/a) = \lim_{L \rightarrow \infty} \frac{\tilde{h}_{\chi,\gamma_t}(b, z, L, P_z; 1/a)}{\sqrt{Z_E(b, 2L + z; 1/a)}}$$

3.2 L dependence of the matrix elements



To balance the statistical uncertainty and the systematic uncertainty, we take $L = 6a$ for MILC12, $L = 8a$ for MILC09, $L = 11a$ for MILC06, $L = 13a$ for MILC04, and $L = 14a$ for MILC03 in most cases. For the cases with larger b on the MILC03 ensemble, we set $L = b$ (more specifically, we set $L = 16a$ for $b = 16a$, and $L = 18a$ for $b = 18a$).

3.3 Action dependence



3.4 Operator mixing







$$\mathcal{M}_{\mathcal{P}\Gamma}(b, z; 1/a) = \frac{\text{Abs} \left[\text{Tr} \left[\mathcal{P} \left\langle q(p) \middle| \bar{q} \Gamma \mathcal{W}(b, z, L) q \middle| q(p) \right\rangle \right] \right]}{\text{Abs} \left[\text{Tr} \left[\Gamma \left\langle q(p) \middle| \bar{q} \Gamma \mathcal{W}(b, z, L) q \middle| q(p) \right\rangle \right] \right]}$$

$\Gamma \backslash \Gamma$	1	y^t	y^x	y^y	y^z	σ^{tx}	σ^{ty}	σ^{tz}	σ^{xy}	σ^{xz}	σ^{yz}	$y^5 t$	$y^5 x$	$y^5 y$	$y^5 z$	y^5
1	1.00	0.05	0.06	0.05	0.07	0.03	0.00	0.11	0.03	0.06	0.11	0.02	0.00	0.02	0.00	0.00
y^t	0.01	1.00	0.04	0.01	0.06	0.04	0.00	0.15	0.00	0.02	0.00	0.10	0.07	0.03	0.01	
y^x	0.06	0.05	1.00	0.05	0.05	0.00	0.00	0.02	0.00	0.15	0.02	0.10	0.00	0.10	0.00	0.00
y^y	0.01	0.01	0.04	1.00	0.05	0.00	0.00	0.04	0.02	0.15	0.07	0.10	0.00	0.03	0.01	
y^z	0.06	0.05	0.03	0.05	1.00	0.02	0.00	0.00	0.02	0.03	0.00	0.02	0.00	0.02	0.00	
σ^{tx}	0.02	0.03	0.00	0.00	0.02	1.00	0.05	0.04	0.01	0.05	0.00	0.00	0.02	0.07	0.04	0.11
σ^{ty}	0.00	0.00	0.00	0.00	0.00	0.04	1.00	0.05	0.04	0.00	0.05	0.01	0.07	0.01	0.06	0.15
σ^{tz}	0.10	0.15	0.02	0.00	0.00	0.04	0.05	1.00	0.00	0.05	0.01	0.00	0.02	0.07	0.01	0.02
σ^{xy}	0.02	0.00	0.00	0.03	0.02	0.01	0.05	0.00	1.00	0.05	0.04	0.07	0.02	0.00	0.04	0.11
σ^{xz}	0.07	0.02	0.15	0.02	0.04	0.05	0.00	0.04	0.05	1.00	0.04	0.02	0.00	0.01	0.00	0.00
σ^{yz}	0.10	0.00	0.02	0.15	0.00	0.00	0.05	0.01	0.04	0.05	1.00	0.07	0.02	0.00	0.01	0.02
$y^5 t$	0.02	0.00	0.11	0.07	0.03	0.00	0.01	0.00	0.07	0.04	0.06	1.00	0.04	0.01	0.05	0.00
$y^5 x$	0.00	0.11	0.00	0.10	0.00	0.02	0.06	0.04	0.02	0.00	0.04	0.05	1.00	0.05	0.04	0.03
$y^5 y$	0.02	0.07	0.11	0.00	0.03	0.07	0.01	0.06	0.00	0.04	0.00	0.01	0.04	1.00	0.05	0.00
$y^5 z$	0.00	0.02	0.00	0.02	0.00	0.02	0.06	0.01	0.02	0.00	0.01	0.05	0.04	0.05	1.00	0.16
y^5	0.00	0.01	0.00	0.01	0.00	0.09	0.06	0.02	0.09	0.00	0.02	0.00	0.04	0.00	0.14	1.00

$\Gamma \backslash \Gamma$	1	y^t	y^x	y^y	y^z	σ^{tx}	σ^{ty}	σ^{tz}	σ^{xy}	σ^{xz}	σ^{yz}	$y^5 t$	$y^5 x$	$y^5 y$	$y^5 z$	y^5
1	1.00	0.00	0.01	0.00	0.01	0.05	0.00	0.19	0.05	0.06	0.19	0.00	0.00	0.00	0.00	0.00
y^t	0.01	1.00	0.07	0.04	0.08	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.20	0.06	0.05	0.00
y^x	0.01	0.07	1.00	0.07	0.05	0.00	0.00	0.00	0.01	0.00	0.00	0.19	0.00	0.19	0.00	0.00
y^y	0.01	0.04	0.07	1.00	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.06	0.20	0.00	0.05
y^z	0.01	0.07	0.03	0.07	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.04	0.00	0.00
σ^{tx}	0.04	0.00	0.00	0.00	0.00	1.00	0.07	0.04	0.02	0.07	0.01	0.00	0.00	0.00	0.20	
σ^{ty}	0.00	0.00	0.00	0.00	0.00	0.06	1.00	0.08	0.06	0.00	0.08	0.00	0.01	0.00	0.01	0.15
σ^{tz}	0.20	0.01	0.00	0.00	0.00	0.04	0.07	1.00	0.01	0.07	0.02	0.00	0.01	0.01	0.00	0.04
σ^{xy}	0.04	0.00	0.00	0.00	0.00	0.02	0.07	0.01	1.00	0.07	0.04	0.00	0.00	0.00	0.00	0.20
σ^{xz}	0.06	0.00	0.01	0.00	0.00	0.08	0.00	0.06	0.08	1.00	0.06	0.01	0.00	0.01	0.00	0.00
σ^{yz}	0.20	0.00	0.00	0.01	0.00	0.01	0.07	0.02	0.04	0.07	1.00	0.01	0.01	0.00	0.00	0.04
$y^5 t$	0.00	0.01	0.20	0.06	0.05	0.00	0.00	0.00	0.01	0.00	0.01	1.00	0.07	0.04	0.08	0.00
$y^5 x$	0.00	0.19	0.00	0.19	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.07	1.00	0.07	0.05	0.00
$y^5 y$	0.00	0.06	0.20	0.01	0.05	0.01	0.00	0.01	0.00	0.00	0.00	0.04	0.07	1.00	0.08	0.00
$y^5 z$	0.00	0.04	0.00	0.04	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.07	0.03	0.07	1.00	0.01
y^5	0.00	0.00	0.00	0.00	0.00	0.17	0.06	0.04	0.17	0.00	0.04	0.00	0.00	0.01	1.00	

3.5 Matrix element at larger b

$$h_{\chi,\gamma_t}^{\overline{\text{MS}}}(b, z, P_z; \mu) = h_{\gamma_t}^{\overline{\text{MS}}}(b_0, 0, 0; \mu) h_{\chi,\gamma_t}^{\text{SDR}}\left(b, z, P_z; \frac{1}{b_0}\right)$$







3.6 TMD wave function matrix element

$$\Psi_{\pi,\gamma_5\gamma_t}(b,z,P_z;1/a) = \lim_{L \rightarrow \infty} \frac{\langle O_{\gamma_5\gamma_t}(b,z,L)|\pi(P_z)\rangle}{\langle O_{\gamma_5\gamma_t}(0,0,0)|\pi(P_z)\rangle \sqrt{Z_E(b,2L+z;1/a)}},$$

$$\Psi_{\pi,\gamma_5\gamma_t}^{\text{SDR}}(b,z,P_z;\frac{1}{b_0}) = \frac{\Psi_{\pi,\gamma_5\gamma_t}(b,z,P_z;1/a)}{\Psi_{\pi,\gamma_5\gamma_t}(b_0,z_0=0,0,1/a)}.$$



$$\begin{aligned}\bar{\Psi}_{\gamma_5\gamma_t}^{\overline{\text{MS}}}(b_0,z_0,0;\mu) &= h_{\chi,\gamma_t}^{\overline{\text{MS}}}(b_0,z_0,0;\mu) \\ &= 1 + \frac{\alpha_s C_F}{2\pi} \left\{ \frac{1}{2} + 3\gamma_E - 3\ln 2 + \frac{3}{2} \ln[\mu^2(b_0^2 + z_0^2)] - 2\frac{z_0}{b_0} \arctan \frac{z_0}{b_0} \right\} + \mathcal{O}(\alpha_s^2), \\ \bar{\Psi}_{\pi,\gamma_5\gamma_t}^{\overline{\text{MS}}}(b,z,P_z;\mu) &= \bar{\psi}_{\gamma_5\gamma_t}^{\overline{\text{MS}}}(b_0,0,0;\mu) \Psi_{\pi,\gamma_t}^{\text{SDR}}(b,z,P_z;\frac{1}{b_0}).\end{aligned}$$



4、Summary

- The square root of the **Wilson loop** combined with the **short distance hadron matrix element** provides a successful method to **remove all ultraviolet divergences** of the quasi TMD-PDF operator
- Quasi TMD-PDF renormalized in **RI/MOM** scheme suffers from **residual linear divergence**.

backup



Hybrid-renormalization

X. Ji. et.al. , NPB964(2021)115311

Self-renormalization

Y.-K. Huo. et.al. ,LPC, NPB969(2021)115443

.....