



Contribution ID: 376

Type: **Parallel Talk**

Form factors for semileptonic B-decays with HISQ light quarks and clover b-quarks in Fermilab interpretation

Monday, 31 July 2023 14:10 (20 minutes)

We compute the complete set of form factors for the $B \rightarrow \pi$, $B \rightarrow K$, and $B_s \rightarrow K$ amplitudes, which are needed to describe semileptonic B -meson decay rates for both the charged and neutral current cases. We use the highly improved staggered quark (HISQ) action for the sea and light valence quarks. The b quark is described by the Wilson-clover action in the Fermilab interpretation. Simulations are carried out on $N_f = 2 + 1 + 1$ MILC HISQ ensembles at approximate lattice spacings from 0.15 fm down to 0.057 fm. We present preliminary results for the form factors.

Topical area

Quark and Lepton Flavor Physics

Primary authors: EL-KHADRA, Aida (University of Illinois at Urbana-Champaign); VAQUERO, Alejandro (University of Utah); KRONFELD, Andreas (Fermilab); LYTTLE, Andrew (University of Illinois at Urbana-Champaign); DETAR, Carleton (University of Utah); GAMIZ, Elvira (University of Granada); JEONG, Hwancheol (Indiana University); GOTTLIEB, Steven (Indiana Univ.); JAY, William (MIT); GELZER, Zechariah (University of Illinois at Urbana-Champaign)

Presenter: JEONG, Hwancheol (Indiana University)

Session Classification: Quark and Lepton Flavor Physics