



Contribution ID: 164

Type: Poster

## Optimizing plasma-downramp profiles and beam transport for emittance preservation in multi-stage plasma accelerators

Plasma-based particle accelerators maintain accelerating fields that are several orders of magnitude higher than conventional accelerators. This allows for more compact accelerator footprints that can deliver particle beams of very high charge ( $> 100$  pC) and large current ( $> \text{kA}$ ) for various applications. For instance, plasma-wakefield accelerators are promising candidates for next-generation TeV-class electron-positron colliders for high-energy physics and secondary light sources. However, to reach the desired TeV energy regime, a staging approach of independent laser-driven plasma accelerators that each preserve low energy spread and beam emittance is required. Maintaining beam emittance over tens and hundreds of stages is a serious challenge but is crucial to achieving a high luminosity in future colliders. We present results for the optimization of plasma-stage downramp profiles and inter-stage beam transport in simulations of multi-stage plasma accelerators, carried out with codes from the Beam pLasma & Accelerator Simulation Toolkit (BLAST) and steered by *optimas*, a Python library for optimization at scale, powered by *libEnsemble*.

---

Supported by the CAMPA collaboration, a project of the U.S. Department of Energy, Office of Science, ASCR and HEP, SciDAC program, and Exascale Computing Project. See comments for full details.

### Working group

WG7 : Linear Colliders

**Primary author:** GARTEN, Marco (LBNL)

**Co-authors:** LEHE, Remi (LBNL); BENEDETTI, Carlo (LBNL); SHAPOVAL, Olga (LBNL); SANDBERG, Ryan (LBNL); HUEBL, Axel (Lawrence Berkeley National Laboratory, USA); VAY, Jean-Luc (Lawrence Berkeley National Laboratory)

**Presenter:** GARTEN, Marco (LBNL)

**Session Classification:** Poster