

NuInt 2024 - 14th International Workshop on Neutrino-Nucleus Interactions

Thursday, 11 April 2024 - Saturday, 20 April 2024

Instituto Principia



Book of Abstracts

Contents

Opening	1
Alleviating the present tension between T2K and NO ν ν A with neutrino New Physics at source	1
Neutral and Charged forbidden contributions to ν - ^{40}Ar cross sections	1
Electron/positron beam identification analysis in FERMILAB	1
Parallelizing the computation of neutrino oscillation parameters in a general background medium, by using the Jacobi Method	2
The search for Majorana neutrinos with nEXO	2
Impact of interaction uncertainties on oscillation measurements	3
Impact of interaction uncertainties on neutrino cross section measurements	3
Systematic uncertainty reduction and residual cross section uncertainties for PRISM	3
Impact of neutrino cross section uncertainties on T2K physics measurements	3
TBA	3
Meson-exchange currents in quasielastic electron and neutrino scattering in a generalized superscaling approach	4
JLab Spectral Functions of Argon in NuWro and Their Implications for MicroBooNE	4
Probing Earth's internal structure with high energy neutrino transmission	5
Random Forests for determining muon numbers in extensive air showers relative to their primary energy using KASCADE-Grande data	5
TBA	6
TBA	6
TBA	6
TBA	6
Constraining Lorentz Invariance Violation Parameters Using Short and Long Baseline Experiments	6

First Measurement of Double-Differential Charged Current $\nu\mu$ -Argon Scattering Cross Sections In Kinematic Imbalance Variables With The MicroBooNE Detector	7
Cross Section Systematics in DUNE	7
LIV Studies for Inverted Hierarchy Scenario	7
Muon-neutrino charged-current cross sections from MicroBooNE: first simultaneous measurements of final states with and without protons for Muon-neutrino scattering on argon	8
Data-driven model validation for neutrino-Argon inclusive measurements at MicroBooNE	8
GiBUU-Based Monte Carlo Simulation for Neutrino Experiments	9
The development of the R-LDH Innovative media for O2 Capturing in Liquid Argon	9
Evaluation of the Leggett-Garg inequality by means of the neutrino oscillations observed in reactor and accelerator experiments	10
Development and application of the nuclear deexcitation simulator NucDeEx for precise prediction of neutrino-nuclear interactions	10
Bounds on Lorentz Violation Parameters: present and future	11
Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method	11
Status of the Measurement of the Elastic and Quasi-elastic Scattering of Protons on Carbon Nuclei in EMPHATIC	12
Probing light vector mediators with coherent scattering at future facilities	12
Ant colony in Sensitivity Studies	13
Nuclear effects in single pion production: hadron final state interactions	13
Refining Neutrino Final State Interactions in carbon and lead: Insights from extrapolating between extremes	14
Studying Neutrino-Nucleus Interactions at SBND with Muon Neutrino CC Events	14
New developments in the Ghent Model for Single Pion Production	14
Neutrons from CCQE-like Antineutrino Interactions in MINERvA's Various Nuclear Targets	15
Influence of complex phases from off-diagonal parameters that violate Lorentz invariance in DUNE	16
Professor Based ReWeight for GENIE Generator	16
Hybrid model for single-pion production incorporated in the NuWro event generator nuclear framework	17
Impact of neutrino cross section uncertainties on NOvA physics measurements	17

Impact of neutrino cross section uncertainties on MicroBooNE and SBN measurements . . .	17
GENIE	17
NuWro	17
ACHILLES	18
INCL+ABLA	18
NEUT	18
GiBUU	18
Characterization of Neutral-current background induced by atmospheric neutrinos using neutrino event generators and the TALYS deexcitation package	18
High-energy neutrino-matter interaction cross-sections in neutrino event generators . . .	18
Flux Overview	19
Fermilab Beams	19
J-PARC neutrino beam performance, monitoring and flux prediction, constraints and un- certainties	19
Spallation Fluxes	19
NA61	19
EMPHATIC	19
ENUBET	19
ν -e Scattering	20
ICARUS	20
MicroBooNE - 1	20
MicroBooNE - 2	20
SBND	20
LIT+coupled cluster, SF from coupled cluster	20
short range correlations, generalized contact formalism	20
Two body currents in Hartree-Fock Mean Field framework	21
DWIA comparisons to recent measurements	21
Telegrapher equation for photon diffusion in LArTPCs with photon removal at the bound- aries	21
MicroBooNE - 1	21
MicroBooNE - 2	21

ICARUS status and plans for measurements related to ν -Ar interactions with no mesons in the final state and/or QE- and 2p2h-like interactions	22
SBND	22
Meson exchange currents in relativistic mean field	22
Short time approximation	22
DWIA comparison to recent experimental data	22
Extended superscaling with two-particle emission in electron and neutrino scattering	22
Improving neutrino reconstruction in LArTPCs by including charge and light signal	22
Electrons for Neutrinos	23
The Hyper-K Intermediate Water Cherenkov Detector and the Water Cherenkov Test Experiment	23
Measuring muon antineutrino charged-current interactions without mesons in the final state, in the NOvA Near Detector	24
CEvNS and BSM physics with the CONNIE experiment.	24
Eta production from nucleons and nuclei	25
Nuclear medium effects in ν - τ -A scattering at DUNE energies	25
Neutrino-induced single pion production and the reanalyzed bubble chamber data	25
Photo-, Electro- and Weak Single Pion Production Model	25
Pion Production at MINERvA	25
ν_e and $\bar{\nu}_e$ low recoil in MINERvA	25
SIS Interactions in MINERvA	26
Pion Production in NOvA	26
$e4\nu$	26
Determination of the spectral functions of argon at JLab	26
Electron Scattering in Generators	26
Empirical fits to inclusive electron-carbon scattering data obtained by deep-learning methods	26
Charged particles interaction measurements in ProtoDUNE	26
ANNIE	27
Low-energy: Theory Overview in Inelastic and CEvNS Cross-sections and Impact of BSM Effects	27
Low-energy Inelastic Cross-sections: Experiment Review and Future Prospects	27

Neutrino-nucleus coherent scattering: status and prospects	27
Lattice QCD	27
NUISANCE and global fits	27
MINERvA form-factor measurement	27
T2K Near Detector Upgrade	28
Resonance and QE contributions to inclusive nucleon structure functions	28
The Physics of SIS – TMC and HT	28
Superscaling in the inelastic region: The SuSAv2-inelastic model	28
Neutrino Cross Sections for collider neutrinos	28
MicroBooNE	28
T2K latest results and prospects on NC and CC pion production	29
DUNE 2x2	29
Impact of Cross Section Uncertainties on BSM Searches	29
Experimental Summary	29
Theory Summary	29
Closing Remarks	29
MINERvA	29
NOvA	30
T2K	30
Double Beta Decay within the Relativistic QRPA	30
Welcome	31
General introduction	31
General theory at the nucleon level	31
Introduction to nuclear effects	31
Relativistic mean field approach	31
Spectral function approach	31
Ab initio approach	31
How to make a cross-section measurement	32
Generator comparisons to data	32
Hands-on exercises with generators	32

How generators work	32
Practicum: analyzing generator output	32
Generators review	32
Practicum continued	33
Meson exchange currents in a relativistic mean field model	33
Introduction to simulation	33
Practicum: running a generator	33
Welcome	33
ICTP-SAIFR	33
Institute Principia	33
Registration	34
Information	34
Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method	34
Ant colony in Sensitivity Studies	34
CEvNS and BSM physics with the CONNIE experiment.	34
Discussion	34
Closing remarks	34

1

Opening

Corresponding Authors: orlandop@unicamp.br, helio@fnal.gov

Poster Session / 6

Alleviating the present tension between T2K and NO ν ν A with neutrino New Physics at source

Authors: Adriano Cherchiglia¹; Orlando Luis Goulart Peres²; Edson Souza³; Fernanda Rodrigues³; Pedro Simoni Pasquini⁴; Rafaela Rossi³

¹ *Universidade Estadual de Campinas*

² *UNICAMP*

³ *University of Campinas*

⁴ *Unicamp*

Corresponding Author: peres.orlando@gmail.com

Since neutrino oscillation was observed, several experiments have been built to measure its parameters. At the present there is tension between T2K and NOVA. We propose a non-standard neutrino interaction at production. In this scenario, computed by quantum field theory formalism can make a better description of combined data. A new phase from this new interaction can make a role in search for violation of charge and parity symmetry.

Poster Session / 7

Neutral and Charged forbidden contributions to ν -⁴⁰Ar cross sections

Author: Arturo Samana¹

¹ *Universidade Estadual de Santa Cruz*

Corresponding Author: arturo.samana@gmail.com

The contributions of neutral and charged forbidden and allowed multipole to ν -⁴⁰Ar cross sections are evaluated within the QRPA and PQRPA models as function of the neutrino energy in the interval of interest for supernova neutrinos. These calculations are important as input for neutrino generator and simulator as the Marley code used for DUNE experiment.

Poster Session / 8

Electron/positron beam identification analysis in FERMILAB

Authors: Carlos Javier Solano Salinas¹; Edgar Chavarria²

¹ *UNMSM, Peru*

² UNI, Peru**Corresponding Author:** csolanos@unmsm.edu.pe

In High Energy Physics it is essential the study of the particles that make up everything (at least the known baryonic universe). Carrying out a study of these particles is necessary devices (detectors). These devices interact with particles through known physical processes and then, through a data acquisition system, one can proceed for further analysis. In the MINERØA experiment, the cross-section measurements are generally done. As a previous step, they used a mini test detector called Test Beam II (TB II) to see the response of new materials (objectives and scintillators) to different energy bands. In this work, the response of the Beam II test detector to electron and positron beams are studied. The response of this detector is studied for energies in the range of 2 to 8 MeV. A data/MC comparison is also performed to see discrepancies with the model used, as well as in the analysis of the behavior of electron and positron beams under different parameters (electromagnetic cascade opening angle, electromagnetic cascade starting module and energy absorbed by the calorimeter). Our method has allowed the differentiation of electron and positron beams, unequivocally for the energies in the range 2-6 MeV and, to a lesser extent, in the range 6-8 MeV. This is a genius result given that previous experiments did not obtain results that could differentiate well electron beams relative to positron beams in these beam energy ranges

Poster Session / 9

Parallelizing the computation of neutrino oscillation parameters in a general background medium, by using the Jacobi Method

Author: Gustavo Valdiviesso¹¹ Universidade Federal de Alfenas**Corresponding Author:** gustavo.valdiviesso@unifal-mg.edu.br

This work describes an alternative tool for the study of matter effects over neutrino propagation, by numerically describing the mixing amplitudes and mass splittings as a function of a single parameter. It allows the description of these observables in the same manner as it is commonly seen in 2-neutrino phenomenology, identifying resonant mixing and level crossing, except in a N-neutrino scenario. This is accomplished by sequentially applying the Jacobi diagonalization method over a defined and smooth path, in infinitesimal steps, transporting the information about the neutrino mixing and parametrization from one background to another. This transport occurs in the parameter space, thus being reliable even over spatial background discontinuities. Due to the nature of the neutrino mass-flavor mixing, it is possible to describe any background effects, in a model-independent way, by making use of the vacuum mixing only. This is computationally interesting since the workload can be split in two stages, separating the diagonalization from the physical interpretation and fitting of any model in question. The procedure is detailed and benchmarked against known analytical solutions, together with a complexity and computational-time analysis.

Poster Session / 10

The search for Majorana neutrinos with nEXO

Author: Caio Licciardi¹¹ Laurentian University**Corresponding Author:** caio.licciardi@snolab.ca

Neutrinos are the only known fermions with zero electric charge, and thus far the unique candidates to act as their own antimatter counterpart. If indeed a neutrino is its own antiparticle, we can find processes in which lepton number conservation, a global symmetry of the Standard Model, is violated. Such a discovery would have great implications in neutrino physics and their interactions. The most sensitive test for this condition, also known as Majorana neutrinos, is through the search of a hypothetical decay known as neutrinoless double beta decay. The primary focus of the nEXO Collaboration is the search for this process using a liquid xenon time projection chamber, at the tonne-scale rooted on the success of the EXO-200 experiment. Our projections result in a half-life sensitivity beyond 10^{28} yr, sufficient to cover a milestone of this search consisting of the inverted ordering of neutrinos masses. This poster presents the search as well as the nEXO detector and its potential for discovery of new physics.

Impact of scattering uncertainties on measurements / 11

Impact of interaction uncertainties on oscillation measurements

Corresponding Author: clarence.wret@physics.ox.ac.uk

Impact of scattering uncertainties on measurements / 12

Impact of interaction uncertainties on neutrino cross section measurements

Corresponding Author: vpandey@fnal.gov

Impact of scattering uncertainties on measurements / 13

Systematic uncertainty reduction and residual cross section uncertainties for PRISM

Corresponding Author: i.caracas@uni-mainz.de

Impact of scattering uncertainties on measurements / 14

Impact of neutrino cross section uncertainties on T2K physics measurements

Corresponding Author: laura.munteanu@cern.ch

Impact of scattering uncertainties on measurements / 15

TBA

Poster Session / 18

Meson-exchange currents in quasielastic electron and neutrino scattering in a generalized superscaling approach

Authors: JOSE ENRIQUE AMARO¹; Paloma Rodríguez Casalé^{None}

¹ UNIV. GRANADA

Corresponding Author: palomacasale@ugr.es

A method that enables the consistent inclusion of meson-exchange currents (MEC) within the framework of the superscaling analysis with relativistic effective mass will be presented. We use a novel definition of the single nucleon tensor, defined as the mean value of the single-nucleon responses by averaging with an energy distribution $n()$ [1]. This single nucleon prefactor is obtained from the 1p-1h matrix element of the OB current combined with the two body current. The averaging definition is extended beyond the scaling region of the ψ variable characteristic of the Fermi gas modifying the momentum distribution through a smeared Fermi distribution [2].

In the generalized scaling analysis, the selected QE data generate a band with scatter of 20% at the maximum (highlighting the extent of scaling violation) that will be parametrized using a simple function $f(\psi^*)$.

Through the inclusion of MEC, we conducted a comparative analysis of the 1p-1h response functions within the context of the RFG, RMF and SuSAM* models. These responses can be expressed in a factorized way as the product of the averaged single nucleon multiplied by the scaling function. A reduction in the transverse response is observed, which is in accordance with previous calculations [3][4].

In this manner, the 1p-1h cross section including MEC is calculated and compared with the calculation without MEC and the experimental data [5][6]. Our examination shows that in the low momentum limit, the predictions of the relativistic model align with those of the non-relativistic model in Fermi gas. The formalism can be extended to calculate the 1p-1h neutrino nuclear responses by including the meson exchange currents using the models already proposed in electron scattering.

[1] P. R. Casale, J. E. Amaro, V. L. Martinez-Consentino and I. Ruiz Simo, Universe 9, no.4, 158 (2023) [arXiv:2303.03890 [nucl-th]].

[2] P. R. Casale, J. E. Amaro and M. B. Barbaro, Symmetry 15 (2023) no.9, 1709 [arXiv:2307.15783 [nucl-th]].

[3] J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, C. Maieron and J. M. Udias, Phys. Rev. C 81 (2010), [arXiv:0906.5598 [nucl-th]].

[4] J. E. Amaro, A. M. Lallena and G. Co, Nucl. Phys. A 578 (1994), 365-396 doi:10.1016/0375-9474(94)90752-8

[5] Benhar, O.; Day, D.; Sick, I. An archive for quasi-elastic electron-nucleus scattering data [arXiv:nucl-ex/0603032]

[6] Benhar, O.; Day, D.; Sick, I. Available online: <https://www.faculty.virginia.edu/qes-archive/> C83 (2011) 045501.

Poster Session / 19

JLab Spectral Functions of Argon in NuWro and Their Implications for MicroBooNE

Author: Rwik Dharmapal Banerjee¹

Co-authors: ARTUR ANKOWSKI²; JAN SOBCZYK²; HEMANT PRASAD ; Krzysztof Graczyk ; Beata KOWAL

¹ University of Wrocław

² UNIVERSITY OF WROCLAW

Corresponding Author: rwiku98@gmail.com

The Short-Baseline Neutrino program in Fermilab aims to resolve the nature of the low-energy excess events observed in LSND and MiniBooNE, and analyze with unprecedented precision neutrino interactions with argon, which requires reliable estimate of neutrino cross sections. We report updates of the NuWro generator that bring the state-of-the-art spectral functions to model the ground state properties of the argon nucleus, and improve the accuracy at low energies by accounting for effects of Coulomb potential and nuclear recoil. We discuss these developments in context of electron and neutrino interactions, by comparing NuWro predictions to experimental data from JLab Hall A and MicroBooNE. The MicroBooNE data are described with the Chi square per degree of freedom of 0.7, compared with 1.0 in the local Fermi gas model. Being obtained using the axial form factor parametrization from MINERvA, our results indicate a consistency between the CCQE measurements in MINERvA and MicroBooNE

Impact of scattering uncertainties on measurements / 20

Probing Earth's internal structure with high energy neutrino transmission

Authors: Reinaldo Francener¹; Victor Gonçalves²

¹ UNICAMP

² UFPEL

Corresponding Author: reinaldofrancener@gmail.com

The opacity of the Earth for the incidence of high energy neutrinos is directly related to the neutrino-nucleon cross sections and the description of the distribution of matter in the Earth's interior. In this work we investigate the sensitivity of neutrino transmission to the use of different models for Earth's density profile. In particular, we compare neutrino transmission using the PREM density model with three-, two- and one-layer models (core, mantle and crust) and constant densities. To calculate the neutrino-nucleon cross section we used the DGLAP predictions, with the CT14 parametrizations. Our results indicate that neutrino transmission is sensitive to different Earth profile density models at magnitudes that depend on the kinematic region tested and the models compared. We also show that the regeneration of the tau neutrino flux by its decay significantly impacts flux transmission in all models studied. The results achieved indicate that Earth profile density models impact the events observed at neutrino observatories such as IceCube.

Poster Session / 21

Random Forests for determining muon numbers in extensive air showers relative to their primary energy using KASCADE-Grande data

Author: omar moises asto rojas¹

Co-authors: Juan Carlos Arteaga-Velázquez²; Carlos Javier Solano Salinas³

¹ Universidad Nacional de Ingeniería

² Instituto de Física y Matemáticas, Universidad Michoacana

³ UNMSM, Peru

Corresponding Author: oasto1310@gmail.com

This study presented the development and application of two Random Forest (RF) models to KASCADE-Grande data with zenith angles $\theta < 40^\circ$, obtained from the KASCADE Cosmic Ray Data Centre. The

aim was to predict the primary energy and correct the muon number for systematic effects on an event-by-event basis. Model training involved KCDC simulations using the high-energy hadronic interaction model QGSjet-II-04. To refine the RF models, we conducted feature engineering on the magnitudes and fine-tuned the hyperparameters. The results enabled us to estimate the number of muons relative to the primary energy in extensive air shower events across the energy range from 10 PeV to 1 EeV. A comparison was made between the experimental results and the expectations from the QGSJET-II-04 model, considering pure protons and iron nuclei, respectively.

Impact of scattering uncertainties on measurements / 22

TBA

Impact of scattering uncertainties on measurements / 23

TBA

Impact of scattering uncertainties on measurements / 24

TBA

Impact of scattering uncertainties on measurements / 25

TBA

27

Constraining Lorentz Invariance Violation Parameters Using Short and Long Baseline Experiments

Authors: Andre Steklain¹; Luciana Hirsch²

¹ *Universidade Tecnológica Federal do Parana*

² *UTFPR*

Corresponding Author: steklain@utfpr.edu.br

The Lorentz Invariance is deeply connected to Special Relativity, which states that the laws of physics are the same for different observers in relative motion. It is the foundation of other successful theories, like quantum field theory, and connected to fundamental symmetries, like charge, parity, and time reversal (CPT), which is essential in the Standard Model of particle physics. Nevertheless, alternative theories proposing that Lorentz Invariance may break in some scales have been considered in the context of neutrino oscillations, as they can explain some anomalies present in experiments like LSND and MiniBoone. The dependence on baseline and energy can distinguish the influence of LIV on these anomalies in contrast with other effects, like Non-Standard Interactions (NSI). In this

work, we perform simulation studies to study the influence of Lorentz-violating parameters in neutrino experiments combining two different baselines. We use the General Long Baseline Experiment Simulator (GLoBES) with a modified probability engine to include LIV parameters.

Poster Session / 28

First Measurement of Double-Differential Charged Current $\nu\mu$ -Argon Scattering Cross Sections In Kinematic Imbalance Variables With The MicroBooNE Detector

Author: Afroditi Papadopoulou^{None}

Corresponding Author: apapadop@fnal.gov

Making high-precision measurements of neutrino oscillation parameters requires an unprecedented understanding of neutrino-nucleus scattering. In this work, we present the first charged current double-differential cross sections in kinematic imbalance variables. These variables characterize both the transverse and total kinematic imbalance in a neutrino interaction. We use events with a single muon above 100 MeV/c, a single final state proton above 300 MeV/c, and no recorded final state pions. Thus, these variables act as a direct probe of nuclear effects such as final state interactions, Fermi motion, and multi-nucleon processes. Our measurement allows us to constrain systematic uncertainties associated with neutrino oscillation results performed by near-future experiments of the Short Baseline Neutrino (SBN) program, as well as by future large-scale experiments like DUNE.

Poster Session / 29

Cross Section Systematics in DUNE

Author: Lars Bathe-Peters¹

¹ University of Oxford

Corresponding Author: lars.bathe-peters@physics.ox.ac.uk

For the operation of precision neutrino experiments, the understanding of neutrino interactions with matter are preconditioned requirements of all detections and measurements of neutrinos. The largest uncertainties in estimating neutrino-nucleus interaction cross sections arise in the incomplete understanding of nuclear effects. In the study of neutrino oscillations and nuclear scattering processes, obtaining an interaction model with associated uncertainties is of substantial interest for the neutrino physics community. This report presents studies of simulated CC 2p-2h interactions, in which a neutrino interacts with a bound pair of nucleons. This interaction mode is very poorly constrained by current data. A comparison of three leading CC 2p-2h models is presented, along with a number of uncertainty parameters that have been implemented to account for model-to-model discrepancies in the DUNE oscillation analysis.

30

LIV Studies for Inverted Hierarchy Scenario

Author: Thiago Cruz¹

Co-authors: Angelo Thomazi¹; Diego Madeira¹; Lucas De Lima¹; Samuel Marcolin¹; Johan Wodzinski¹; Andre Steklain²; Luciana Hirsch³

¹ *Universidade Tecnológica Federal do Paraná*² *Universidade Tecnológica Federal do Paraná*³ *UTFPR***Corresponding Author:** thiagocruz397@gmail.com

The Standard Model is highly successful in describing the behavior of particles and the nongravitational forces that govern their interactions. However, its limitations hinder our ability to explain certain phenomena, like the observed matter-antimatter asymmetry. Such limitations led to the development of extensions to the Standard Model. Unified theories, such as string theory and loop quantum gravity, suggest that Lorentz Invariance Violation may occur at the Planck Scale, but direct measurements at this scale are currently impossible. Nevertheless, this violation could be observable at a lower energy scale accessible to current experiments under the Standard Model Extension (SME) framework. Long-baseline experiments with neutrinos are crucial in advancing our understanding of physics beyond the Standard Model. Nevertheless, such studies mainly concern the normal hierarchy of neutrino masses. In this study, we investigate the influence of LIV parameters on the inverted hierarchy scenario. We present the probabilities, event rates, and sensitivity studies for long-baseline neutrino experiments.

Poster Session / 31

Muon-neutrino charged-current cross sections from MicroBooNE: first simultaneous measurements of final states with and without protons for Muon-neutrino scattering on argon

Author: Benjamin Bogart^{None}**Corresponding Author:** bbogart@umich.edu

A detailed understanding of muon neutrino charged-current interactions on argon is crucial to the study of neutrino oscillations in current and future experiments using liquid argon time projection chambers. To help fill this need, MicroBooNE has produced a comprehensive set of cross section measurements which simultaneously probe the leptonic and hadronic systems by dividing the inclusive channel into final states with and without protons. Data-driven model validation utilizing the conditional constraint formalism is employed to detect mismodeling that may bias the nominal flux averaged cross section results, which are extracted with the Wiener-SVD unfolding method. The results are compared to widely used event generator predictions revealing significant mismodeling of final states without protons, possibly due to insufficient treatment of final state interactions. These are first differential muon neutrino-argon cross section measurements made simultaneously for final states with and without protons, and provide novel information that will help stimulate the improvement of event generator modeling.

Poster Session / 32

Data-driven model validation for neutrino-Argon inclusive measurements at MicroBooNE

Author: London Cooper-troendle^{None}**Corresponding Author:** lcoopert@fnal.gov

Neutrino-nucleus cross section measurements are needed to improve interaction modeling to enable upcoming precision oscillation measurements and searches for physics beyond the standard model. There are two methods for extracting cross sections, which rely on using either the real or nominal flux prediction for the measurement. We examine the different challenges faced by these methods. Furthermore, the necessity for model validation in both procedures is addressed, and differences between “traditional” fake-data based validation and data-driven validation are discussed.

Data-driven model validation leverages goodness-of-fit tests enhanced by the conditional constraint procedure. This procedure aims to validate a model for a specific measurement so that any bias introduced in unfolding will be within the quoted uncertainties of the measurement. Results are shown for the first measurement of the 3D differential cross section $d^2\sigma(E_\nu)/d\cos(\theta_\mu)dP_\mu$ for inclusive muon-neutrino charged-current scattering on argon in MicroBooNE using a nominal-flux-prediction unfolding and data-driven model validation.

Poster Session / 33

GiBUU-Based Monte Carlo Simulation for Neutrino Experiments

Author: Leonidas Aliaga Soplin¹

¹ *University of Texas at Arlington*

Corresponding Author: leoliaga@gmail.com

This poster presents a Monte Carlo simulation implemented with the GiBUU model tailored for neutrino experiments. Specifically, we focus on its implementation, generating events in a generic liquid argon time projection chamber and comparing them with other neutrino event generators such as GENIE. The simulation generates realistic neutrino event samples, contributing to the prediction and interpretation of experimental outcomes. Our results demonstrate the robust performance of the GiBUU-based simulation framework and validate its performance against the original GiBUU cross-section model. Additionally, we outline our current efforts in developing infrastructure to address systematic uncertainties in the GiBUU model. By advancing simulation techniques, implementing neutrino-nucleus scattering models following different physics approaches, and building techniques to address systematic uncertainties, this work contributes to the refinement and reliability of neutrino experimental analyses.

Poster Session / 34

The development of the R-LDH Innovative media for O2 Capturing in Liquid Argon

Authors: Ana Amélia Machado¹; Ana Maria Caffer²; André Augusto³; Cezar Rodrigo Aguiar dos Santos¹; Cris Adriano¹; Cristhiane Reis⁴; Daniel Souza Correia⁵; David Montanari⁶; Dilson Cardoso⁷; Dirceu Noriler⁸; Elisabete Assaf⁷; Ettore Segreto¹; Frederico Demolin¹; G. S. Freitas¹; Gustavo Wiederhecker¹; H. B. Pizzi¹; Heriques Frandini¹; Italo O. Mazali⁹; Ivana Cruz⁷; José Mansur Assaf⁷; Krystal Brant⁵; Magda Fontes⁵; Mark Adamowski⁶; N. F. Oliveira Jr.¹⁰; P. G. Pagliuso¹; Pedro Bianchi⁸; Renato Soccol⁸; Rosembergue G. Gonçalves⁷; Roza Doubnik⁶; Thiago P. M. Alegre¹

¹ *Instituto de Física "Gleb Wataghin", UNICAMP*

² *UNICAMP*

³ *Equatorial sistemas S.A., Grupo Akaer*

⁴ *University of São João del Rei*

⁵ *Brazilian Center for Research in Physics*

⁶ *Fermi National Accelerator Laboratory*

⁷ *Advanced Materials and Energy Research Center, Federal University of São Carlos*

⁸ *School of Chemical Engineering, UNICAMP*

⁹ *Instituto de Química - UNICAMP*

¹⁰ *Instituto de Física, Universidade de São Paulo*

Corresponding Author: anamariacaffer@gmail.com

In this work, we have explored the potential of oxygen capturing in Liquid Argon (LAr) of the innovative CuO dispersive layered double hydroxide media (R-LDH). Low temperature experiments in the LAr Purification Cryostat (PuLArC) at IFGW/Unicamp were performed using LAr circulation through two filters, one containing the R-LDH material and the other the BASF commercial copper material (Cu-02265 - proposed as a reference O₂ getter media by Fermilab) for comparison. Interestingly, the experiments performed in PuLArC revealed that the R-LDH innovative media was capable of capturing O₂ from recirculating LAr in PuLArC, reducing the O₂ contaminants concentration to 80% of its initial values after 200 min of LAr circulation. As for the reference media BASF Cu-S0226, this media reduced the O₂ concentration to 40% of its initial value in the same time window. This result demonstrated a putative higher potential of the innovative R-LDH media for O₂ capturing in LAr and invoke further tests of this media in the PuLArC and in larger scale LAr cryostats, possibly at Fermilab and CERN.

Poster Session / 35

Evaluation of the Leggett-Garg inequality by means of the neutrino oscillations observed in reactor and accelerator experiments

Authors: Ricardo Zamora¹; Mario A. Acero²

¹ *Universidad del atlántico*

² *Universidad del Atlántico*

Corresponding Author: rjzamora@mail.uniatlantico.edu.co

The study of neutrino oscillation and its quantumness was studied by means of the Leggett-Garg inequality, which is based on the concept of macrorealism. This research considered the results from the Daya Bay and RENO reactor experiments, as well as the MINOS and NOvA accelerator experiments. It was found that Daya Bay and MINOS exhibit a strong violation of the Leggett-Garg inequality, while the indications from RENO and NOvA data are weaker.

The phenomenon of neutrino oscillation is significant in this context as it allows for the study of quantum phenomena, such as the survival probability of a neutrino flavor, at macroscopic distances. These studies are made possible by the aforementioned experiments.

The results demonstrate that the violation of the Leggett-Garg inequality is more pronounced for a smaller baseline-to-energy ratio in all the data sets considered. This is an important factor to consider when searching for evidence of quantum mechanical decoherence in neutrino oscillations. The findings imply that there is a characteristic value for each neutrino flavor beyond which the inequality is not satisfied.

Poster Session / 36

Development and application of the nuclear deexcitation simulator NucDeEx for precise prediction of neutrino-nuclear interactions

Author: Seisho Abe¹

¹ *Kamioka Observatory, ICRR, the University of Tokyo*

Corresponding Author: seisho@icrr.u-tokyo.ac.jp

In recent years, neutron multiplicity associated with neutrino-nucleus interactions has become important observable in large neutrino detectors such as Super-Kamiokande, KamLAND, and JUNO. The neutron multiplicity can be measured by detecting gamma rays emitted by neutron capture by taking delayed coincidence. It is expected to improve the results of various physics analyses by using the measured neutron multiplicity to enhance flavor identification or signal-to-background ratio. However, predicting neutron multiplicity is a challenging topic because neutrino-nucleus interactions involve highly uncertain nuclear effects.

Among the various processes involved in neutrino-nucleus interactions, nuclear deexcitation plays an important role in neutron multiplicity. This process emits various particles while transitioning to the ground state when the residual nucleus has exciting energy after the nucleon is knocked out. One issue is that most widely used neutrino interaction generators omit this process or describe it with a simplified model. Another issue is that the energy of deexcited particles is as low as a few MeV and, therefore, unobservable, i.e., un-constrainable, by most accelerator neutrino detectors due to higher detection thresholds. This feature of deexcitation requires us to rely on precise nuclear theory and experiments to verify it.

In this study, I developed a nuclear deexcitation simulator, NucDeEx, based on the nuclear calculation software TALYS. Since TALYS contains sophisticated nuclear models and parameters, NucDeEx can precisely simulate the nuclear deexcitation process. In addition, NucDeEx can be easily integrated with the neutrino interaction generators and other hadron simulators, such as Geant4 and the hadron cascade model INCL. The source code of NucDeEx and the interfaces and build scripts necessary for use with the above software are available on the web. Thus, a wide range of applications are expected. In this talk, I will present an overview of NucDeEx, its performance evaluation with nuclear experiments, the impact of integrating NucDeEx into neutrino interaction generators, and its application and prospects for other hadron simulators.

Paper: Phys. Rev. D 109, 036009

NucDeEx GitHub: <https://github.com/SeishoAbe/NucDeEx>

37

Bounds on Lorentz Violation Parameters: present and future

Author: Gabrieli Salmória¹

Co-authors: Luciana Hirsch¹; André Steklain¹; Isadora Martins¹; Pedro Bertazo¹

¹ UTFPR

Corresponding Author: gabrielisalmoria@alunos.utfpr.edu.br

String theory and loop quantum gravity suggest that Lorentz Invariance Violation may occur at the Planck scale. Nevertheless, this violation could be observable at a lower energy scale accessible to current experiments under the Standard Model Extension (SME) framework. This study aims to investigate and catalog current bounds for the Lorentz Invariance Violation (LIV) parameters using data from experiments. We also will study new bounds that new experiments will impose on these parameters. The methodology of this work is quantitative, using bibliographic research and data collection from academic materials and articles. This work is expected to result in a repository of parameter values for neutrino physics, facilitating data access and interpretation.

Poster Session / 38

Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method

Authors: Alan Hahn^{None}; Daniel Souza Correia^{None}; Ana Amelia Machado^{None}; David Montanari^{None}; Dilson Cardoso^{None}; Dirceu Noriler^{None}; Ettore Segreto^{None}; Flor de Maria Blaszczyk^{None}; Helio da Motta^{None}; Jose Mansur Assaf^{None}; Mark Adamowski^{None}; Pascoal Jose Giglio Pagliuso^{None}; Pedro Bianchi Neto^{None}; Renato Soccol Junior^{None}; Roza Doubnik^{None}; Sergey Koshelev^{None}; Thiago P. M. Alegre^{None}

Corresponding Author: danielsouzacorreia@hotmail.com

Liquid argon (LAr) is at the core of Time Projection Chamber (TPC) used in neutrino experiments. Nitrogen, oxygen, and water are the main contaminants in LAr that compromise the quality of physics a LArTPC can deliver. It is crucial to keep them as low as possible. In particular nitrogen is known to absorb the LAr scintillation light, one of the 2 key observables in LArTPCs. This poster outlines an experiment performed in the ICEBERG test stand at the Noble Liquid Test Facility (NLTF, Fermilab) to capture nitrogen from LAr using Li-FAU, an innovative method developed and previously tested in the Liquid Argon Purification Cryostat (PuLArC) at IFGW/Unicamp. Confirming the results obtained in PuLArC, this test shows that Li-FAU filtered the nitrogen injected multiple times into 2,625 L of LAr down to <1 ppm concentration over cycles of 96 h.

Poster Session / 39

Status of the Measurement of the Elastic and Quasi-elastic Scattering of Protons on Carbon Nuclei in EMPHATIC

Author: Robert Chirco¹

¹ EMPHATIC Collaboration

Corresponding Author: rchirco@hawk.iit.edu

In long-baseline neutrino oscillation experiments, Monte Carlo (MC) simulations based on hadron interactions and decays are used to predict the neutrino flux. The 10%-level systematic uncertainty of the predicted neutrino fluxes from these simulations is dominated by uncertainties in hadron interaction cross sections due to limited hadron scattering data. EMPHATIC aims to reduce the neutrino flux uncertainty by providing additional data. Using a table-top-sized spectrometer located at the Fermilab Test Beam Facility (FTBF), its physics program includes precise measurements of hadron scattering and production cross sections at various beam momenta and target species that are relevant for GeV-scale neutrino production. Using simulation, we have developed a simple single-track reconstruction algorithm that has a momentum resolution of 3-4%. We will demonstrate the progress in developing one of EMPHATIC's first track reconstruction algorithms – an important step in making a new single-track forward scattering measurement ($p + C \rightarrow p + C$ at several beam momenta) using Phase 1 data collected between 2022 and 2023.

Poster Session / 40

Probing light vector mediators with coherent scattering at future facilities

Author: Lucas Ramos¹

Co-authors: Enrico Bertuzzo²; Giovanni di Cortona

¹ IF-USP

² University of São Paulo

Corresponding Author: lucas.magno.ramos@usp.br

Future experiments dedicated to the detection of Coherent Elastic Neutrino-Nucleus Scattering may be powerful tools in probing light new physics. In this paper we study the sensitivity on light Z' mediators of two proposed experiments: a directional low pressure Time Projection Chamber detector, ν BDX-DRIFT, that will utilize neutrinos produced at the Long Baseline Neutrino Facility, and several possible experiments to be installed at the European Spallation Source. We compare the results obtained with existing limits from fixed-target, accelerator, solar neutrino and reactor experiments. Furthermore, we show that these experiments have the potential to test unexplored regions that, in some case, could explain the anomalous magnetic moment of the muon or peculiar spectral features in the cosmic neutrino spectrum observed by IceCube.

41

Ant colony in Sensitivity Studies

Author: Felipe Wieler^{None}

Co-authors: André Steklain ; Cássio Amador ; Isadora Martins ; Luiz Accorsi

Corresponding Authors: luizpauloavale@gmail.com, felipew@alunos.utfpr.edu.br, steklain@utfpr.edu.br

This work proposes an adaptation of the parameter verification method for CPT violation and CPT conservation in LIV parameters, as implemented in previous works by other authors using the GLOBES software. The adaptation will be based on the ant colony methodology (ACM) developed in the author's previous works.

The main objective is to reproduce the graphs and values obtained by other authors using the aforementioned ant colony methodology.

The ACM implementation will be calibrated, executed across various scenarios, and its results analyzed and compared to existing literature. The goal is to validate ACM as a tool for LIV parameter verification and efficient limit simulation.

Poster Session / 42

Nuclear effects in single pion production: hadron final state interactions

Author: Javier García-Marcos¹

Co-authors: Tania Franco Muñoz²; Matthias Hooft³; Kajetan Niewczas³; Alexis Nikolakopoulos⁴; Raul González Jiménez²; Natalie Jachowicz³; Jose Manuel Udias Moineo²

¹ *Universidad Complutense de Madrid and Ghent University*

² *Universidad Complutense de Madrid*

³ *Ghent University*

⁴ *FNAL*

Corresponding Author: javier31@ucm.es

Single pion production (SPP) is an important interaction mechanism in accelerator-based neutrino experiments, contributing significantly to total cross-sections and impacting precision measurements. Its study is essential for understanding neutrino oscillations, distinguishing signal from background, and refining theoretical models, which will help to reduce systematic errors in the neutrino energy reconstruction 1.

Therefore, a realistic SPP model that covers the whole energy range in neutrino experiments and accounts for final state interactions (FSI) is crucial. The current operator within the Hybrid model consists of several Feynmann diagrams including four nucleon resonances (P_{33} , D_{33} , P_{11} and S_{11}) and background terms from Chiral Perturbation Theory (ChPT) and an extension to the high energy region performed via Regge phenomenology [2]. This model is embedded in a sophisticated nuclear framework: the bound state is described as a Dirac solution within relativistic mean field (RMF) potentials. The final nucleon is in the relativistic distorted wave impulse approximation (RDWIA), a Dirac continuum state computed with energy dependent RMF potentials (EDRMF), so orthogonality between nucleon initial and final states is naturally implemented [3,4].

In recent work [5], we addressed the effect that the asymptotic approximation in the current operator has on inclusive and semi-inclusive cross sections. This is a step forward into a more complete and consistent treatment of SPP on the nucleus.

In a next step we implement FSI for the final pion, i.e., we describe the pion wave function with a RDWIA formalism. The pion continuum states are hence computed as solutions of the Klein-Gordon equation with a suitable nucleus-pion potential. The treatment of the final hadron wave functions,

with distortion effects from the residual nucleus, is the only way of taking into account elastic FSI within a consistent, fully relativistic and quantum-mechanical framework.

In this contribution we will present our latest results from Ref. [5] and new updates towards the inclusion of the pion distortion within our model.

- [1] L. Alvarez-Ruso *et al.*, Prog. Part. Nuc. Phys. 100 (2018)
- [2] R. González-Jiménez *et al.*, Phys. Rev. D 95, 113007 (2017)
- [3] R. González-Jiménez *et al.*, Phys. Rev. C 100, 045501 (2019)
- [4] A. Nikolakopoulos *et al.*, Phys. Rev. D 107, 053007 (2023)
- [5] J. García-Marcos *et al.*, Phys. Rev. C 109, 024608 (2024)

Poster Session / 43

Refining Neutrino Final State Interactions in carbon and lead: Insights from extrapolating between extremes

Author: Oscar Moreno Palacios^{None}

Corresponding Author: oemorenopalaci@email.wm.edu

Neutrino interactions with nucleons can lead to Final State Interactions (FSI), where the resulting hadrons scatter within the nucleus. However, current FSI models are often extrapolated to heavier nuclei, leading to possible discrepancies with experimental data. The MINERvA experiment employs five sets of nuclear targets, and it features an electromagnetic calorimeter (ECAL) situated at the back side of the detector, consisting of alternating layers of 2mm-thick lead and plastic scintillator. The substantial amount of lead interleaved within the ECAL, approximately 4 tons (fiducial) in total, offers the opportunity to isolate a high-statistics sample of neutrino-lead interactions. By analyzing the data collected by MINERvA's ECAL, we aim to improve our understanding of neutrino interactions with heavy nuclei and refine FSI modeling techniques. This poster provides insight into FSI model expectations scaling between hydrocarbon and lead, focusing on adapting simulation weights to mimic GENIE 3's A-dependent pion fate, with the MINERvA experiment's simulation baseline simulation based on GENIE 2.12.6. This ensures consistency with updated physics frameworks and enabling more accurate comparisons with experimental data.

Poster Session / 44

Studying Neutrino-Nucleus Interactions at SBND with Muon Neutrino CC Events

Author: Brinden Carlson¹

¹ University of Florida

Corresponding Author: bcarlson1@ufl.edu

The Short-Baseline Near Detector (SBND) is a 100-ton scale Liquid Argon Time Projection Chamber (LArTPC) neutrino detector positioned in the Booster Neutrino Beam (BNB) at Fermilab, as part of the Short-Baseline Neutrino (SBN) program. Located only 110 m from the neutrino production target, SBND is expected to record millions of neutrino interaction events every year allowing neutrino-argon cross-section measurements with unprecedented precision. This poster will present progress towards the first measurement of the muon-neutrino charged-current inclusive cross section in SBND.

Poster Session / 45

New developments in the Ghent Model for Single Pion Production

Author: Matthias Hooft¹

Co-authors: Javier García-Marcos²; Alexis Nikolakopoulos³; Natalie Jachowicz¹; Kajetan Niewczas¹

¹ *Ghent University*

² *Universidad Complutense de Madrid and Ghent University*

³ *FNAL*

Corresponding Author: matthias.hooft@ugent.be

Single-pion production constitutes an important contribution to neutrino-nucleus cross sections in the region regime covered by current and future accelerator-based neutrino-oscillation experiments. To analyse and interpret the data of these experiments, it is absolutely necessary to have good models for these cross sections available.

The Ghent model 1 for single-pion production consists of a low-energy part described by the three-level vertices of resonances on top of a background described by a Lagrangian from chiral perturbation theory (ChPT). This model can be extended to higher energies using Regge theory.

In order to improve the performance of the model, we reformulated our description in terms of helicity amplitudes and a multipole expansion. The advantage of this procedure, is that cross section contributions are separated in parts with fixed quantum numbers, providing more flexibility for further refinement of the model. Using this scheme makes it possible to effectively unitarize the whole background, taking into account the pion-nucleon scattering phases provided by pion-nucleon scattering data. Further, we modified the delta decay width and formfactors to incorporate Watson's theorem [2] and included meson exchange diagrams in the background, to obtain a better agreement with other models and data.

In this contribution we will present the results of these efforts in the vector sector. Work on implementing these developments in the axial part of neutrino-induced single pion production is in progress. In a next step, we will incorporate the updated description of the pion production off the nucleon in our nuclear model [3] to study the impact of these modifications on the total neutrino cross section and the comparison with data.

[1] R. González-Jiménez, N. Jachowicz, K. Niewczas, J. Nys, V. Pandey, T. Van Cuyck, and N. Van Dessel Phys. Rev. D 95, 113007 (2017)

[2] L. Alvarez-Ruso, E. Hernández, J. Nieves, and M.J. Vicente Vacas Physical Review D93, 014016 (2016)

[3] A. Nikolakopoulos, R. González-Jiménez, N. Jachowicz, and J.M. Udías Phys. Rev. D 107, 053007 (2023)

Poster Session / 46

Neutrons from CCQE-like Antineutrino Interactions in MINERvA's Various Nuclear Targets

Author: David Last^{None}

Corresponding Author: dlast@fnal.gov

The MINERvA collaboration has demonstrated both an ability to detect neutrons from GeV-scale antineutrino interactions, and entangled deficiencies in the modeling of their production and detection. These deficiencies present problems for oscillation experiments as misunderstanding the neutron content in the final state will bias energy estimators which assume a particular final state (e.g. CCQE-like) or rely on calorimetric information. This poster describes a cross section analysis to further the study of GeV-scale neutron production. The analysis studies the neutron production in

CCQE-like interactions across MINERvA's range of nuclear targets: carbon, iron, water, and lead. It will provide insight into the effect of the nuclear environment on neutron production and subsequent detection in antineutrino CCQE-like interactions, across a broad range of nuclei.

47

Influence of complex phases from off-diagonal parameters that violate Lorentz invariance in DUNE

Authors: Andre Steklain¹; Aretha Camila Mercuri¹; Isabel Ribas¹; João Victor Oliveira de Lima²; Luciana Hirsch¹; Nathan Bergmann³

¹ *UTFPR*

² *Universidade Tecnológica Federal do Paraná*

³ *Universidade Tecnológica Federal do Paraná*

Corresponding Author: aretha.camila@gmail.com

The existence of oscillating massive neutrinos poses a challenge to the standard model of particle physics (SM). Even standard oscillation theory cannot replicate the experimental results, such as the anomalies in the LSND and MiniBooNE data, pushing physicists to look for new alternatives, such as exploring physics Beyond the Standard Model (BSM). Utilizing a model involving Lorentz Invariance Violation (LIV) represents a possibility for understanding these phenomena. The fact that neutrinos are sensitive to measuring LIV is very relevant. To assess the numerical implications of a novel Hamiltonian introducing Lorentz-breaking parameters within the framework of the DUNE experiment, we employ the General Long Baseline Experiment Simulator (GLOBES) software. By modifying probability generators, we simulate the event count in the detector, focusing mainly on investigating the parameter phases from off-diagonal terms that conserve CPT symmetry while breaking Lorentz invariance.

Poster Session / 48

Professor Based ReWeight for GENIE Generator

Authors: Qiyu Yan^{None}; Costas Andreopoulos¹; Xianguo Lu²; Marco Roda¹; Julia Tena Vidal¹

¹ *University of Liverpool*

² *University of Oxford*

Corresponding Author: yanqiyu17@mails.ucas.ac.cn

Interaction generators for neutrinos are essential tools to predict the final states of neutrino interactions from atmospheric and accelerator sources. Those final states would be important input to quantify the relation between the visible energy in our detector and the neutrino energy, whose distribution is affected by oscillation. This understanding is crucial for experiments such as JUNO, DUNE, and Hyper-Kamiokande. GENIE is one of the neutrino generators that specialise in the GeV region. Much work is being done to tune GENIE models' parameters to obtain for the best description of experimental datasets. In addition, the parameters extracted from the tuning have very well motivated statistical uncertainties that will make the analyses more robust as based on better motivated inputs. Once the initial inputs are defined, it will be crucial to understand the how the uncertainties would affect the predictions. Reweighting is a powerful approach to propagate those model uncertainties through GENIE. There are many restrictions in the current reweight approaches, the main being that only a subset of parameters can be reweightable. This work aims to utilize the Professor tool to model GENIE as response functions. This approach unifies the workflow of tuning and reweight, allowing us to propagate the uncertainty obtained from the tuning using a reweight infrastructure. This will enable us to do reweight all parameters, including previously unweightable parameters, e.g. hadronization parameters.

Poster Session / 49**Hybrid model for single-pion production incorporated in the NuWro event generator nuclear framework**

Authors: Qiyu Yan^{None}; Kajetan Niewczas¹; Alexis Nikolakopoulos²; Raul González Jiménez³; Natalie Jachowicz¹; Xianguo Lu⁴; Jan Sobczyk^{None}

¹ *Ghent University*

² *FNAL*

³ *Universidad Complutense de Madrid*

⁴ *University of Warwick*

Corresponding Author: yanqiyu17@mails.ucas.ac.cn

Precise modeling of GeV neutrino interactions with nuclei underpins our understanding of data from atmospheric and accelerator neutrino experiments. Single pion production, a critical channel in these interactions, demands accurate representation as required by future experiments like DUNE and Hyper-K. The current NuWro single pion production model, while valuable, relies on a single Delta resonance and a non-resonant background obtained from Pythia. While Pythia excels in describing interactions with very high W values, its effectiveness diminishes at both lower and intermediate W values, leading to limitations in capturing single pion production accurately. To address these limitations, we propose a new single pion production model in NuWro. This model incorporates additional resonances and utilizes the Chiral Perturbation Theory (ChPT) method for a more comprehensive description of the non-resonant background, specifically focusing on the lower and intermediate W region. This refined approach demonstrably improves agreement with MINERvA transverse kinematic imbalance data compared to the previous model, paving the way for a more nuanced understanding of neutrino interactions.

Impact of scattering uncertainties on measurements / 50**Impact of neutrino cross section uncertainties on NOvA physics measurements**

Corresponding Authors: michael.dolce@tufts.edu, mdolce@fnal.gov

Impact of scattering uncertainties on measurements / 51**Impact of neutrino cross section uncertainties on MicroBooNE and SBN measurements**

Corresponding Author: lcoopert@fnal.gov

Neutrino interaction generators / 52**GENIE**

Corresponding Author: marco.roda@liverpool.ac.uk

Neutrino interaction generators / 53

NuWro

Corresponding Author: jan.sobczyk@uwr.edu.pl

Neutrino interaction generators / 54

ACHILLES

Corresponding Author: isaacson@fnal.gov

Neutrino interaction generators / 55

INCL+ABLA

Neutrino interaction generators / 56

NEUT

Corresponding Author: p.stowell@sheffield.ac.uk

Neutrino interaction generators / 57

GiBUU

Corresponding Authors: mosel@theo.physik.uni-giessen.de, mosel@physik.uni-giessen.de

Neutrino interaction generators / 58

Characterization of Neutral-current background induced by atmospheric neutrinos using neutrino event generators and the TALYS deexcitation package

Corresponding Authors: chengjie@ncepu.edu.cn, chengjie@ihep.ac.cn

Neutrino interaction generators / 59

High-energy neutrino-matter interaction cross-sections in neutrino event generators

Corresponding Author: alfonso.garciasoto@fas.harvard.edu

Neutrino Fluxes / 60

Flux Overview

Corresponding Author: yonehara@fnal.gov

Neutrino Fluxes / 61

Fermilab Beams

Corresponding Author: zarko@fnal.gov

Neutrino Fluxes / 62

J-PARC neutrino beam performance, monitoring and flux prediction, constraints and uncertainties

Corresponding Author: lucas.machado@na.infn.it

Neutrino Fluxes / 63

Spallation Fluxes

Corresponding Author: yefremen@utk.edu

Neutrino Fluxes / 64

NA61

Corresponding Authors: aolivier@fnal.gov, aolivie4@nd.edu

Neutrino Fluxes / 65

EMPHATIC

Corresponding Author: leoliaga@gmail.com

Neutrino Fluxes / 66

ENUBET

Corresponding Authors: andrea.longhin@unipd.it, andrea.longhin@lnf.infn.it

Neutrino Fluxes / 67

nu-e Scattering

Corresponding Author: lzazueta@fnal.gov

0pi / QE / 2p2h / 68

ICARUS

0pi / QE / 2p2h / 69

MicroBooNE - 1

Corresponding Author: afurmans@fnal.gov

0pi / QE / 2p2h / 70

MicroBooNE - 2

Corresponding Author: gardiner@fnal.gov

0pi / QE / 2p2h / 71

SBND

Corresponding Author: munjung@uchicago.edu

0pi / QE / 2p2h / 72

LIT+coupled cluster, SF from coupled cluster

Corresponding Author: jsobczyk@uni-mainz.de

0pi / QE / 2p2h / 73

short range correlations, generalized contact formalism

0pi / QE / 2p2h / 74

Two body currents in Hartree-Fock Mean Field framework

Corresponding Author: kajetan.niewczas@ugent.be

0pi / QE / 2p2h / 75

DWIA comparisons to recent measurements

Corresponding Authors: alexis.nikolakopoulos@ugent.be, anikolak@fnal.gov

76

Telegrapher equation for photon diffusion in LArTPCs with photon removal at the boundaries

Authors: Andre Steklain¹; Marcio Adames²

¹ *Universidade Tecnológica Federal do Parana*

² *UTFPR*

Corresponding Author: marcioadames@utfpr.edu.br

LArTPCs are important detectors in several experiments and light propagation predictions inside the LArTPC play an important role in their capabilities. This work explores an analytical approach to predict light propagation. We present an analytical solution to a relativistic photon diffusion equation in terms of the physical parameters relevant to the DUNE detectors' physics and account for photon absorption at the boundaries through physical considerations, instead of solving it as a boundary value problem. We then compare our results to Geant4 simulations and find similar outcomes.

0pi / QE / 2p2h / 77

MicroBooNE - 1

Corresponding Author: afurmans@fnal.gov

0pi / QE / 2p2h / 78

MicroBooNE - 2

Corresponding Author: gardiner@fnal.gov

0pi / QE / 2p2h / 79

ICARUS status and plans for measurements related to nu-Ar interactions with no mesons in the final state and/or QE- and 2p2h-like interactions

Corresponding Author: betan009@fnal.gov

0pi / QE / 2p2h / 80

SBND

Corresponding Author: munjung@uchicago.edu

0pi / QE / 2p2h / 81

Meson exchange currents in relativistic mean field

Corresponding Authors: tania.franco86@gmail.com, taniafra@ucm.es

0pi / QE / 2p2h / 82

Short time approximation

Corresponding Author: landreoli@wustl.edu

0pi / QE / 2p2h / 83

DWIA comparison to recent experimental data

Corresponding Author: anikolak@fnal.gov

0pi / QE / 2p2h / 84

Extended superscaling with two-particle emission in electron and neutrino scattering

Corresponding Author: palomacasale@ugr.es

Poster Session / 85

Improving neutrino reconstruction in LArTPCs by including charge and light signal

Author: Laura Paulucci¹

Co-authors: Flavio Cavanna²; Franciole Marinho ; Luis Gustavo Porto Paixao³

¹ UFABC

² Fermilab

³ UNICAMP

Corresponding Author: lpaulucc@fnal.gov

The use of liquid argon time projection chambers for neutrino experiments provides good tracking resolution and PID capabilities. Analysis are typically based on the ionization electron (charge) collection but argon is also a prolific scintillator. We discuss how to better model the combined charge and scintillation light emission in argon and how the use of both signals together can improve neutrino energy resolution and impact the sensibility to parameters such as neutrino delta-CP phase and mass hierarchy.

Poster Session / 86

Electrons for Neutrinos

Author: Julia Tena Vidal¹

¹ Tel Aviv University

Corresponding Author: jttenavidal@tauex.tau.ac.il

Future long-baseline neutrino oscillation experiments, such as the Deep Underground Neutrino Experiment (DUNE), aim to measure neutrino oscillation parameters with unprecedented precision. Such sensitivity demands precise characterization of the incoming neutrino energy, which can only be determined via comprehensive cross-section models. In particular DUNE, which flux peaks at 2.5~GeV, will be dominated by pion-production events. Far from ideal, current neutrino pion-production Monte Carlo simulations strongly rely on empirical models which fail to describe neutrino pion-production data at the few-GeV energy range on deuterium and heavier targets. The Electrons for Neutrinos collaboration (e4nu) proposes a novel alternative to leverage neutrino simulations by exploiting the similarities between electrons and neutrino interactions with matter. The e4nu collaboration is working towards a single-differential semi-inclusive pion production measurement with electron-scattering CLAS6 data at 1, 2, 4~GeV on Carbon. The results offer a new insight to exclusive pion-production properties and pion final-state-interactions in the nuclear environment and can be used to validate theory models and neutrino event-generators.

Poster Session / 87

The Hyper-K Intermediate Water Cherenkov Detector and the Water Cherenkov Test Experiment

Author: Mauricio Barbi¹

¹ University of Regina

Corresponding Author: barbi@uregina.ca

The Intermediate Water Cherenkov Detector (IWCD) is a planned component of the Hyper-K long baseline neutrino experiment with vertical mobility capable of scanning a region of the JPARC

accelerator-made neutrino beam spectrum. It will contribute to control systematic uncertainties arising from neutrino interaction cross sections and extrapolation of the beam flux between the Hyper-K near and far detectors. A key aspect of IWCD is its high granularity and fast time response thanks to an advanced photo-detection system based on multi-photomultiplier (mPMT) devices developed to deliver excellent Cherenkov ring reconstruction. The Water Cherenkov Test Experiment (WCTE) at CERN is designed to test this system, along with the capabilities of the calibration and reconstruction techniques planned for IWCD. It will use a beam produced at the CERN accelerator to also measure lepton interactions relevant to constraint uncertainties from physics models at a range of energies covering an important region of the IWCD phase space. This poster presents a short summary of both IWCD and WCTE, including their designs, plans and goals.

Poster Session / 88

Measuring muon antineutrino charged-current interactions without mesons in the final state, in the NOvA Near Detector

Author: Kevin Vockerodt¹

¹ *Queen Mary University of London*

Corresponding Author: k.j.vockerodt@qmul.ac.uk

NOvA is a long-baseline neutrino experiment based at Fermilab in the US, with the primary aim of measuring neutrino and antineutrino oscillations. This will enhance our understanding of electroweak interactions by measuring the neutrino mixing angles, CP-violating phase and neutrino mass ordering. To measure these oscillations, we first need to have a deep understanding of how neutrinos and antineutrinos interact with matter. Antineutrino interaction cross sections are, at present, particularly poorly constrained, and processes such as meson exchange currents are not well understood in the antineutrino sector.

This analysis will develop a cross-section measurement of muon antineutrino interactions without mesons (e.g. pions or kaons) in the final state, in the NOvA near detector. A high-statistics, high-purity sample is obtained through a cut-based selection process implementing machine learning techniques. The sample is dominated by quasi-elastic and meson exchange current interactions which are sensitive to nuclear effects such as final-state interactions. The cross section will be extracted as a function of the incoming neutrino energy and the kinematics of the outgoing particles.

Poster Session / 89

CEvNS and BSM physics with the CONNIE experiment.

Author: Pedro Ventura¹

Co-authors: Carla Bonifazi ; Irina Nasteva ²

¹ *IF-USP*

² *IF-UFRJ*

Corresponding Author: pedrozventura@gmail.com

The CONNIE experiment is dedicated to detecting Coherent Elastic Neutrino-Nucleus Scattering (CEvNS) of reactor antineutrinos using high-resistivity silicon CCDs at the Angra-2 reactor, and underwent significant advancements in 2021. Introducing two Skipper-CCD sensors with sub-electron readout noise, CONNIE became the pioneering project to employ Skipper-CCDs for reactor neutrino detection and achieved a record detection threshold of 15 eV. This work presents an overview of the sensors performance and the latest findings derived from 300 days of data collected during 2021-2022, totaling an exposure of 18.4 g-days. A comparison of event rates during reactor-on and off periods

reveals no excess, resulting in upper limits on the neutrino interaction rates at 95% confidence level, consistent with earlier findings. Additionally, searches for Physics Beyond the Standard Model yield constraints on simplified models with light vector mediators. New studies include a dark matter search employing diurnal modulation techniques and a quest for relativistic millicharged particles produced in the reactor. The promising outcomes achieved with a minimal sensor mass underscore the potential of Skipper-CCDs in probing rare neutrino interactions, motivating the future plans to expand the detector mass with a Multi-Chip-Module of Skipper-CCDs.

Pion Production/SIS/DIS / 90

Eta production from nucleons and nuclei

Pion Production/SIS/DIS / 91

Nuclear medium effects in ν - τ -A scattering at DUNE energies

Pion Production/SIS/DIS / 92

Neutrino-induced single pion production and the reanalyzed bubble chamber data

Corresponding Author: mariano@fisica.unlp.edu.ar

Pion Production/SIS/DIS / 93

Photo-, Electro- and Weak Single Pion Production Model

Corresponding Author: minoo.kabirnezhad@physics.ox.ac.uk

Pion Production/SIS/DIS / 94

Pion Production at MINERvA

Corresponding Authors: deborahh@yorku.ca, sultana@fnal.gov, mehreen@uga.edu

Pion Production/SIS/DIS / 95

ν_e and $\bar{\nu}_e$ low recoil in MINERvA

Corresponding Author: kevin@rochester.edu

Pion Production/SIS/DIS / 96

SIS Interactions in MINERvA

Corresponding Author: al3667@drexel.edu

Pion Production/SIS/DIS / 97

Pion Production in NOvA

Corresponding Author: wetstein@iastate.edu

Charged Particle/Testbeam Measurements / 98

e4nu

Corresponding Authors: jtenavidal@tauex.tau.ac.il, jtenavid@fnal.gov

Charged Particle/Testbeam Measurements / 99

Determination of the spectral functions of argon at JLab

Corresponding Authors: benhar@vt.edu, omar.benhar@roma1.infn.it

Charged Particle/Testbeam Measurements / 100

Electron Scattering in Generators

Corresponding Author: gardiner@fnal.gov

Charged Particle/Testbeam Measurements / 101

Empirical fits to inclusive electron-carbon scattering data obtained by deep-learning methods

Corresponding Author: krzysztof.graczyk@uwr.edu.pl

Charged Particle/Testbeam Measurements / 102

Charged particles interaction measurements in ProtoDUNE

Corresponding Authors: franciole@gmail.com, fmarinho@fnal.gov

Charged Particle/Testbeam Measurements / 103

ANNIE

Corresponding Author: mastbaum@physics.rutgers.edu

Low-energy Scattering / 104

Low-energy: Theory Overview in Inelastic and CEvNS Cross-sections and Impact of BSM Effects

Low-energy Scattering / 105

Low-energy Inelastic Cross-sections: Experiment Review and Future Prospects

Low-energy Scattering / 106

Neutrino-nucleus coherent scattering: status and prospects

Corresponding Author: enectali@northwestern.edu

Uncertainties and Prospects for Future Improvements / 107

Lattice QCD

Corresponding Author: asmeyer.physics@gmail.com

Uncertainties and Prospects for Future Improvements / 108

NUISANCE and global fits

Author: Clarence Wret¹

Co-authors: Callum Wilkinson²; Patrick Stowell³

¹ *Oxford*

² *Lawrence Berkeley National Laboratory*

³ *University of Sheffield*

Corresponding Authors: p.stowell@sheffield.ac.uk, cwilkinson@lbl.gov, clarence.wret@physics.ox.ac.uk

Uncertainties and Prospects for Future Improvements / 109

MINERvA form-factor measurement

Corresponding Authors: aolivier@fnal.gov, aolivie4@nd.edu

Uncertainties and Prospects for Future Improvements / 110

T2K Near Detector Upgrade

Corresponding Author: ulyse.virginet@lpnhe.in2p3.fr

Pion Production/SIS/DIS / 111

Resonance and QE contributions to inclusive nucleon structure functions

Corresponding Author: christy@jlab.org

Pion Production/SIS/DIS / 112

The Physics of SIS – TMC and HT

Corresponding Author: morfin@fnal.gov

Pion Production/SIS/DIS / 113

Superscaling in the inelastic region: The SuSAv2-inelastic model

Corresponding Author: jgroasa@us.es

Pion Production/SIS/DIS / 114

Neutrino Cross Sections for collider neutrinos

Corresponding Authors: yuseon.jeong@cern.ch, jeong.yuseon@gmail.com

Pion Production/SIS/DIS / 115

MicroBooNE

Corresponding Author: bbogart@umich.edu

Pion Production/SIS/DIS / 116

T2K latest results and prospects on NC and CC pion production

Corresponding Author: cesar.jesus-valls@ipmu.jp

Pion Production/SIS/DIS / 117

DUNE 2x2

Corresponding Authors: cyifan@slac.stanford.edu, yifanch@fnal.gov

Impact of scattering uncertainties on measurements / 118

Impact of Cross Section Uncertainties on BSM Searches

Corresponding Authors: jkopp@cern.ch, jkopp@fnal.gov

Closeout / 119

Experimental Summary

Corresponding Author: jpaley@fnal.gov

Closeout / 120

Theory Summary

Corresponding Author: artur.ankowski@uwr.edu.pl

Closeout / 121

Closing Remarks

Corresponding Author: natalie.jachowicz@ugent.be

0pi / QE / 2p2h / 122

MINERvA

Corresponding Author: druterbo@pas.rochester.edu

0pi / QE / 2p2h / 123

NOvA

Corresponding Authors: sfsanche@iastate.edu, sfsanche@fnal.gov

0pi / QE / 2p2h / 124

T2K

Corresponding Author: dolan@llr.in2p3.fr

Poster Session / 125

Double Beta Decay within the Relativistic QRPA

Authors: Lucas Carvalho dos Santos Maia¹; Cláudio de Conti²

Co-author: Arturo Rodolfo Samana¹

¹ Universidade Estadual de Santa Cruz

² Universidade Estadual Paulista

Corresponding Authors: arturo.samana@gmail.com, claudio.conti@unesp.br, lucacarvalho291@gmail.com

In nature there are about 50 nuclear systems where the single beta-decay is energetically forbidden, and double beta-decay turns out to be only possible mode of disintegration. It is the nuclear pairing force which causes such an “anomaly”, by making the mass of the odd-odd isobar, (N-1,Z+1), to be greater than the masses of its even-even neighbors, (N,Z) and (N-2,Z-2). The modes by which the double beta decay can take place are connected with the neutrino and antineutrino distinction. The Quasi-Particle Random Phase Approximation (QRPA) has turned out to be the most simple model for calculating the nuclear wave function involved in the single and double beta-decay transitions. In this work we perform a self-consistent relativistic QRPA (RQRPA) calculation of double beta-decay based on relativistic BCS (RBCS) mean field theory results for odd-odd intermediate nuclei ⁴⁸Sc, ⁷⁶As, ⁸²Br, ¹⁰⁰Tc, ¹²⁸I, and ¹³⁰I. We use the parameter set NL3 for interactions between protons, neutrons, σ , ω , ρ mesons and photons. The RQRPA equations are solved for the residual pion and rho interaction by employing the same parameters used in the RBCS, and experimental values for the pion and nucleons. The RQRPA results for the double beta-decay matrix elements are similar to those obtained within the QRPA and the shell model. Motivated by the results obtained here, we intend to calculate the neutrinoless decay in order to estimate the neutrino mass.

References

- F. Krmpotic, FIZIKA B 14 (2005) 2, 139.
- N. Paar, D. Vretenar, T. Marketin and P. Ring, Phys. Rev. C77 (2008) 024608.
- B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. 16 (1986) 1.
- C. De Conti, A. P. Galeão and F. Krmpotic', Phys. Lett. B 444 (1999) 14.
- C. De Conti, A. P. Galeão and F. Krmpotic', Phys. Lett. B 494 (2000) 46.
- D. Vretenar, N. Paar, P. Ring, and G. A. Lalazissis, Phys. Rev. C 63 (2001) 047301.

B. V. Carlson and D. Hirata, Phys. Rev. C 62 (2000) 054310.

Theoretical introduction / 126

Welcome

Corresponding Author: helio@cbpf.br

Theoretical introduction / 127

General introduction

Corresponding Author: deborahh@yorku.ca

Theoretical introduction / 128

General theory at the nucleon level

Corresponding Author: anikolak@fnal.gov

Theoretical introduction / 129

Introduction to nuclear effects

Corresponding Author: artur.ankowski@uwr.edu.pl

Theoretical introduction / 130

Relativistic mean field approach

Corresponding Author: anikolak@fnal.gov

Theoretical introduction / 131

Spectral function approach

Corresponding Author: artur.ankowski@uwr.edu.pl

Theoretical introduction / 132

Ab initio approach

Corresponding Author: landreoli@wustl.edu

Experimental introduction / 133

How to make a cross-section measurement

Corresponding Author: cherdack@fnal.gov

Experimental introduction / 134

Generator comparisons to data

Corresponding Authors: clarence.wret@physics.ox.ac.uk, cwilkinson@lbl.gov

Experimental introduction / 135

Hands-on exercises with generators

Corresponding Authors: clarence.wret@physics.ox.ac.uk, cwilkinson@lbl.gov

Introduction to Monte Carlo generators / 136

How generators work

Corresponding Author: kajetan.niewczas@ugent.be

Introduction to Monte Carlo generators / 137

Practicum: analyzing generator output

Author: Clarence Wret¹

Co-author: Callum Wilkinson²

¹ *Oxford*

² *Lawrence Berkeley National Laboratory*

Corresponding Authors: clarence.wret@physics.ox.ac.uk, cwilkinson@lbl.gov

Introduction to Monte Carlo generators / 138

Generators review

Corresponding Author: gardiner@fnal.gov

Introduction to Monte Carlo generators / 139

Practicum continued

Corresponding Authors: clarence.wret@physics.ox.ac.uk, cwilkinson@lbl.gov

0pi / QE / 2p2h / 140

Meson exchange currents in a relativistic mean field model

Corresponding Author: taniafra@ucm.es

Theoretical introduction / 141

Introduction to simulation

Corresponding Author: cwilkinson@lbl.gov

Introduction to Monte Carlo generators / 142

Practicum: running a generator

Corresponding Authors: clarence.wret@physics.ox.ac.uk, cwilkinson@lbl.gov

Opening / 143

Welcome

Corresponding Authors: peres.orlando@gmail.com, helio@fnal.gov

Opening / 144

ICTP-SAIFR

Opening / 145

Institute Principia

Registration / 146

Registration

Opening / 147

Information

Corresponding Authors: peres.orlando@gmail.com, helio@fnal.gov

Poster Winners / 148

Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method

Corresponding Author: danielsouzacorreia@hotmail.com

Poster Winners / 149

Ant colony in Sensitivity Studies

Corresponding Author: felipew@alunos.utfpr.edu.br

Poster Winners / 150

CEvNS and BSM physics with the CONNIE experiment.

Corresponding Author: pedrozventura@gmail.com

Discussion / 151

Discussion

Closeout / 152

Closing remarks