

Electron scattering from ^{12}C in the Short-Time Approximation

NUINT 2024, São Paulo, Brazil

April 17, 2024

Quantum Monte Carlo Group @ WashU

Jason Bub (GS) Graham Chambers-Wall (GS) Garrett King (GS)

Lorenzo Andreoli (PD) Abraham Flores (PD) Sam Novario (PD)

Saori Pastore and Maria Piarulli

Lorenzo Andreoli



Washington University in St. Louis



Outline



- Ab initio description of nuclei:
 - Nuclear interaction
 - GS wave function
 - Electroweak interaction of leptons with nucleons and clusters of correlated nucleons
- Short-time approximation
- Results
- Conclusions and outlook



Electron-nucleus scattering

Theoretical understanding of nuclear effects is extremely important for neutrino experimental programs: oscillation experiments require accurate calculations of cross sections

Electron scattering can be used to test our nuclear model: (same nuclear effects, no need to reconstruct energies, abundant experimental data)



Lepton-nucleus cross sections $\omega \sim 10^2$ MeV



Many-body nuclear problem

Many-body Nuclear Hamiltonian in coordinate space: Argonne v₁₈ + Urbana X

$$H = \sum_i T_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} v_{ijk}$$

$$\psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_A, s_1, s_2, \dots, s_A, t_1, t_2, \dots, t_A)$$

ψ are complex spin-isospin vectors in $3A$ dimensions with components $2^A \times \frac{A!}{Z!(A-Z)!}$

${}^4\text{He}$: 96

${}^6\text{Li}$: 1280

${}^8\text{Li}$: 14336

${}^{12}\text{C}$: 540572



Many-body nuclear problem

Many-body Nuclear Hamiltonian in coordinate space: Argonne v₁₈ + Urbana X

$$H = \sum_i T_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} v_{ijk}$$

$$\psi(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_A, s_1, s_2, \dots, s_A, t_1, t_2, \dots, t_A)$$

ψ are complex spin-isospin vectors in $3A$ dimensions with components $2^A \times \frac{A!}{Z!(A-Z)!}$



<http://exascaleage.org/np/>

- ${}^4\text{He}$: 96
- ${}^6\text{Li}$: 1280
- ${}^8\text{Li}$: 14336
- ${}^{12}\text{C}$: 540572



Many-body nuclear problem

Many-body Nuclear Hamiltonian in coordinate space: Argonne v₁₈ + Urbana X

$$H = \sum_i T_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} v_{ijk}$$

Quantum Monte Carlo method:
Use nuclear wave functions that minimize the expectation value of E

$$E_V = \frac{\langle \psi | H | \psi \rangle}{\langle \psi | \psi \rangle} \geq E_0$$

The evaluation is performed using Metropolis sampling



Nuclear Wave Functions

Variational wave function for nucleus in J state

$$|\psi\rangle = \mathcal{S} \prod_{i < j}^A \left[1 + U_{ij} + \sum_{k \neq i, j}^A U_{ijk} \right] \left[\prod_{i < j} f_c(r_{ij}) \right] |\Phi(JMTT_3)\rangle$$

Two-body spin- and isospin-dependent correlations

$$U_{ij} = \sum_p f^p(r_{ij}) O_{ij}^p$$

$$O_{ij}^p = [1, \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j, S_{ij}] \otimes [1, \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j]$$

$$U_{ijk} = \epsilon v_{ijk}(\bar{r}_{ij}, \bar{r}_{jk}, \bar{r}_{ki})$$

The trial wave function can be improved by eliminating spurious contaminations via 6 propagation in imaginary time (GFMC)



Nuclear Wave Functions

Variational wave function for nucleus in J state

$$|\psi\rangle = \mathcal{S} \prod_{i < j}^A \left[1 + U_{ij} + \sum_{k \neq i, j}^A U_{ijk} \right] \left[\prod_{i < j} f_c(r_{ij}) \right] |\Phi(JMTT_3)\rangle$$

Two-body spin- and isospin-dependent correlations

$$U_{ij} = \sum_p f^p(r_{ij}) O_{ij}^p$$

$$O_{ij}^p = [1, \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j, S_{ij}] \otimes [1, \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j]$$

$$U_{ijk} = \epsilon v_{ijk}(\bar{r}_{ij}, \bar{r}_{jk}, \bar{r}_{ki})$$

The trial wave function can be improved by eliminating spurious contaminations via 6 propagation in imaginary time (GFMC)



Nuclear Wave Functions

Variational wave function for nucleus in J state

$$|\psi\rangle = \mathcal{S} \prod_{i < j}^A \left[1 + U_{ij} + \sum_{k \neq i, j}^A U_{ijk} \right] \left[\prod_{i < j} f_c(r_{ij}) \right] |\Phi(JMTT_3)\rangle$$

Two-body spin- and isospin-dependent correlations

$$U_{ij} = \sum_p f^p(r_{ij}) O_{ij}^p$$

$$O_{ij}^p = [1, \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j, S_{ij}] \otimes [1, \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j]$$

$$U_{ijk} = \epsilon v_{ijk}(\bar{r}_{ij}, \bar{r}_{jk}, \bar{r}_{ki})$$

The trial wave function can be improved by eliminating spurious contaminations via 6 propagation in imaginary time (GFMC)



Nuclear Wave Functions

Variational wave function for nucleus in J state

$$|\psi\rangle = \mathcal{S} \prod_{i < j}^A \left[1 + U_{ij} + \sum_{k \neq i, j}^A U_{ijk} \right] \left[\prod_{i < j} f_c(r_{ij}) \right] |\Phi(JMTT_3)\rangle$$

Two-body spin- and isospin-dependent correlations

$$U_{ij} = \sum_p f^p(r_{ij}) O_{ij}^p$$

$$O_{ij}^p = [1, \boldsymbol{\sigma}_i \cdot \boldsymbol{\sigma}_j, S_{ij}] \otimes [1, \boldsymbol{\tau}_i \cdot \boldsymbol{\tau}_j]$$

$$U_{ijk} = \epsilon v_{ijk}(\bar{r}_{ij}, \bar{r}_{jk}, \bar{r}_{ki})$$


The trial wave function can be improved by eliminating spurious contaminations via 6 propagation in imaginary time (GFMC)



Many-body nuclear interaction

Many-body Nuclear Hamiltonian: Argonne v₁₈ + Urbana X

$$H = \sum_i T_i + \sum_{i < j} v_{ij} + \sum_{i < j < k} v_{ijk}$$







Electromagnetic interactions

Phenomenological Hamiltonian for NN and NNN

The interaction with external probes is described in terms on one- and two-body charge and current operators



Charge operators

$$\rho = \sum_{i=1}^A \rho_i + \sum_{i < j} \rho_{ij} + \dots$$

Current operators


$$\mathbf{j} = \sum_{i=1}^A \mathbf{j}_i + \sum_{i < j} \mathbf{j}_{ij} + \dots$$

Two-body currents are a manifestation of two-nucleon correlations



Electromagnetic interactions

- One body-currents: non-relativistic reduction of covariant nucleons' isoscalar and isovector currents
- Two-body currents: modeled on MEC currents constrained by commutation relation with the nuclear Hamiltonian (Pastore et al. PRC84(2011)024001, PRC87(2013)014006)
- Argonne v18 two-nucleon and Urbana potentials, together with these currents, provide a quantitatively successful description of many nuclear electroweak observables, including charge radii, electromagnetic moments and transition rates, charge and magnetic form factors of nuclei with up to $A = 12$ nucleons





Lepton-Nucleus scattering: Inclusive Processes

Electromagnetic Nuclear Response Functions

$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$

Longitudinal response induced by the charge operator $O_L = \rho$

Transverse response induced by the current operator $O_T = j$

5 responses in neutrino-nucleus scattering

$$\frac{d^2\sigma}{d\omega d\Omega} = \sigma_M [v_L R_L(\mathbf{q}, \omega) + v_T R_T(\mathbf{q}, \omega)]$$


One can exploit integral properties of the response functions to avoid explicit calculation of the final states: CC + Lorentz Integral Transform (see Joanna's talk), GFMC + Euclidean



Short-time approximation

S. Pastore, J. Carlson, S. Gandolfi, R. Schiavilla, and R. B. Wiringa PRC101(2020)044612

Describe electroweak scattering from $A >= 12$ without losing two-body physics, account for exclusive processes, Incorporate relativistic effects



Response functions


$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$



Short-time approximation

S. Pastore, J. Carlson, S. Gandolfi, R. Schiavilla, and R. B. Wiringa PRC101(2020)044612

Describe electroweak scattering from $A >= 12$ without losing two-body physics, account for exclusive processes, Incorporate relativistic effects



Response functions

$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$

$$R_\alpha(q, \omega) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i(\omega + E_i)t} \langle \Psi_i | O_\alpha^\dagger(\mathbf{q}) e^{-iHt} O_\alpha(\mathbf{q}) | \Psi_i \rangle$$


The sum over all final states is replaced by a two nucleon propagator



Short-time approximation

S. Pastore, J. Carlson, S. Gandolfi, R. Schiavilla, and R. B. Wiringa PRC101(2020)044612

Describe electroweak scattering from $A >= 12$ without losing two-body physics, account for exclusive processes, Incorporate relativistic effects



Response functions

$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$

$$R_\alpha(q, \omega) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i(\omega+E_i)t} \langle \Psi_i | O_\alpha^\dagger(\mathbf{q}) e^{-iHt} O_\alpha(\mathbf{q}) | \Psi_i \rangle$$

The sum over all final states is replaced by a two nucleon propagator


$$\begin{aligned} O^\dagger e^{-iHt} O &= \left(\sum_i O_i^\dagger + \sum_{i < j} O_{ij}^\dagger \right) e^{-iHt} \left(\sum_{i'} O_{i'} + \sum_{i' < j'} O_{i'j'} \right) \\ &= \sum_i O_i^\dagger e^{-iHt} O_i + \sum_{i \neq j} O_i^\dagger e^{-iHt} O_j \\ &\quad + \sum_{i \neq j} \left(O_i^\dagger e^{-iHt} O_{ij} + O_{ij}^\dagger e^{-iHt} O_i \right. \\ &\quad \left. + O_{ij}^\dagger e^{-iHt} O_{ij} \right) + \dots \end{aligned}$$



Short-time approximation

S. Pastore, J. Carlson, S. Gandolfi, R. Schiavilla, and R. B. Wiringa PRC101(2020)044612

Describe electroweak scattering from $A >= 12$ without losing two-body physics, account for exclusive processes, Incorporate relativistic effects



Response functions

$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$

$$R_\alpha(q, \omega) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i(\omega+E_i)t} \langle \Psi_i | O_\alpha^\dagger(\mathbf{q}) e^{-iHt} O_\alpha(\mathbf{q}) | \Psi_i \rangle$$

The sum over all final states is replaced by a two nucleon propagator


$$\begin{aligned} O^\dagger e^{-iHt} O &= \left(\sum_i O_i^\dagger + \sum_{i < j} O_{ij}^\dagger \right) e^{-iHt} \left(\sum_{i'} O_{i'} + \sum_{i' < j'} O_{i'j'} \right) \\ &= \sum_i O_i^\dagger e^{-iHt} O_i + \sum_{i \neq j} O_i^\dagger e^{-iHt} O_j \\ &\quad + \boxed{\sum_{i \neq j} (O_i^\dagger e^{-iHt} O_{ij} + O_{ij}^\dagger e^{-iHt} O_i)} \text{ Interference} \\ &\quad + O_{ij}^\dagger e^{-iHt} O_{ij} + \dots \end{aligned}$$



Short-time approximation

S. Pastore, J. Carlson, S. Gandolfi, R. Schiavilla, and R. B. Wiringa PRC101(2020)044612

Quasielastic scattering cross sections are expressed in terms of response function



Response functions

$$R_\alpha(q, \omega) = \sum_f \delta(\omega + E_0 - E_f) |\langle f | O_\alpha(\mathbf{q}) | 0 \rangle|^2$$


Response densities

$$R^{\text{STA}}(q, \omega) \sim \int \delta(\omega + E_0 - E_f) de dE_{cm} \mathcal{D}(e, E_{cm}; q)$$

STA: scattering of external probes from pairs of correlated nucleons



Transverse response density




Electron scattering from ${}^4\text{He}$ in the STA:

- Provides “more” exclusive information in terms of nucleon-pair kinematics via the Response Densities as functions of (E, e)
- Give access to particular kinematics for the struck nucleon pair



Back-to-back kinematic



We can select a particular kinematic, and assess the contributions from different particle identities



Benchmark

L.A, J. Carlson, A. Lovato, S. Pastore, N. Rocco, RB Wiringa PRC105(2022)014002

- We benchmarked three different methods based on the same description of nuclear dynamics of the initial target state
- Compared to the experimental data for the longitudinal and transverse electromagnetic response functions of ${}^3\text{He}$, and the inclusive cross sections of both ${}^3\text{He}$ and ${}^3\text{H}$
- Comparing the results allows for a precise quantification of the uncertainties inherent to factorization schemes



Benchmark

L.A. et al. PRC105(2022)014002

Green's function Monte Carlo


$$|\Psi_0\rangle \propto \lim_{\tau \rightarrow \infty} \exp[-(H - E_0)\tau] |\Psi_T\rangle$$

$$E_\alpha(\mathbf{q}, \tau) = \int_{\omega_{\text{th}}}^{\infty} d\omega e^{-\omega\tau} R_\alpha(\mathbf{q}, \omega), \quad \alpha = L, T$$

$$E_\alpha(\mathbf{q}, \tau) = \left\langle \Psi_0 \left| J_\alpha^\dagger(\mathbf{q}) e^{-(H - E_0)\tau} J_\alpha(\mathbf{q}) \right| \Psi_0 \right\rangle - |F_\alpha(\mathbf{q})|^2 e^{-\omega_{el}\tau}$$

Stort-time approximation

$$\begin{aligned} R_\alpha(\mathbf{q}, \omega) &= \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{i(\omega + E_0)t} \\ &\times \langle \Psi_0 | J_\alpha^\dagger(\mathbf{q}) e^{-iHt} J_\alpha(\mathbf{q}) | \Psi_0 \rangle \end{aligned}$$



Spectral function







$$|\Psi_f\rangle = |\mathbf{p}\rangle \otimes |\Psi_n^{A-1}\rangle$$

$$\begin{aligned} R_\alpha(\mathbf{q}, \omega) &= \sum_{\tau_k=p,n} \int \frac{d^3k}{(2\pi)^3} dE [P_{\tau_k}(\mathbf{k}, E) \\ &\times \frac{m_N^2}{e(\mathbf{k})e(\mathbf{k} + \mathbf{q})} \sum_i \langle k | j_{i,\alpha}^\dagger | k + q \rangle \langle p | j_{i,\alpha} | k \rangle \\ &\times \delta(\tilde{\omega} + e(\mathbf{k}) - e(\mathbf{k} + \mathbf{q}))] \end{aligned}$$



Benchmark







Longitudinal and transverse response function in ${}^3\text{He}$





Benchmark







Longitudinal and transverse response function in ${}^3\text{He}$





Cross sections







^3H





Cross sections

^3H





Relativistic corrections

Necessary to include relativistic correction at higher momentum q .

We are currently working on including relativistic corrections within the STA formalism:


R. Weiss, J. Carlson (LANL)

- Relativistic kinematic: allowed by STA factorization scheme
- Relativistic currents: expansion for a large value of the momentum transfer \mathbf{q}





Responses for ^{12}C

Longitudinal and transverse response for $300 < q < 800 \text{ MeV}$:





Back-to-back kinematic




We can select a particular kinematic, and assess the contributions from different particle identities




Two-body contributions

Transverse response density at $q=570$ MeV:











Cross sections results for ^{12}C



Preliminary



Cross sections results for ^{12}C






Preliminary






Cross sections results for ^{12}C

$E_e = 0.62 \text{ GeV}, \theta = 36^\circ$




$E_e = 1.108 \text{ GeV}, \theta = 37.5^\circ$





GENIE validation using e-scattering

Z = 2, A = 4, Beam Energy = 0.64 GeV, Angle = $60^\circ \pm 0.25^\circ$



- STA responses used to build the cross sections
- Cross sections are used to generate events in GENIE
- Electromagnetic processes (for which data are available) are used to validate the generator
- Next step: use response densities

$$\frac{d^2 \sigma}{d\omega d\Omega} = \sigma_M [v_L R_L(\mathbf{q}, \omega) + v_T R_T(\mathbf{q}, \omega)]$$


Barrow, Gardiner, Pastore, Betancourt et al. PRD 103 (2021) 5, 052001

GENIE HadronTensorModell Class: https://internal.dunescience.org/doxygen/classgenie_1_1HadronTensorModell.html



EW interactions:

- The current work on EM interactions allows for a thorough evaluation of the method, and a comparison with the abundant experimental data for electron-nucleus scattering
- **G. King**: neutral weak currents quasi-elastic responses evaluated for ${}^2\text{H}$





Conclusion:

- The STA responses for ^{12}C are in good agreement with the data, and are accurate up to moderate values of q (and consequently to moderate values of incoming electron beam for cross sections calculations)
- It can describe electroweak scattering from $A \geq 12$ without losing two-body physics, and is exportable to other QMC methods to address larger nuclei, e.g. AFDMC
- Incorporate relativistic effects, pion production
- Use of information from response densities in event generators: collaboration with GENIE Monte Carlo event generator (S. Gardiner, J. Barrow)

Collaborators:

G. King, S. Pastore, M. Piarulli

R. Weiss, J. Carlson

J. Barrow, M. Betancourt, S. Gardiner

Computational Resources awarded by the DOE ALCC,
INCITE, SciDAC, NERSC

Thank you!

 Washington University in St. Louis



 Fermilab

 URA