

# **NuInt 2024 - 14th International Workshop on Neutrino-Nucleus Interactions**



## **Report of Contributions**

Contribution ID: 1

Type: **not specified**

## Opening

**Presenters:** DA MOTTA FILHO, Helio (CBPF); PERES, Orlando (UNICAMP)

Contribution ID: 6

Type: **Poster**

## Alleviating the present tension between T2K and NO $\nu \nu A$ with neutrino New Physics at source

Since neutrino oscillation was observed, several experiments have been built to measure its parameters. At the present there is tension between T2K and NOVA. We propose a non-standard neutrino interaction at production. In this scenario, computed by quantum field theory formalism can make a better description of combined data. A new phase from this new interaction can make a role in search for violation of charge and parity symmetry.

**Primary authors:** CHERCHIGLIA, Adriano (Universidade Estadual de Campinas); Prof. PERES, Orlando Luis Goulart (UNICAMP); Mr SOUZA, Edson (University of Campinas); Dr RODRIGUES, Fernanda (University of Campinas); PASQUINI, Pedro Simoni (Unicamp); Mrs ROSSI, Rafaela (University of Campinas)

**Presenter:** Prof. PERES, Orlando Luis Goulart (UNICAMP)

**Session Classification:** Poster Session

Contribution ID: 7

Type: **Poster**

## Neutral and Charged forbidden contributions to $\nu-^{40}\text{Ar}$ cross sections

The contributions of neutral and charged forbidden and allowed multipole to  $\nu-^{40}\text{Ar}$  cross sections are evaluated within the QRPA and PQRPA models as function of the neutrino energy in the interval of interest for supernova neutrinos. These calculations are important as input for neutrino generator and simulator as the Marley code used for DUNE experiment.

**Primary author:** SAMANA, Arturo (Universidade Estadual de Santa Cruz)

**Presenter:** SAMANA, Arturo (Universidade Estadual de Santa Cruz)

**Session Classification:** Poster Session

Contribution ID: 8

Type: **Poster**

## Electron/positron beam identification analysis in FERMILAB

In High Energy Physics it is essential the study of the particles that make up everything (at least the known baryonic universe). Carrying out a study of these particles is necessary devices (detectors). These devices interact with particles through known physical processes and then, through a data acquisition system, one can proceed for further analysis. In the MINERvA experiment, the cross-section measurements are generally done. As a previous step, they used a mini test detector called Test Beam II (TB II) to see the response of new materials (objectives and scintillators) to different energy bands. In this work, the response of the Beam II test detector to electron and positron beams are studied. The response of this detector is studied for energies in the range of 2 to 8 GeV. A data/MC comparison is also performed to see discrepancies with the model used, as well as in the analysis of the behavior of electron and positron beams under different parameters (electromagnetic cascade opening angle, electromagnetic cascade starting module and energy absorbed by the calorimeter). Our method has allowed the differentiation of electron and positron beams, unequivocally for the energies in the range 2-6 GeV and, to a lesser extent, in the range 6-8 GeV. This is a genius result given that previous experiments did not obtain results that could differentiate well electron beams relative to positron beams in these beam energy ranges

**Primary authors:** Dr SOLANO SALINAS, Carlos Javier (UNMSM, Peru); Mr CHAVARRIA, Edgar (UNI, Peru)

**Presenter:** Dr SOLANO SALINAS, Carlos Javier (UNMSM, Peru)

**Session Classification:** Poster Session

Contribution ID: 9

Type: **Poster**

## **Parallelizing the computation of neutrino oscillation parameters in a general background medium, by using the Jacobi Method**

This work describes an alternative tool for the study of matter effects over neutrino propagation, by numerically describing the mixing amplitudes and mass splittings as a function of a single parameter. It allows the description of these observables in the same manner as it is commonly seen in 2-neutrino phenomenology, identifying resonant mixing and level crossing, except in a  $N$ -neutrino scenario. This is accomplished by sequentially applying the Jacobi diagonalization method over a defined and smooth path, in infinitesimal steps, transporting the information about the neutrino mixing and parametrization from one background to another. This transport occurs in the parameter space, thus being reliable even over spatial background discontinuities. Due to the nature of the neutrino mass-flavor mixing, it is possible to describe any background effects, in a model-independent way, by making use of the vacuum mixing only. This is computationally interesting since the workload can be split in two stages, separating the diagonalization from the physical interpretation and fitting of any model in question. The procedure is detailed and benchmarked against known analytical solutions, together with a complexity and computational-time analysis.

**Primary author:** Prof. VALDIVIESSO, Gustavo (Universidade Federal de Alfenas)

**Presenter:** Prof. VALDIVIESSO, Gustavo (Universidade Federal de Alfenas)

**Session Classification:** Poster Session

Contribution ID: 10

Type: **Poster**

## The search for Majorana neutrinos with nEXO

Neutrinos are the only known fermions with zero electric charge, and thus far the unique candidates to act as their own antimatter counterpart. If indeed a neutrino is its own antiparticle, we can find processes in which lepton number conservation, a global symmetry of the Standard Model, is violated. Such a discovery would have great implications in neutrino physics and their interactions. The most sensitive test for this condition, also known as Majorana neutrinos, is through the search of a hypothetical decay known as neutrinoless double beta decay. The primary focus of the nEXO Collaboration is the search for this process using a liquid xenon time projection chamber, at the tonne-scale rooted on the success of the EXO-200 experiment. Our projections result in a half-life sensitivity beyond  $10^{28}$  yr, sufficient to cover a milestone of this search consisting of the inverted ordering of neutrinos masses. This poster presents the search as well as the nEXO detector and its potential for discovery of new physics.

**Primary author:** LICCIARDI, Caio (Laurentian University)

**Presenter:** LICCIARDI, Caio (Laurentian University)

**Session Classification:** Poster Session

Contribution ID: 11

Type: **not specified**

## **Impact of interaction uncertainties on oscillation measurements**

*Monday, 15 April 2024 09:15 (30 minutes)*

**Presenter:** WRET, Clarence (Oxford)

**Session Classification:** Impact of scattering uncertainties on measurements



Contribution ID: 12

Type: **not specified**

## **Impact of interaction uncertainties on neutrino cross section measurements**

*Monday, 15 April 2024 09:45 (20 minutes)*

**Presenter:** PANDEY, Vishvas (Fermilab)

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 13

Type: **not specified**

## **Systematic uncertainty reduction and residual cross section uncertainties for PRISM**

*Monday, 15 April 2024 10:05 (20 minutes)*

**Presenter:** CARACAS, Ioana

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 14

Type: **not specified**

## **Impact of neutrino cross section uncertainties on T2K physics measurements**

*Monday, 15 April 2024 10:55 (20 minutes)*

**Presenter:** MUNTEANU, Laura (CERN)

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 15

Type: **not specified**

**TBA**

**Session Classification:** Impact of scattering uncertainties on measurements

**Track Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 18

Type: Poster

## Meson-exchange currents in quasielastic electron and neutrino scattering in a generalized superscaling approach

A method that enables the consistent inclusion of meson-exchange currents (MEC) within the framework of the superscaling analysis with relativistic effective mass will be presented. We use a novel definition of the single nucleon tensor, defined as the mean value of the single-nucleon responses by averaging with an energy distribution  $n()$  [1]. This single nucleon prefactor is obtained from the 1p-1h matrix element of the OB current combined with the two body current. The averaging definition is extended beyond the scaling region of the  $\psi$  variable characteristic of the Fermi gas modifying the momentum distribution through a smeared Fermi distribution [2].

In the generalized scaling analysis, the selected QE data generate a band with scatter of 20% at the maximum (highlighting the extent of scaling violation) that will be parametrized using a simple function  $f(\psi^*)$ .

Through the inclusion of MEC, we conducted a comparative analysis of the 1p-1h response functions within the context of the RFG, RMF and SuSAM  $\ast$  models. These responses can be expressed in a factorized way as the product of the averaged single nucleon multiplied by the scaling function. A reduction in the transverse response is observed, which is in accordance with previous calculations [3][4].

In this manner, the 1p-1h cross section including MEC is calculated and compared with the calculation without MEC and the experimental data [5][6]. Our examination shows that in the low momentum limit, the predictions of the relativistic model align with those of the non-relativistic model in Fermi gas. The formalism can be extended to calculate the 1p-1h neutrino nuclear responses by including the meson exchange currents using the models already proposed in electron scattering.

[1] P. R. Casale, J. E. Amaro, V. L. Martinez-Consentino and I. Ruiz Simo, Universe 9, no.4, 158 (2023) [arXiv:2303.03890 [nucl-th]].

[2] P. R. Casale, J. E. Amaro and M. B. Barbaro, Symmetry 15 (2023) no.9, 1709 [arXiv:2307.15783 [nucl-th]].

[3] J. E. Amaro, M. B. Barbaro, J. A. Caballero, T. W. Donnelly, C. Maieron and J. M. Udias, Phys. Rev. C 81 (2010), [arXiv:0906.5598 [nucl-th]].

[4] J. E. Amaro, A. M. Lallena and G. Co, Nucl. Phys. A 578 (1994), 365-396 doi:10.1016/0375-9474(94)90752-8

[5] Benhar, O.; Day, D.; Sick, I. An archive for quasi-elastic electron-nucleus scattering data [arXiv:nucl-ex/0603032]

[6] Benhar, O.; Day, D.; Sick, I. Available online: <https://www.faculty.virginia.edu/qes-archive/> C83 (2011) 045501.

**Primary authors:** AMARO, JOSE ENRIQUE (UNIV. GRANADA); RODRÍGUEZ CASALÉ, Paloma

**Presenter:** RODRÍGUEZ CASALÉ, Paloma

**Session Classification:** Poster Session

Contribution ID: 19

Type: **Poster**

## JLab Spectral Functions of Argon in NuWro and Their Implications for MicroBooNE

The Short-Baseline Neutrino program in Fermilab aims to resolve the nature of the low-energy excess events observed in LSND and MiniBooNE, and analyze with unprecedented precision neutrino interactions with argon, which requires reliable estimate of neutrino cross sections. We report updates of the NuWro generator that bring the state-of-the-art spectral functions to model the ground state properties of the argon nucleus, and improve the accuracy at low energies by accounting for effects of Coulomb potential and nuclear recoil. We discuss these developments in context of electron and neutrino interactions, by comparing NuWro predictions to experimental data from JLab Hall A and MicroBooNE. The MicroBooNE data are described with the Chi square per degree of freedom of 0.7, compared with 1.0 in the local Fermi gas model. Being obtained using the axial form factor parametrization from MINERvA, our results indicate a consistency between the CCQE measurements in MINERvA and MicroBooNE

**Primary author:** BANERJEE, Rwik Dharmapal (University of Wrocław)

**Co-authors:** Mr ANKOWSKI, ARTUR (UNIVERSITY OF WROCLAW); Mr SOBCZYK, JAN (UNIVERSITY OF WROCLAW); Mr PRASAD, HEMANT; Mr GRACZYK, Krzysztof; Mrs KOWAL, Beata

**Presenter:** BANERJEE, Rwik Dharmapal (University of Wrocław)

**Session Classification:** Poster Session

Contribution ID: 20

Type: **Poster**

## Probing Earth's internal structure with high energy neutrino transmission

The opacity of the Earth for the incidence of high energy neutrinos is directly related to the neutrino-nucleon cross sections and the description of the distribution of matter in the Earth's interior. In this work we investigate the sensitivity of neutrino transmission to the use of different models for Earth's density profile. In particular, we compare neutrino transmission using the PREM density model with three-, two- and one-layer models (core, mantle and crust) and constant densities. To calculate the neutrino-nucleon cross section we used the DGLAP predictions, with the CT14 parametrizations. Our results indicate that neutrino transmission is sensitive to different Earth profile density models at magnitudes that depend on the kinematic region tested and the models compared. We also show that the regeneration of the tau neutrino flux by its decay significantly impacts flux transmission in all models studied. The results achieved indicate that Earth profile density models impact the events observed at neutrino observatories such as IceCube.

**Primary authors:** FRANCENER, Reinaldo (UNICAMP); GONÇALVES, Victor (UFPEL)

**Presenter:** FRANCENER, Reinaldo (UNICAMP)

**Session Classification:** Impact of scattering uncertainties on measurements

**Track Classification:** Neutrino flux predictions and measurements



Contribution ID: 21

Type: **Poster**

## Random Forests for determining muon numbers in extensive air showers relative to their primary energy using KASCADE-Grande data

This study presented the development and application of two Random Forest (RF) models to KASCADE-Grande data with zenith angles  $\theta < 40^\circ$ , obtained from the KASCADE Cosmic Ray Data Centre. The aim was to predict the primary energy and correct the muon number for systematic effects on an event-by-event basis. Model training involved KCDC simulations using the high-energy hadronic interaction model QGSjet-II-04. To refine the RF models, we conducted feature engineering on the magnitudes and fine-tuned the hyperparameters. The results enabled us to estimate the number of muons relative to the primary energy in extensive air shower events across the energy range from 10 PeV to 1 EeV. A comparison was made between the experimental results and the expectations from the QGSJET-II-04 model, considering pure protons and iron nuclei, respectively.

**Primary author:** ASTO ROJAS, omar moises (Universidad Nacional de Ingeniería)

**Co-authors:** Dr ARTEAGA-VELÁZQUEZ, Juan Carlos (Instituto de Física y Matemáticas, Universidad Michoacana); SOLANO SALINAS, Carlos Javier (UNMSM, Peru)

**Presenter:** ASTO ROJAS, omar moises (Universidad Nacional de Ingeniería)

**Session Classification:** Poster Session

Contribution ID: 22

Type: **not specified**

**TBA**

**Session Classification:** Impact of scattering uncertainties on measurements

**Track Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 23

Type: **not specified**

**TBA**

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 24

Type: **not specified**

**TBA**

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 25

Type: **not specified**

**TBA**

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 27

Type: **Poster**

## Constraining Lorentz Invariance Violation Parameters Using Short and Long Baseline Experiments

The Lorentz Invariance is deeply connected to Special Relativity, which states that the laws of physics are the same for different observers in relative motion. It is the foundation of other successful theories, like quantum field theory, and connected to fundamental symmetries, like charge, parity, and time reversal (CPT), which is essential in the Standard Model of particle physics. Nevertheless, alternative theories proposing that Lorentz Invariance may break in some scales have been considered in the context of neutrino oscillations, as they can explain some anomalies present in experiments like LSND and MiniBoone. The dependence on baseline and energy can distinguish the influence of LIV on these anomalies in contrast with other effects, like Non-Standard Interactions (NSI). In this work, we perform simulation studies to study the influence of Lorentz-violating parameters in neutrino experiments combining two different baselines. We use the General Long Baseline Experiment Simulator (GLOBES) with a modified probability engine to include LIV parameters.

**Primary authors:** STEKLAIN, Andre (Universidade Tecnológica Federal do Parana); HIRSCH, Luciana (UTFPR)

**Presenter:** STEKLAIN, Andre (Universidade Tecnológica Federal do Parana)

Contribution ID: 28

Type: **Poster**

## **First Measurement of Double-Differential Charged Current $\nu\mu$ -Argon Scattering Cross Sections In Kinematic Imbalance Variables With The MicroBooNE Detector**

Making high-precision measurements of neutrino oscillation parameters requires an unprecedented understanding of neutrino-nucleus scattering. In this work, we present the first charged current double-differential cross sections in kinematic imbalance variables. These variables characterize both the transverse and total kinematic imbalance in a neutrino interaction. We use events with a single muon above 100 MeV/c, a single final state proton above 300 MeV/c, and no recorded final state pions. Thus, these variables act as a direct probe of nuclear effects such as final state interactions, Fermi motion, and multi-nucleon processes. Our measurement allows us to constrain systematic uncertainties associated with neutrino oscillation results performed by near-future experiments of the Short Baseline Neutrino (SBN) program, as well as by future large-scale experiments like DUNE.

**Primary author:** PAPADOPOULOU, Afroditi

**Presenter:** PAPADOPOULOU, Afroditi

**Session Classification:** Poster Session

Contribution ID: 29

Type: **Poster**

## Cross Section Systematics in DUNE

For the operation of precision neutrino experiments, the understanding of neutrino interactions with matter are preconditioned requirements of all detections and measurements of neutrinos. The largest uncertainties in estimating neutrino-nucleus interaction cross sections arise in the incomplete understanding of nuclear effects. In the study of neutrino oscillations and nuclear scattering processes, obtaining an interaction model with associated uncertainties is of substantial interest for the neutrino physics community. This report presents studies of simulated CC 2p-2h interactions, in which a neutrino interacts with a bound pair of nucleons. This interaction mode is very poorly constrained by current data. A comparison of three leading CC 2p-2h models is presented, along with a number of uncertainty parameters that have been implemented to account for model-to-model discrepancies in the DUNE oscillation analysis.

**Primary author:** BATHE-PETERS, Lars (University of Oxford)

**Presenter:** BATHE-PETERS, Lars (University of Oxford)

**Session Classification:** Poster Session



Contribution ID: 30

Type: **Poster**

## LIV Studies for Inverted Hierarchy Scenario

The Standard Model is highly successful in describing the behavior of particles and the nongravitational forces that govern their interactions. However, its limitations hinder our ability to explain certain phenomena, like the observed matter-antimatter asymmetry. Such limitations led to the development of extensions to the Standard Model. Unified theories, such as string theory and loop quantum gravity, suggest that Lorentz Invariance Violation may occur at the Planck Scale, but direct measurements at this scale are currently impossible. Nevertheless, this violation could be observable at a lower energy scale accessible to current experiments under the Standard Model Extension (SME) framework. Long-baseline experiments with neutrinos are crucial in advancing our understanding of physics beyond the Standard Model. Nevertheless, such studies mainly concern the normal hierarchy of neutrino masses. In this study, we investigate the influence of LIV parameters on the inverted hierarchy scenario. We present the probabilities, event rates, and sensitivity studies for long-baseline neutrino experiments.

**Primary author:** CRUZ, Thiago (Universidade Tecnológica Federal do Paraná)

**Co-authors:** THOMAZI, Angelo (Universidade Tecnológica Federal do Paraná); MADEIRA, Diego (Universidade Tecnológica Federal do Paraná); DE LIMA, Lucas (Universidade Tecnológica Federal do Paraná); MARCOLIN, Samuel (Universidade Tecnológica Federal do Paraná); WODZYNSKI, Johan (Universidade Tecnológica Federal do Paraná); Prof. STEKLAIN, Andre (Universidade Tecnológica Federal do Paraná); Prof. HIRSCH, Luciana (UTFPR)

**Presenter:** CRUZ, Thiago (Universidade Tecnológica Federal do Paraná)

Contribution ID: 31

Type: **Poster**

## **Muon-neutrino charged-current cross sections from MicroBooNE: first simultaneous measurements of final states with and without protons for Muon-neutrino scattering on argon**

A detailed understanding of muon neutrino charged-current interactions on argon is crucial to the study of neutrino oscillations in current and future experiments using liquid argon time projection chambers. To help fill this need, MicroBooNE has produced a comprehensive set of cross section measurements which simultaneously probe the leptonic and hadronic systems by dividing the inclusive channel into final states with and without protons. Data-driven model validation utilizing the conditional constraint formalism is employed to detect mismodeling that may bias the nominal flux averaged cross section results, which are extracted with the Wiener-SVD unfolding method. The results are compared to widely used event generator predictions revealing significant mismodeling of final states without protons, possibly due to insufficient treatment of final state interactions. These are first differential muon neutrino-argon cross section measurements made simultaneously for final states with and without protons, and provide novel information that will help stimulate the improvement of event generator modeling.

**Primary author:** BOGART, Benjamin

**Presenter:** BOGART, Benjamin

**Session Classification:** Poster Session

Contribution ID: 32

Type: **Poster**

## Data-driven model validation for neutrino-Argon inclusive measurements at MicroBooNE

Neutrino-nucleus cross section measurements are needed to improve interaction modeling to enable upcoming precision oscillation measurements and searches for physics beyond the standard model. There are two methods for extracting cross sections, which rely on using either the real or nominal flux prediction for the measurement. We examine the different challenges faced by these methods. Furthermore, the necessity for model validation in both procedures is addressed, and differences between “traditional” fake-data based validation and data-driven validation are discussed. Data-driven model validation leverages goodness-of-fit tests enhanced by the conditional constraint procedure. This procedure aims to validate a model for a specific measurement so that any bias introduced in unfolding will be within the quoted uncertainties of the measurement. Results are shown for the first measurement of the 3D differential cross section  $d^2\sigma(E_\nu)/d\cos(\theta_\mu)dP_\mu$  for inclusive muon-neutrino charged-current scattering on argon in MicroBooNE using a nominal-flux-prediction unfolding and data-driven model validation.

**Primary author:** COOPER-TROENDLE, London

**Presenter:** COOPER-TROENDLE, London

**Session Classification:** Poster Session

Contribution ID: 33

Type: **Poster**

## GiBUU-Based Monte Carlo Simulation for Neutrino Experiments

This poster presents a Monte Carlo simulation implemented with the GiBUU model tailored for neutrino experiments. Specifically, we focus on its implementation, generating events in a generic liquid argon time projection chamber and comparing them with other neutrino event generators such as GENIE. The simulation generates realistic neutrino event samples, contributing to the prediction and interpretation of experimental outcomes. Our results demonstrate the robust performance of the GiBUU-based simulation framework and validate its performance against the original GiBUU cross-section model. Additionally, we outline our current efforts in developing infrastructure to address systematic uncertainties in the GiBUU model. By advancing simulation techniques, implementing neutrino-nucleus scattering models following different physics approaches, and building techniques to address systematic uncertainties, this work contributes to the refinement and reliability of neutrino experimental analyses.

**Primary author:** Dr ALIAGA SOPLIN, Leonidas (University of Texas at Arlington)

**Presenter:** Dr ALIAGA SOPLIN, Leonidas (University of Texas at Arlington)

**Session Classification:** Poster Session

Contribution ID: 34

Type: **Poster**

## The development of the R-LDH Innovative media for O<sub>2</sub> Capturing in Liquid Argon

In this work, we have explored the potential of oxygen capturing in Liquid Argon (LAr) of the innovative CuO dispersive layered double hydroxide media (R-LDH). Low temperature experiments in the LAr Purification Cryostat (PuLArC) at IFGW/Unicamp were performed using LAr circulation through two filters, one containing the R-LDH material and the other the BASF commercial copper material (Cu-02265 - proposed as a reference O<sub>2</sub> getter media by Fermilab) for comparison. Interestingly, the experiments performed in PuLArC revealed that the R-LDH innovative media was capable of capturing O<sub>2</sub> from recirculating LAr in PuLArC, reducing the O<sub>2</sub> contaminants concentration to 80% of its initial values after 200 min of LAr circulation. As for the reference media BASF Cu-S0226, this media reduced the O<sub>2</sub> concentration to 40% of its initial value in the same time window. This result demonstrated a putative higher potential of the innovative R-LDH media for O<sub>2</sub> capturing in LAr and invoke further tests of this media in the PuLArC and in larger scale LAr cryostats, possibly at Fermilab and CERN.

**Primary authors:** MACHADO, Ana Amélia (Instituto de Física “Gleb Wataghin”, UNICAMP); Ms CAFFER, Ana Maria (UNICAMP); AUGUSTO, André (Equatorial sistemas S.A., Grupo Akaer); DOS SANTOS, Cezar Rodrigo Aguiar (Instituto de Física “Gleb Wataghin”, UNICAMP); Dr ADRIANO, Cris (Instituto de Física “Gleb Wataghin”, UNICAMP); Dr REIS, Cristhiane (University of São João del Rei); SOUZA CORREIA, Daniel (Brazilian Center for Research in Physics); MONTANARI, David (Fermi National Accelerator Laboratory); Dr CARDOSO, Dilson (Advanced Materials and Energy Research Center, Federal University of São Carlos); NORILER, Dirceu (School of Chemical Engineering, UNICAMP); Dr ASSAF, Elisabete (Advanced Materials and Energy Research Center, Federal University of São Carlos); Dr SEGRETO, Ettore (Instituto de Física “Gleb Wataghin”, UNICAMP); DEMOLIN, Frederico (Instituto de Física “Gleb Wataghin”, UNICAMP); Ms FREITAS, G. S. (Instituto de Física “Gleb Wataghin”, UNICAMP); WIEDERHECKER, Gustavo (Instituto de Física “Gleb Wataghin”, UNICAMP); Ms PIZZI, H. B. (Instituto de Física “Gleb Wataghin”, UNICAMP); FRANDINI, Heriques (Instituto de Física “Gleb Wataghin”, UNICAMP); Dr MAZALI, Italo O. (Instituto de Química - UNICAMP); CRUZ, Ivana (Advanced Materials and Energy Research Center, Federal University of São Carlos); Dr MANSUR ASSAF, José (Advanced Materials and Energy Research Center, Federal University of São Carlos); BRANT, Krystal (Brazilian Center for Research in Physics); Dr FONTES, Magda (Brazilian Center for Research in Physics); ADAMOWSKI, Mark (Fermi National Accelerator Laboratory); OLIVEIRA JR., N. F. ( Instituto de Física, Universidade de São Paulo); Dr PAGLIUSO, P. G. (Instituto de Física “Gleb Wataghin”, UNICAMP); BIANCHI, Pedro (School of Chemical Engineering, UNICAMP); SOCCOL, Renato (School of Chemical Engineering, UNICAMP); Dr GONÇALVES, Rosembergue G. (Advanced Materials and Energy Research Center, Federal University of São Carlos); DOUBNIK, Roza (Fermi National Accelerator Laboratory); Dr ALEGRE, Thiago P. M. (Instituto de Física “Gleb Wataghin”, UNICAMP)

**Presenter:** Ms CAFFER, Ana Maria (UNICAMP)

**Session Classification:** Poster Session

Contribution ID: 35

Type: **Poster**

## Evaluation of the Leggett-Garg inequality by means of the neutrino oscillations observed in reactor and accelerator experiments

The study of neutrino oscillation and its quantumness was studied by means of the Leggett-Garg inequality, which is based on the concept of macrorealism. This research considered the results from the Daya Bay and RENO reactor experiments, as well as the MINOS and NOvA accelerator experiments. It was found that Daya Bay and MINOS exhibit a strong violation of the Leggett-Garg inequality, while the indications from RENO and NOvA data are weaker.

The phenomenon of neutrino oscillation is significant in this context as it allows for the study of quantum phenomena, such as the survival probability of a neutrino flavor, at macroscopic distances. These studies are made possible by the aforementioned experiments.

The results demonstrate that the violation of the Leggett-Garg inequality is more pronounced for a smaller baseline-to-energy ratio in all the data sets considered. This is an important factor to consider when searching for evidence of quantum mechanical decoherence in neutrino oscillations. The findings imply that there is a characteristic value for each neutrino flavor beyond which the inequality is not satisfied.

**Primary authors:** ZAMORA, Ricardo (Universidad del atlántico); Mr ACERO, Mario A. (Universidad del Atlántico)

**Presenter:** ZAMORA, Ricardo (Universidad del atlántico)

**Session Classification:** Poster Session

Contribution ID: 36

Type: **Poster**

## Development and application of the nuclear deexcitation simulator NucDeEx for precise prediction of neutrino-nuclear interactions

In recent years, neutron multiplicity associated with neutrino-nucleus interactions has become important observable in large neutrino detectors such as Super-Kamiokande, KamLAND, and JUNO. The neutron multiplicity can be measured by detecting gamma rays emitted by neutron capture by taking delayed coincidence. It is expected to improve the results of various physics analyses by using the measured neutron multiplicity to enhance flavor identification or signal-to-background ratio. However, predicting neutron multiplicity is a challenging topic because neutrino-nucleus interactions involve highly uncertain nuclear effects.

Among the various processes involved in neutrino-nucleus interactions, nuclear deexcitation plays an important role in neutron multiplicity. This process emits various particles while transitioning to the ground state when the residual nucleus has exciting energy after the nucleon is knocked out. One issue is that most widely used neutrino interaction generators omit this process or describe it with a simplified model. Another issue is that the energy of deexcited particles is as low as a few MeV and, therefore, unobservable, i.e., un-constrainable, by most accelerator neutrino detectors due to higher detection thresholds. This feature of deexcitation requires us to rely on precise nuclear theory and experiments to verify it.

In this study, I developed a nuclear deexcitation simulator, NucDeEx, based on the nuclear calculation software TALYS. Since TALYS contains sophisticated nuclear models and parameters, NucDeEx can precisely simulate the nuclear deexcitation process. In addition, NucDeEx can be easily integrated with the neutrino interaction generators and other hadron simulators, such as Geant4 and the hadron cascade model INCL. The source code of NucDeEx and the interfaces and build scripts necessary for use with the above software are available on the web. Thus, a wide range of applications are expected. In this talk, I will present an overview of NucDeEx, its performance evaluation with nuclear experiments, the impact of integrating NucDeEx into neutrino interaction generators, and its application and prospects for other hadron simulators.

Paper: Phys. Rev. D 109, 036009

NucDeEx GitHub: <https://github.com/SeishoAbe/NucDeEx>

**Primary author:** ABE, Seisho (Kamioka Observatory, ICRR, the University of Tokyo)

**Presenter:** ABE, Seisho (Kamioka Observatory, ICRR, the University of Tokyo)

**Session Classification:** Poster Session

Contribution ID: 37

Type: **Poster**

## **Bounds on Lorentz Violation Parameters: present and future**

String theory and loop quantum gravity suggest that Lorentz Invariance Violation may occur at the Planck scale. Nevertheless, this violation could be observable at a lower energy scale accessible to current experiments under the Standard Model Extension (SME) framework. This study aims to investigate and catalog current bounds for the Lorentz Invariance Violation (LIV) parameters using data from experiments. We also will study new bounds that new experiments will impose on these parameters. The methodology of this work is quantitative, using bibliographic research and data collection from academic materials and articles. This work is expected to result in a repository of parameter values for neutrino physics, facilitating data access and interpretation.

**Primary author:** Mrs SALMÓRIA, Gabrieli (UTFPR)

**Co-authors:** Dr HIRSCH, Luciana (UTFPR); Dr STEKLAIN, André (UTFPR); Mrs MARTINS, Isadora (UTFPR); Mr BERTAZO, Pedro (UTFPR)

**Presenter:** Mrs SALMÓRIA, Gabrieli (UTFPR)



Contribution ID: 38

Type: **Poster**

## Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method

Liquid argon (LAr) is at the core of Time Projection Chamber (TPC) used in neutrino experiments. Nitrogen, oxygen, and water are the main contaminants in LAr that compromise the quality of physics a LArTPC can deliver. It is crucial to keep them as low as possible. In particular nitrogen is known to absorb the LAr scintillation light, one of the 2 key observables in LArTPCs. This poster outlines an experiment performed in the ICEBERG test stand at the Noble Liquid Test Facility (NLTF, Fermilab) to capture nitrogen from LAr using Li-FAU, an innovative method developed and previously tested in the Liquid Argon Purification Cryostat (PuLArC) at IFGW/Unicamp. Confirming the results obtained in PuLArC, this test shows that Li-FAU filtered the nitrogen injected multiple times into 2,625 L of LAr down to <1 ppm concentration over cycles of 96 h.

**Primary authors:** HAHN, Alan; SOUZA CORREIA, Daniel; MACHADO, Ana Amelia; MONTANARI, David; CARDOSO, Dilson; NORILER, Dirceu; SEGRETO, Ettore; BLASZCZYK, Flor de Maria; DA MOTTA, Helio; MANSUR ASSAF, Jose; ADAMOWSKI, Mark; GIGLIO PAGLIUSO, Pascoal Jose; BIANCHI NETO, Pedro; SOCCOL JUNIOR, Renato; DOUBNIK, Roza; KOSHELEV, Sergey; ALGRE, Thiago P. M.

**Presenter:** SOUZA CORREIA, Daniel

**Session Classification:** Poster Session

Contribution ID: 39

Type: **Poster**

## Status of the Measurement of the Elastic and Quasi-elastic Scattering of Protons on Carbon Nuclei in EMPHATIC

In long-baseline neutrino oscillation experiments, Monte Carlo (MC) simulations based on hadron interactions and decays are used to predict the neutrino flux. The 10%-level systematic uncertainty of the predicted neutrino fluxes from these simulations is dominated by uncertainties in hadron interaction cross sections due to limited hadron scattering data. EMPHATIC aims to reduce the neutrino flux uncertainty by providing additional data. Using a table-top-sized spectrometer located at the Fermilab Test Beam Facility (FTBF), its physics program includes precise measurements of hadron scattering and production cross sections at various beam momenta and target species that are relevant for GeV-scale neutrino production. Using simulation, we have developed a simple single-track reconstruction algorithm that has a momentum resolution of 3-4%. We will demonstrate the progress in developing one of EMPHATIC's first track reconstruction algorithms – an important step in making a new single-track forward scattering measurement ( $p + C \rightarrow p + C$  at several beam momenta) using Phase 1 data collected between 2022 and 2023.

**Primary author:** CHIRCO, Robert (EMPHATIC Collaboration)

**Presenter:** CHIRCO, Robert (EMPHATIC Collaboration)

**Session Classification:** Poster Session

Contribution ID: 40

Type: **Poster**

## Probing light vector mediators with coherent scattering at future facilities

Future experiments dedicated to the detection of Coherent Elastic Neutrino-Nucleus Scattering may be powerful tools in probing light new physics. In this paper we study the sensitivity on light  $Z'$  mediators of two proposed experiments: a directional low pressure Time Projection Chamber detector,  $\nu$ BDX-DRIFT, that will utilize neutrinos produced at the Long Baseline Neutrino Facility, and several possible experiments to be installed at the European Spallation Source. We compare the results obtained with existing limits from fixed-target, accelerator, solar neutrino and reactor experiments. Furthermore, we show that these experiments have the potential to test unexplored regions that, in some case, could explain the anomalous magnetic moment of the muon or peculiar spectral features in the cosmic neutrino spectrum observed by IceCube.

**Primary author:** RAMOS, Lucas (IF-USP)

**Co-authors:** BERTUZZO, Enrico (University of São Paulo); Dr DI CORTONA, Giovanni

**Presenter:** RAMOS, Lucas (IF-USP)

**Session Classification:** Poster Session

Contribution ID: 41

Type: **Poster**

## Ant colony in Sensitivity Studies

This work proposes an adaptation of the parameter verification method for CPT violation and CPT conservation in LIV parameters, as implemented in previous works by other authors using the GLoBES software. The adaptation will be based on the ant colony methodology (ACM) developed in the author's previous works.

The main objective is to reproduce the graphs and values obtained by other authors using the aforementioned ant colony methodology.

The ACM implementation will be calibrated, executed across various scenarios, and its results analyzed and compared to existing literature. The goal is to validate ACM as a tool for LIV parameter verification and efficient limit simulation.

**Primary author:** WIELER, Felipe

**Co-authors:** STEKLAIN, André; AMADOR, Cássio; MARTINS, Isadora; ACCORSI, Luiz

**Presenters:** WIELER, Felipe; STEKLAIN, André; AMADOR, Cássio; MARTINS, Isadora; ACCORSI, Luiz

Contribution ID: 42

Type: **Poster**

## Nuclear effects in single pion production: hadron final state interactions

Single pion production (SPP) is an important interaction mechanism in accelerator-based neutrino experiments, contributing significantly to total cross-sections and impacting precision measurements. Its study is essential for understanding neutrino oscillations, distinguishing signal from background, and refining theoretical models, which will help to reduce systematic errors in the neutrino energy reconstruction <sup>1</sup>.

Therefore, a realistic SPP model that covers the whole energy range in neutrino experiments and accounts for final state interactions (FSI) is crucial. The current operator within the Hybrid model consists of several Feynmann diagrams including four nucleon resonances ( $P_{33}$ ,  $D_{33}$ ,  $P_{11}$  and  $S_{11}$ ) and background terms from Chiral Perturbation Theory (ChPT) and an extension to the high energy region performed via Regge phenomenology [2]. This model is embedded in a sophisticated nuclear framework: the bound state is described as a Dirac solution within relativistic mean field (RMF) potentials. The final nucleon is in the relativistic distorted wave impulse approximation (RDWIA), a Dirac continuum state computed with energy dependent RMF potentials (EDRMF), so orthogonality between nucleon initial and final states is naturally implemented [3,4].

In recent work [5], we addressed the effect that the asymptotic approximation in the current operator has on inclusive and semi-inclusive cross sections. This is a step forward into a more complete and consistent treatment of SPP on the nucleus.

In a next step we implement FSI for the final pion, i.e., we describe the pion wave function with a RDWIA formalism. The pion continuum states are hence computed as solutions of the Klein-Gordon equation with a suitable nucleus-pion potential. The treatment of the final hadron wave functions, with distortion effects from the residual nucleus, is the only way of taking into account elastic FSI within a consistent, fully relativistic and quantum-mechanical framework.

In this contribution we will present our latest results from Ref. [5] and new updates towards the inclusion of the pion distortion within our model.

1 L. Alvarez-Ruso *et al.*, Prog. Part. Nuc. Phys. 100 (2018)

[2] R. González-Jiménez *et al.*, Phys. Rev. D 95, 113007 (2017)

[3] R. González-Jiménez *et al.*, Phys. Rev. C 100, 045501 (2019)

[4] A. Nikolakopoulos *et al.*, Phys. Rev. D 107, 053007 (2023)

[5] J. García-Marcos *et al.*, Phys. Rev. C 109, 024608 (2024)

**Primary author:** GARCÍA-MARCOS, Javier (Universidad Complutense de Madrid and Ghent University)

**Co-authors:** FRANCO MUÑOZ, Tania (Universidad Complutense de Madrid); HOOFT, Matthias (Ghent University); NIEWCZAS, Kajetan (Ghent University); NIKOLAKOPOULOS, Alexis (FNAL); GONZÁLEZ JIMÉNEZ, Raul (Universidad Complutense de Madrid); JACHOWICZ, Natalie (Ghent University); UDIAS MOINELO, Jose Manuel (Universidad Complutense de Madrid)

**Presenter:** GARCÍA-MARCOS, Javier (Universidad Complutense de Madrid and Ghent University)

**Session Classification:** Poster Session

Contribution ID: 43

Type: **Poster**

## Refining Neutrino Final State Interactions in carbon and lead: Insights from extrapolating between extremes

Neutrino interactions with nucleons can lead to Final State Interactions (FSI), where the resulting hadrons scatter within the nucleus. However, current FSI models are often extrapolated to heavier nuclei, leading to possible discrepancies with experimental data. The MINERvA experiment employs five sets of nuclear targets, and it features an electromagnetic calorimeter (ECAL) situated at the back side of the detector, consisting of alternating layers of 2mm-thick lead and plastic scintillator. The substantial amount of lead interleaved within the ECAL, approximately 4 tons (fiducial) in total, offers the opportunity to isolate a high-statistics sample of neutrino-lead interactions. By analyzing the data collected by MINERvA's ECAL, we aim to improve our understanding of neutrino interactions with heavy nuclei and refine FSI modeling techniques. This poster provides insight into FSI model expectations scaling between hydrocarbon and lead, focusing on adapting simulation weights to mimic GENIE 3's A-dependent pion fate, with the MINERvA experiment's simulation baseline simulation based on GENIE 2.12.6. This ensures consistency with updated physics frameworks and enabling more accurate comparisons with experimental data.

**Primary author:** MORENO PALACIOS, Oscar

**Presenter:** MORENO PALACIOS, Oscar

**Session Classification:** Poster Session

Contribution ID: 44

Type: **Poster**

## Studying Neutrino-Nucleus Interactions at SBND with Muon Neutrino CC Events

The Short-Baseline Near Detector (SBND) is a 100-ton scale Liquid Argon Time Projection Chamber (LArTPC) neutrino detector positioned in the Booster Neutrino Beam (BNB) at Fermilab, as part of the Short-Baseline Neutrino (SBN) program. Located only 110 m from the neutrino production target, SBND is expected to record millions of neutrino interaction events every year allowing neutrino-argon cross-section measurements with unprecedented precision. This poster will present progress towards the first measurement of the muon-neutrino charged-current inclusive cross section in SBND.

**Primary author:** CARLSON, Brinden (University of Florida)

**Presenter:** CARLSON, Brinden (University of Florida)

**Session Classification:** Poster Session

Contribution ID: 45

Type: **Poster**

## New developments in the Ghent Model for Single Pion Production

Single-pion production constitutes an important contribution to neutrino-nucleus cross sections in the region regime covered by current and future accelerator-based neutrino-oscillation experiments. To analyse and interpret the data of these experiments, it is absolutely necessary to have good models for these cross sections available.

The Ghent model 1 for single-pion production consists of a low-energy part described by the three-level vertices of resonances on top of a background described by a Lagrangian from chiral perturbation theory (ChPT). This model can be extended to higher energies using Regge theory.

In order to improve the performance of the model, we reformulated our description in terms of helicity amplitudes and a multipole expansion. The advantage of this procedure, is that cross section contributions are separated in parts with fixed quantum numbers, providing more flexibility for further refinement of the model. Using this scheme makes it possible to effectively unitarize the whole background, taking into account the pion-nucleon scattering phases provided by pion-nucleon scattering data. Further, we modified the delta decay width and formfactors to incorporate Watson's theorem [2] and included meson exchange diagrams in the background, to obtain a better agreement with other models and data.

In this contribution we will present the results of these efforts in the vector sector. Work on implementing these developments in the axial part of neutrino-induced single pion production is in progress. In a next step, we will incorporate the updated description of the pion production off the nucleon in our nuclear model [3] to study the impact of these modifications on the total neutrino cross section and the comparison with data.

[1] R. González-Jiménez, N. Jachowicz, K. Niewczas, J. Nys, V. Pandey, T. Van Cuyck, and N. Van Dessel Phys. Rev. D 95, 113007 (2017)

[2] L. Alvarez-Ruso, E. Hernández, J. Nieves, and M.J. Vicente Vacas Physical Review D93, 014016 (2016)

[3] A. Nikolakopoulos, R. González-Jiménez, N. Jachowicz, and J.M. Udías Phys. Rev. D 107, 053007 (2023)

**Primary author:** HOOFT, Matthias (Ghent University)

**Co-authors:** GARCÍA-MARCOS, Javier (Universidad Complutense de Madrid and Ghent University); NIKOLAKOPOULOS, Alexis (FNAL); JACHOWICZ, Natalie (Ghent University); NIEWCZAS, Kajetan (Ghent University)

**Presenter:** HOOFT, Matthias (Ghent University)

**Session Classification:** Poster Session



Contribution ID: 46

Type: **Poster**

## Neutrons from CCQE-like Antineutrino Interactions in MINERvA's Various Nuclear Targets

The MINERvA collaboration has demonstrated both an ability to detect neutrons from GeV-scale antineutrino interactions, and entangled deficiencies in the modeling of their production and detection. These deficiencies present problems for oscillation experiments as misunderstanding the neutron content in the final state will bias energy estimators which assume a particular final state (e.g. CCQE-like) or rely on calorimetric information. This poster describes a cross section analysis to further the study of GeV-scale neutron production. The analysis studies the neutron production in CCQE-like interactions across MINERvA's range of nuclear targets: carbon, iron, water, and lead. It will provide insight into the effect of the nuclear environment on neutron production and subsequent detection in antineutrino CCQE-like interactions, across a broad range of nuclei.

**Primary author:** LAST, David

**Presenter:** LAST, David

**Session Classification:** Poster Session

Contribution ID: 47

Type: **Poster**

## **Influence of complex phases from off-diagonal parameters that violate Lorentz invariance in DUNE**

The existence of oscillating massive neutrinos poses a challenge to the standard model of particle physics (SM). Even standard oscillation theory cannot replicate the experimental results, such as the anomalies in the LSND and MiniBooNE data, pushing physicists to look for new alternatives, such as exploring physics Beyond the Standard Model (BSM). Utilizing a model involving Lorentz Invariance Violation (LIV) represents a possibility for understanding these phenomena. The fact that neutrinos are sensitive to measuring LIV is very relevant. To assess the numerical implications of a novel Hamiltonian introducing Lorentz-breaking parameters within the framework of the DUNE experiment, we employ the General Long Baseline Experiment Simulator (GLOBES) software. By modifying probability generators, we simulate the event count in the detector, focusing mainly on investigating the parameter phases from off-diagonal terms that conserve CPT symmetry while breaking Lorentz invariance.

**Primary authors:** STEKLAIN, Andre (UTFPR); MERCURI, Aretha Camila (UTFPR); Mrs RIBAS, Isabel (UTFPR); OLIVEIRA DE LIMA, João Victor (Universidade Tecnológica Federal do Paraná); HIRSCH, Luciana (UTFPR); BERGMANN, Nathan (Universidade Tecnológica Federal do Paraná)

**Presenter:** MERCURI, Aretha Camila (UTFPR)

Contribution ID: 48

Type: **Poster**

## Professor Based ReWeight for GENIE Generator

Interaction generators for neutrinos are essential tools to predict the final states of neutrino interactions from atmospheric and accelerator sources. Those final states would be important input to quantify the relation between the visible energy in our detector and the neutrino energy, whose distribution is affected by oscillation. This understanding is crucial for experiments such as JUNO, DUNE, and Hyper-Kamiokande. GENIE is one of the neutrino generators that specialise in the GeV region. Much work is being done to tune GENIE models' parameters to obtain for the best description of experimental datasets. In addition, the parameters extracted from the tuning have very well motivated statistical uncertainties that will make the analyses more robust as based on better motivated inputs. Once the initial inputs are defined, it will be crucial to understand the how the uncertainties would affect the predictions. Reweighting is a powerful approach to propagate those model uncertainties through GENIE. There are many restrictions in the current reweight approaches, the main being that only a subset of parameters can be reweightable. This work aims to utilize the Professor tool to model GENIE as response functions. This approach unifies the workflow of tuning and reweight, allowing us to propagate the uncertainty obtained from the tuning using a reweight infrastructure. This will enable us to do reweight all parameters, including previously unweightable parameters, e.g. hadronization parameters.

**Primary authors:** YAN, Qiyu; ANDREOPOULOS, Costas (University of Liverpool); LU, Xianguo (University of Oxford); RODA, Marco (University of Liverpool); TENA VIDAL, Julia (University of Liverpool)

**Presenter:** YAN, Qiyu

**Session Classification:** Poster Session

Contribution ID: 49

Type: **Poster**

## Hybrid model for single-pion production incorporated in the NuWro event generator nuclear framework

Precise modeling of GeV neutrino interactions with nuclei underpins our understanding of data from atmospheric and accelerator neutrino experiments. Single pion production, a critical channel in these interactions, demands accurate representation as required by future experiments like DUNE and Hyper-K. The current NuWro single pion production model, while valuable, relies on a single Delta resonance and a non-resonant background obtained from Pythia. While Pythia excels in describing interactions with very high  $W$  values, its effectiveness diminishes at both lower and intermediate  $W$  values, leading to limitations in capturing single pion production accurately. To address these limitations, we propose a new single pion production model in NuWro. This model incorporates additional resonances and utilizes the Chiral Perturbation Theory (ChPT) method for a more comprehensive description of the non-resonant background, specifically focusing on the lower and intermediate  $W$  region. This refined approach demonstrably improves agreement with MINERvA transverse kinematic imbalance data compared to the previous model, paving the way for a more nuanced understanding of neutrino interactions.

**Primary authors:** YAN, Qiyu; NIEWCZAS, Kajetan (Ghent University); NIKOLAKOPOULOS, Alexis (FNAL); GONZÁLEZ JIMÉNEZ, Raul (Universidad Complutense de Madrid); JACHOWICZ, Natalie (Ghent University); LU, Xianguo (University of Warwick); SOBCZYK, Jan

**Presenter:** YAN, Qiyu

**Session Classification:** Poster Session

Contribution ID: 50

Type: **not specified**

## **Impact of neutrino cross section uncertainties on NOvA physics measurements**

*Monday, 15 April 2024 11:35 (20 minutes)*

**Presenters:** DOLCE, Michael; DOLCE, Michael

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 51

Type: **not specified**

## **Impact of neutrino cross section uncertainties on MicroBooNE and SBN measurements**

*Monday, 15 April 2024 11:15 (20 minutes)*

**Presenter:** COOPER-TROENDLE, London

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: 52

Type: **not specified**

## GENIE

*Monday, 15 April 2024 14:40 (20 minutes)*

**Presenter:** RODA, Marco (University of Liverpool)

**Session Classification:** Neutrino interaction generators

Contribution ID: 53

Type: **not specified**

## **NuWro**

*Monday, 15 April 2024 14:20 (20 minutes)*

**Presenter:** SOBCZYK, Jan

**Session Classification:** Neutrino interaction generators



Contribution ID: 54

Type: **not specified**

# ACHILLES

*Monday, 15 April 2024 15:50 (25 minutes)*

**Presenter:** ISAACSON, Joshua (FNAL)

**Session Classification:** Neutrino interaction generators

Contribution ID: 55

Type: **not specified**

## INCL+ABLA

*Monday, 15 April 2024 17:05 (25 minutes)*

**Presenter:** Dr ERSHOVA, Anna

**Session Classification:** Neutrino interaction generators

Contribution ID: 56

Type: **not specified**

## NEUT

*Monday, 15 April 2024 15:00 (20 minutes)*

**Presenter:** STOWELL, Patrick (University of Sheffield)

**Session Classification:** Neutrino interaction generators

Contribution ID: 57

Type: **not specified**

## GiBUU

*Monday, 15 April 2024 14:00 (20 minutes)*

**Presenters:** MOSEL, Ulrich (University of Giessen); MOSEL, Ulrich (Universitaet Giessen)

**Session Classification:** Neutrino interaction generators

Contribution ID: 58

Type: **not specified**

## **Characterization of Neutral-current background induced by atmospheric neutrinos using neutrino event generators and the TALYS deexcitation package**

*Monday, 15 April 2024 16:40 (25 minutes)*

**Presenters:** CHENG, Jie; CHENG, Jie (IHEP)

**Session Classification:** Neutrino interaction generators

Contribution ID: 59

Type: **not specified**

## **High-energy neutrino-matter interaction cross-sections in neutrino event generators**

*Monday, 15 April 2024 16:15 (25 minutes)*

**Presenter:** GARCIA SOTO, Alfonso Andres (Harvard University)

**Session Classification:** Neutrino interaction generators

Contribution ID: **60**

Type: **not specified**

## Flux Overview

*Tuesday, 16 April 2024 08:45 (20 minutes)*

**Presenter:** YONEHARA, Katsuya (Fermilab)

**Session Classification:** Neutrino Fluxes

Contribution ID: **61**

Type: **not specified**

## Fermilab Beams

*Tuesday, 16 April 2024 09:05 (20 minutes)*

**Presenter:** PAVLOVIC, Zarko (Fermilab)

**Session Classification:** Neutrino Fluxes



Contribution ID: 62

Type: **not specified**

## **J-PARC neutrino beam performance, monitoring and flux prediction, constraints and uncertainties**

*Tuesday, 16 April 2024 09:25 (20 minutes)*

**Presenter:** NASCIMENTO MACHADO, Lucas (University of Naples - INFN Naples)

**Session Classification:** Neutrino Fluxes

Contribution ID: **63**

Type: **not specified**

## Spallation Fluxes

*Tuesday, 16 April 2024 09:45 (20 minutes)*

**Presenter:** EFREMENKO, Yuri (University of Tennessee)

**Session Classification:** Neutrino Fluxes

Contribution ID: **64**

Type: **not specified**

## **NA61**

*Tuesday, 16 April 2024 10:35 (20 minutes)*

**Presenters:** OLIVIER, Andrew; OLIVIER, Andrew (University of Notre Dame)

**Session Classification:** Neutrino Fluxes

Contribution ID: 65

Type: **not specified**

## EMPHATIC

*Tuesday, 16 April 2024 10:55 (20 minutes)*

**Presenter:** ALIAGA SOPLIN, Leonidas (University of Texas at Arlington)

**Session Classification:** Neutrino Fluxes

Contribution ID: **66**

Type: **not specified**

## ENUBET

*Tuesday, 16 April 2024 11:15 (20 minutes)*

**Presenters:** LONGHIN, Andrea (University and INFN Padova); LONGHIN, Andrea (INFN LNF)

**Session Classification:** Neutrino Fluxes

Contribution ID: 67

Type: **not specified**

## nu-e Scattering

*Tuesday, 16 April 2024 11:35 (20 minutes)*

**Presenter:** ZAZUETA, Luis (Syracuse University)

**Session Classification:** Neutrino Fluxes

Contribution ID: **68**

Type: **not specified**

## ICARUS

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **69**

Type: **not specified**

## **MicroBooNE - 1**

**Presenter:** FURMANSKI, Andrew

**Session Classification:** 0pi / QE / 2p2h



Contribution ID: 70

Type: **not specified**

## MicroBooNE - 2

**Presenter:** GARDINER, Steven (Fermilab)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 71

Type: **not specified**

## SBND

**Presenter:** JUNG, Mun Jung (the University of Chicago)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 72

Type: **not specified**

## **LIT+coupled cluster, SF from coupled cluster**

*Tuesday, 16 April 2024 16:00 (25 minutes)*

**Presenter:** SOBCZYK, Joanna (JGU Mainz)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 73

Type: **not specified**

## **short range correlations, generalized contact formalism**

*Tuesday, 16 April 2024 16:25 (25 minutes)*

**Presenter:** WEISS, Ronen

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 74

Type: **not specified**

## Two body currents in Hartree-Fock Mean Field framework

*Tuesday, 16 April 2024 17:15 (25 minutes)*

**Presenter:** NIEWCZAS, Kajetan (Ghent University)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 75

Type: **not specified**

## **DWIA comparisons to recent measurements**

**Presenters:** NIKOLAKOPOULOS, Alexis (UGent); NIKOLAKOPOULOS, Alexis (FNAL)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 76

Type: **Poster**

## Telegrapher equation for photon diffusion in LArTPCs with photon removal at the boundaries

LArTPCs are important detectors in several experiments and light propagation predictions inside the LArTPC play an important role in their capabilities. This work explores an analytical approach to predict light propagation. We present an analytical solution to a relativistic photon diffusion equation in terms of the physical parameters relevant to the DUNE detectors' physics and account for photon absorption at the boundaries through physical considerations, instead of solving it as a boundary value problem. We then compare our results to Geant4 simulations and find similar outcomes.

**Primary authors:** STEKLAIN, Andre (Universidade Tecnologica Federal do Parana); ADAMES, Marcio (UTFPR)

**Presenter:** ADAMES, Marcio (UTFPR)

Contribution ID: 77

Type: **not specified**

## **MicroBooNE - 1**

*Wednesday, 17 April 2024 08:45 (20 minutes)*

**Presenter:** FURMANSKI, Andrew

**Session Classification:** 0pi / QE / 2p2h



Contribution ID: 78

Type: **not specified**

## MicroBooNE - 2

*Wednesday, 17 April 2024 09:05 (20 minutes)*

**Presenter:** GARDINER, Steven (Fermilab)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 79

Type: **not specified**

## **ICARUS status and plans for measurements related to $\nu$ -Ar interactions with no mesons in the final state and/or QE- and 2p2h-like interactions**

*Wednesday, 17 April 2024 09:25 (30 minutes)*

**Presenter:** BETANCOURT, Minerba (Fermilab)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **80**

Type: **not specified**

## **SBND**

*Wednesday, 17 April 2024 09:55 (20 minutes)*

**Presenter:** JUNG, Mun Jung (the University of Chicago)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **81**

Type: **not specified**

## **Meson exchange currents in relativistic mean field**

**Presenters:** FRANCO MUÑOZ, Tania (Universidad Complutense de Madrid); FRANCO MUÑOZ, Tania (Universidad Complutense de Madrid)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **82**

Type: **not specified**

## **Short time approximation**

*Wednesday, 17 April 2024 10:45 (25 minutes)*

**Presenter:** ANDREOLI, Lorenzo

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **83**

Type: **not specified**

## **DWIA comparison to recent experimental data**

*Wednesday, 17 April 2024 11:35 (25 minutes)*

**Presenter:** NIKOLAKOPOULOS, Alexis (FNAL)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 84

Type: **not specified**

## **Extended superscaling with two-particle emission in electron and neutrino scattering**

*Wednesday, 17 April 2024 11:10 (25 minutes)*

**Presenter:** RODRÍGUEZ CASALÉ, Paloma

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 85

Type: **Poster**

## Improving neutrino reconstruction in LArTPCs by including charge and light signal

The use of liquid argon time projection chambers for neutrino experiments provides good tracking resolution and PID capabilities. Analysis are typically based on the ionization electron (charge) collection but argon is also a prolific scintillator. We discuss how to better model the combined charge and scintillation light emission in argon and how the use of both signals together can improve neutrino energy resolution and impact the sensibility to parameters such as neutrino delta-CP phase and mass hierarchy.

**Primary author:** PAULUCCI, Laura (UFABC)

**Co-authors:** CAVANNA, Flavio (Fermilab); MARINHO, Franciole; PORTO PAIXAO, Luis Gustavo (UNICAMP)

**Presenter:** PAULUCCI, Laura (UFABC)

**Session Classification:** Poster Session



Contribution ID: 86

Type: **Poster**

## Electrons for Neutrinos

Future long-baseline neutrino oscillation experiments, such as the Deep Underground Neutrino Experiment (DUNE), aim to measure neutrino oscillation parameters with unprecedented precision.

Such sensitivity demands precise characterization of the incoming neutrino energy, which can only be determined via comprehensive cross-section models. In particular DUNE, which flux peaks at 2.5-GeV, will be dominated by pion-production events. Far from ideal, current neutrino pion-production Monte Carlo simulations strongly rely on empirical models which fail to describe neutrino pion-production data at the few-GeV energy range on deuterium and heavier targets. The Electrons for Neutrinos collaboration (e4nu) proposes a novel alternative to leverage neutrino simulations by exploiting the similarities between electrons and neutrino interactions with matter. The e4nu collaboration is working towards a single-differential semi-inclusive pion production measurement with electron-scattering CLAS6 data at 1, 2, 4-GeV on Carbon. The results offer a new insight to exclusive pion-production properties and pion final-state-interactions in the nuclear environment and can be used to validate theory models and neutrino event-generators.

**Primary author:** Dr TENA VIDAL, Julia (Tel Aviv University)

**Presenter:** Dr TENA VIDAL, Julia (Tel Aviv University)

**Session Classification:** Poster Session

Contribution ID: 87

Type: **Poster**

## The Hyper-K Intermediate Water Cherenkov Detector and the Water Cherenkov Test Experiment

The Intermediate Water Cherenkov Detector (IWCD) is a planned component of the Hyper-K long baseline neutrino experiment with vertical mobility capable of scanning a region of the JPARC accelerator-made neutrino beam spectrum. It will contribute to control systematic uncertainties arising from neutrino interaction cross sections and extrapolation of the beam flux between the Hyper-K near and far detectors. A key aspect of IWCD is its high granularity and fast time response thanks to an advanced photo-detection system based on multi-photomultiplier (mPMT) devices developed to deliver excellent Cherenkov ring reconstruction. The Water Cherenkov Test Experiment (WCTE) at CERN is designed to test this system, along with the capabilities of the calibration and reconstruction techniques planned for IWCD. It will use a beam produced at the CERN accelerator to also measure lepton interactions relevant to constraint uncertainties from physics models at a range of energies covering an important region of the IWCD phase space. This poster presents a short summary of both IWCD and WCTE, including their designs, plans and goals.

**Primary author:** BARBI, Mauricio (University of Regina)

**Presenter:** BARBI, Mauricio (University of Regina)

**Session Classification:** Poster Session

Contribution ID: 88

Type: **Poster**

## Measuring muon antineutrino charged-current interactions without mesons in the final state, in the NOvA Near Detector

NOvA is a long-baseline neutrino experiment based at Fermilab in the US, with the primary aim of measuring neutrino and antineutrino oscillations. This will enhance our understanding of electroweak interactions by measuring the neutrino mixing angles, CP-violating phase and neutrino mass ordering. To measure these oscillations, we first need to have a deep understanding of how neutrinos and antineutrinos interact with matter. Antineutrino interaction cross sections are, at present, particularly poorly constrained, and processes such as meson exchange currents are not well understood in the antineutrino sector.

This analysis will develop a cross-section measurement of muon antineutrino interactions without mesons (e.g. pions or kaons) in the final state, in the NOvA near detector. A high-statistics, high-purity sample is obtained through a cut-based selection process implementing machine learning techniques. The sample is dominated by quasi-elastic and meson exchange current interactions which are sensitive to nuclear effects such as final-state interactions. The cross section will be extracted as a function of the incoming neutrino energy and the kinematics of the outgoing particles.

**Primary author:** VOCKERODT, Kevin (Queen Mary University of London)

**Presenter:** VOCKERODT, Kevin (Queen Mary University of London)

**Session Classification:** Poster Session

Contribution ID: 89

Type: **Poster**

## **CEvNS and BSM physics with the CONNIE experiment.**

The CONNIE experiment is dedicated to detecting Coherent Elastic Neutrino-Nucleus Scattering (CEvNS) of reactor antineutrinos using high-resistivity silicon CCDs at the Angra-2 reactor, and underwent significant advancements in 2021. Introducing two Skipper-CCD sensors with sub-electron readout noise, CONNIE became the pioneering project to employ Skipper-CCDs for reactor neutrino detection and achieved a record detection threshold of 15 eV. This work presents an overview of the sensors performance and the latest findings derived from 300 days of data collected during 2021-2022, totaling an exposure of 18.4 g-days. A comparison of event rates during reactor-on and off periods reveals no excess, resulting in upper limits on the neutrino interaction rates at 95% confidence level, consistent with earlier findings. Additionally, searches for Physics Beyond the Standard Model yield constraints on simplified models with light vector mediators. New studies include a dark matter search employing diurnal modulation techniques and a quest for relativistic millicharged particles produced in the reactor. The promising outcomes achieved with a minimal sensor mass underscore the potential of Skipper-CCDs in probing rare neutrino interactions, motivating the future plans to expand the detector mass with a Multi-Chip-Module of Skipper-CCDs.

**Primary author:** VENTURA, Pedro (IF-USP)

**Co-authors:** BONIFAZI, Carla; NASTEVA, Irina (IF-UFRJ)

**Presenter:** VENTURA, Pedro (IF-USP)

**Session Classification:** Poster Session

Contribution ID: **90**

Type: **not specified**

## **Eta production from nucleons and nuclei**

*Thursday, 18 April 2024 08:45 (25 minutes)*

**Presenter:** Dr FATIMA, Atika

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 91

Type: **not specified**

## **Nuclear medium effects in $\nu_\tau$ -A scattering at DUNE energies**

*Thursday, 18 April 2024 09:10 (25 minutes)*

**Presenter:** Dr ZAIDI, Farhana

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 92

Type: **not specified**

## Neutrino-induced single pion production and the reanalyzed bubble chamber data

*Thursday, 18 April 2024 09:35 (25 minutes)*

**Presenter:** MARIANO, Alejandro (Departamento de Física, Universidad Nacional de La plata, Argentina)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 93

Type: **not specified**

## **Photo-, Electro- and Weak Single Pion Production Model**

*Thursday, 18 April 2024 10:00 (25 minutes)*

**Presenter:** KABIRNEZHAD, Minoos (Oxford University)

**Session Classification:** Pion Production/SIS/DIS



Contribution ID: 94

Type: **not specified**

## Pion Production at MINERvA

*Thursday, 18 April 2024 10:55 (20 minutes)*

**Presenters:** HARRIS, Deborah (York University and Fermilab); SULTANA, Mehreen (University of Rochester); SULTANA, Mehreen (University of Georgia)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 95

Type: **not specified**

## **nu\_e and nubar\_e low recoil in MINERvA**

*Thursday, 18 April 2024 11:15 (20 minutes)*

**Presenter:** MCFARLAND, Kevin (University of Rochester)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 96

Type: **not specified**

## **SIS Interactions in MINERvA**

*Thursday, 18 April 2024 11:35 (20 minutes)*

**Presenter:** LOZANO SÁNCHEZ, Adrian

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 97

Type: **not specified**

## **Pion Production in NOvA**

*Thursday, 18 April 2024 11:55 (30 minutes)*

**Presenter:** WETSTEIN, Matt (Iowa State University)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: **98**

Type: **not specified**

## **e4nu**

*Thursday, 18 April 2024 14:00 (30 minutes)*

**Presenters:** TENA VIDAL, Julia (Tel Aviv University); TENA VIDAL, Julia

**Session Classification:** Charged Particle/Testbeam Measurements

Contribution ID: 99

Type: **not specified**

## **Determination of the spectral functions of argon at JLab**

*Thursday, 18 April 2024 14:30 (30 minutes)*

**Presenters:** BENHAR, Omar; BENHAR, Omar (INFN - Rome)

**Session Classification:** Charged Particle/Testbeam Measurements

Contribution ID: **100**

Type: **not specified**

## **Electron Scattering in Generators**

*Thursday, 18 April 2024 15:00 (30 minutes)*

**Presenter:** GARDINER, Steven (Fermilab)

**Session Classification:** Charged Particle/Testbeam Measurements

Contribution ID: **101**

Type: **not specified**

# **Empirical fits to inclusive electron-carbon scattering data obtained by deep-learning methods**

*Thursday, 18 April 2024 16:00 (30 minutes)*

**Presenter:** GRACZYK, Krzysztof

**Session Classification:** Charged Particle/Testbeam Measurements



Contribution ID: **102**

Type: **not specified**

## **Charged particles interaction measurements in ProtoDUNE**

*Thursday, 18 April 2024 16:30 (30 minutes)*

**Presenters:** MARINHO, Franciole (ITA); MARINHO, Franciole

**Session Classification:** Charged Particle/Testbeam Measurements

Contribution ID: **103**

Type: **not specified**

## **ANNIE**

*Thursday, 18 April 2024 17:00 (30 minutes)*

**Presenter:** MASTBAUM, Andrew (Rutgers University)

**Session Classification:** Charged Particle/Testbeam Measurements

Contribution ID: **104**

Type: **not specified**

## **Low-energy: Theory Overview in Inelastic and CEvNS Cross-sections and Impact of BSM Effects**

**Session Classification:** Low-energy Scattering

Contribution ID: 105

Type: **not specified**

## **Low-energy Inelastic Cross-sections: Experiment Review and Future Prospects**

**Session Classification:** Low-energy Scattering

Contribution ID: **106**

Type: **not specified**

## **Neutrino-nucleus coherent scattering: status and prospects**

*Friday, 19 April 2024 09:00 (30 minutes)*

**Presenter:** FIGUEROA-FELICIANO, Enectali (Northwestern University)

**Session Classification:** Low-energy Scattering

Contribution ID: **107**

Type: **not specified**

## Lattice QCD

*Friday, 19 April 2024 10:50 (25 minutes)*

**Presenter:** MEYER, Aaron (Lawrence Livermore National Laboratory)

**Session Classification:** Uncertainties and Prospects for Future Improvements

Contribution ID: **108**

Type: **not specified**

## **NUISANCE and global fits**

*Friday, 19 April 2024 11:35 (25 minutes)*

**Primary author:** WRET, Clarence (Oxford)

**Co-authors:** WILKINSON, Callum (Lawrence Berkeley National Laboratory); STOWELL, Patrick (University of Sheffield)

**Presenter:** WRET, Clarence (Oxford)

**Session Classification:** Uncertainties and Prospects for Future Improvements

Contribution ID: **109**

Type: **not specified**

## **MINERvA form-factor measurement**

*Friday, 19 April 2024 11:15 (20 minutes)*

**Presenters:** OLIVIER, Andrew; OLIVIER, Andrew (University of Notre Dame)

**Session Classification:** Uncertainties and Prospects for Future Improvements



Contribution ID: **110**

Type: **not specified**

## **T2K Near Detector Upgrade**

*Friday, 19 April 2024 12:00 (20 minutes)*

**Presenter:** VIRGINET, Ulysse (LPNHE, Sorbonne Université, IN2P3/CNRS)

**Session Classification:** Uncertainties and Prospects for Future Improvements

Contribution ID: 111

Type: **not specified**

## **Resonance and QE contributions to inclusive nucleon structure functions**

*Friday, 19 April 2024 14:00 (25 minutes)*

**Presenter:** CHRISTY, Eric (Hampton University)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 112

Type: **not specified**

## **The Physics of SIS – TMC and HT**

*Friday, 19 April 2024 14:25 (25 minutes)*

**Presenter:** MORFIN, Jorge (Fermilab)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 113

Type: **not specified**

## **Superscaling in the inelastic region: The SuSAv2-inelastic model**

*Friday, 19 April 2024 14:50 (25 minutes)*

**Presenter:** GONZÁLEZ ROSA, Jesús (University of Seville)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: 114

Type: **not specified**

## Neutrino Cross Sections for collider neutrinos

*Friday, 19 April 2024 15:45 (25 minutes)*

**Presenters:** JEONG, Yu Seon (CERN); JEONG, Yu Seon (Chung-Ang University)

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: **115**

Type: **not specified**

## **MicroBooNE**

*Friday, 19 April 2024 16:10 (25 minutes)*

**Presenter:** BOGART, Benjamin

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: **116**

Type: **not specified**

## **T2K latest results and prospects on NC and CC pion production**

*Friday, 19 April 2024 16:35 (30 minutes)*

**Presenter:** JESUS VALLS, Cesar

**Session Classification:** Pion Production/SIS/DIS

Contribution ID: **117**

Type: **not specified**

## **DUNE 2x2**

*Friday, 19 April 2024 17:05 (20 minutes)*

**Presenters:** CHEN, Yifan; CHEN, Yifan

**Session Classification:** Pion Production/SIS/DIS



Contribution ID: **118**

Type: **not specified**

## **Impact of Cross Section Uncertainties on BSM Searches**

*Friday, 19 April 2024 09:30 (35 minutes)*

**Presenters:** KOPP, Joachim; KOPP, Joachim

**Session Classification:** Impact of scattering uncertainties on measurements

Contribution ID: **119**

Type: **not specified**

## Experimental Summary

*Saturday, 20 April 2024 08:55 (30 minutes)*

**Presenter:** PALEY, Jonathan (Fermilab)

**Session Classification:** Closeout

Contribution ID: **120**

Type: **not specified**

## Theory Summary

*Saturday, 20 April 2024 09:25 (30 minutes)*

**Presenter:** ANKOWSKI, Artur (UNIVERSITY OF WROCLAW)

**Session Classification:** Closeout

Contribution ID: **121**

Type: **not specified**

## Closing Remarks

*Saturday, 20 April 2024 11:25 (10 minutes)*

**Presenter:** JACHOWICZ, Natalie (Ghent University)

**Session Classification:** Closeout

Contribution ID: **122**

Type: **not specified**

## **MINERvA**

*Tuesday, 16 April 2024 14:00 (30 minutes)*

**Presenter:** RUTERBORIES, Daniel (University of Rochester)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **123**

Type: **not specified**

## **NOvA**

*Tuesday, 16 April 2024 14:30 (30 minutes)*

**Presenters:** SANCHEZ FALERO, Sebastian (Iowa State University); SANCHEZ FALERO, Sebastian (Iowa State University)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **124**

Type: **not specified**

## **T2K**

*Tuesday, 16 April 2024 15:00 (30 minutes)*

**Presenter:** DOLAN, Stephen (CERN)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: 125

Type: **Poster**

## Double Beta Decay within the Relativistic QRPA

In nature there are about 50 nuclear systems where the single beta-decay is energetically forbidden, and double beta-decay turns out to be only possible mode of disintegration. It is the nuclear pairing force which causes such an “anomaly”, by making the mass of the odd-odd isobar,  $(N-1, Z+1)$ , to be greater than the masses of its even-even neighbors,  $(N, Z)$  and  $(N-2, Z-2)$ . The modes by which the double beta decay can take place are connected with the neutrino and antineutrino distinction. The Quasi-Particle Random Phase Approximation (QRPA) has turned out to be the most simple model for calculating the nuclear wave function involved in the single and double beta-decay transitions. In this work we perform a self-consistent relativistic QRPA (RQRPA) calculation of double beta-decay based on relativistic BCS (RBCS) mean field theory results for odd-odd intermediate nuclei  $^{48}\text{Sc}$ ,  $^{76}\text{As}$ ,  $^{82}\text{Br}$ ,  $^{100}\text{Tc}$ ,  $^{128}\text{I}$ , and  $^{130}\text{I}$ . We use the parameter set NL3 for interactions between protons, neutrons,  $\sigma$ ,  $\omega$ ,  $\rho$  mesons and photons. The RQRPA equations are solved for the residual pion and rho interaction by employing the same parameters used in the RBCS, and experimental values for the pion and nucleons. The RQRPA results for the double beta-decay matrix elements are similar to those obtained within the QRPA and the shell model. Motivated by the results obtained here, we intend to calculate the neutrinoless decay in order to estimate the neutrino mass.

### References

- F. Krmpotic, FIZIKA B 14 (2005) 2, 139.
- N. Paar, D. Vretenar, T. Marketin and P. Ring, Phys. Rev. C 77 (2008) 024608.
- B. D. Serot and J. D. Walecka, Adv. Nucl. Phys. 16 (1986) 1.
- C. De Conti, A. P. Galeão and F. Krmpotic', Phys. Lett. B 444 (1999) 14.
- C. De Conti, A. P. Galeão and F. Krmpotic', Phys. Lett. B 494 (2000) 46.
- D. Vretenar, N. Paar, P. Ring, and G. A. Lalazissis, Phys. Rev. C 63 (2001) 047301.
- B. V. Carlson and D. Hirata, Phys. Rev. C 62 (2000) 054310.

**Primary authors:** CARVALHO DOS SANTOS MAIA, Lucas (Universidade Estadual de Santa Cruz); DE CONTI, Cláudio (Universidade Estadual Paulista)

**Co-author:** SAMANA, Arturo Rodolfo (Universidade Estadual de Santa Cruz)

**Presenters:** CARVALHO DOS SANTOS MAIA, Lucas (Universidade Estadual de Santa Cruz); DE CONTI, Cláudio (Universidade Estadual Paulista); SAMANA, Arturo Rodolfo (Universidade Estadual de Santa Cruz)

**Session Classification:** Poster Session



Contribution ID: **126**

Type: **not specified**

## Welcome

*Thursday, 11 April 2024 09:00 (10 minutes)*

**Presenter:** DA MOTTA, Hélio (CBPF)

**Session Classification:** Theoretical introduction

Contribution ID: 127

Type: **not specified**

## General introduction

*Thursday, 11 April 2024 09:10 (30 minutes)*

**Presenter:** HARRIS, Deborah (York University and Fermilab)

**Session Classification:** Theoretical introduction

Contribution ID: 128

Type: **not specified**

## General theory at the nucleon level

*Thursday, 11 April 2024 10:05 (45 minutes)*

**Presenter:** NIKOLAKOPOULOS, Alexis (FNAL)

**Session Classification:** Theoretical introduction

Contribution ID: **129**

Type: **not specified**

## **Introduction to nuclear effects**

*Thursday, 11 April 2024 11:05 (1 hour)*

**Presenter:** Dr ANKOWSKI, Artur (University of Wroclaw)

**Session Classification:** Theoretical introduction

Contribution ID: 130

Type: **not specified**

## Relativistic mean field approach

*Thursday, 11 April 2024 13:45 (45 minutes)*

**Presenter:** NIKOLAKOPOULOS, Alexis (FNAL)

**Session Classification:** Theoretical introduction

Contribution ID: 131

Type: **not specified**

## Spectral function approach

*Thursday, 11 April 2024 14:45 (45 minutes)*

**Presenter:** Dr ANKOWSKI, Artur (University of Wrocław)

**Session Classification:** Theoretical introduction

Contribution ID: 132

Type: **not specified**

## **Ab initio approach**

*Thursday, 11 April 2024 16:00 (45 minutes)*

**Presenter:** ANDREOLI, Lorenzo

**Session Classification:** Theoretical introduction

Contribution ID: 133

Type: **not specified**

## How to make a cross-section measurement

*Saturday, 13 April 2024 09:00 (45 minutes)*

**Presenter:** CHERDACK, Daniel (University of Houston)

**Session Classification:** Experimental introduction



Contribution ID: 134

Type: **not specified**

## Generator comparisons to data

*Saturday, 13 April 2024 10:15 (1 hour)*

**Presenters:** WILKINSON, Callum (Lawrence Berkeley National Laboratory); WRET, Clarence (Oxford)

**Session Classification:** Experimental introduction

Contribution ID: 135

Type: **not specified**

## **Hands-on exercises with generators**

*Saturday, 13 April 2024 11:30 (1 hour)*

**Presenters:** WILKINSON, Callum (Lawrence Berkeley National Laboratory); WRET, Clarence (Oxford)

**Session Classification:** Experimental introduction

Contribution ID: 136

Type: **not specified**

## How generators work

*Friday, 12 April 2024 09:00 (45 minutes)*

**Presenter:** NIEWCZAS, Kajetan (Ghent University)

**Session Classification:** Introduction to Monte Carlo generators

Contribution ID: 137

Type: **not specified**

## **Practicum: analyzing generator output**

*Friday, 12 April 2024 14:00 (1h 30m)*

**Primary author:** WRET, Clarence (Oxford)

**Co-author:** WILKINSON, Callum (Lawrence Berkeley National Laboratory)

**Presenter:** WRET, Clarence (Oxford)

**Session Classification:** Introduction to Monte Carlo generators

Contribution ID: **138**

Type: **not specified**

## **Generators review**

*Friday, 12 April 2024 10:00 (1h 45m)*

**Presenter:** GARDINER, Steven (Fermilab)

**Session Classification:** Introduction to Monte Carlo generators

Contribution ID: 139

Type: **not specified**

## Practicum continued

*Friday, 12 April 2024 15:45 (1h 15m)*

**Presenters:** WILKINSON, Callum (Lawrence Berkeley National Laboratory); WRET, Clarence (Oxford)

**Session Classification:** Introduction to Monte Carlo generators

Contribution ID: 140

Type: **not specified**

## **Meson exchange currents in a relativistic mean field model**

*Tuesday, 16 April 2024 16:50 (25 minutes)*

**Presenter:** FRANCO MUÑOZ, Tania (Universidad Complutense de Madrid)

**Session Classification:** 0pi / QE / 2p2h

Contribution ID: **141**

Type: **not specified**

## **Introduction to simulation**

*Thursday, 11 April 2024 12:05 (5 minutes)*

**Presenter:** WILKINSON, Callum (Lawrence Berkeley National Laboratory)

**Session Classification:** Theoretical introduction



Contribution ID: 142

Type: **not specified**

## **Practicum: running a generator**

*Friday, 12 April 2024 11:45 (45 minutes)*

**Presenters:** WILKINSON, Callum (Lawrence Berkeley National Laboratory); WRET, Clarence (Oxford)

**Session Classification:** Introduction to Monte Carlo generators

Contribution ID: **143**

Type: **not specified**

## Welcome

*Monday, 15 April 2024 08:45 (10 minutes)*

**Presenters:** PERES, Orlando Luis Goulart (UNICAMP); DA MOTTA FILHO, Helio (CBPF)

**Session Classification:** Opening

Contribution ID: **144**

Type: **not specified**

## **ICTP-SAIFR**

*Monday, 15 April 2024 08:55 (10 minutes)*

**Presenter:** BERKOVITS, Nathan (ICTP-SAIFR)

**Session Classification:** Opening

Contribution ID: 145

Type: **not specified**

## Institute Principia

*Monday, 15 April 2024 09:05 (5 minutes)*

**Presenter:** FRANCISCO, Gerson (Instituto Principia)

**Session Classification:** Opening

Contribution ID: **146**

Type: **not specified**

## **Registration**

*Monday, 15 April 2024 08:25 (20 minutes)*

**Session Classification:** Registration

Contribution ID: **147**

Type: **not specified**

## Information

*Monday, 15 April 2024 09:10 (5 minutes)*

**Presenters:** DA MOTTA FILHO, Helio (CBPF); PERES, Orlando Luis Goulart (UNICAMP)

**Session Classification:** Opening

Contribution ID: 148

Type: **not specified**

## **Nitrogen removal from liquid Argon using Li-FAU, an innovative Brazilian method**

*Friday, 19 April 2024 10:05 (5 minutes)*

**Presenter:** SOUZA CORREIA, Daniel

**Session Classification:** Poster Winners

Contribution ID: **149**

Type: **not specified**

## **Ant colony in Sensitivity Studies**

*Friday, 19 April 2024 10:10 (5 minutes)*

**Presenter:** WIELER, Felipe

**Session Classification:** Poster Winners



Contribution ID: 150

Type: **not specified**

## **CEvNS and BSM physics with the CONNIE experiment.**

*Friday, 19 April 2024 10:15 (5 minutes)*

**Presenter:** VENTURA, Pedro (IF-USP)

**Session Classification:** Poster Winners

Contribution ID: **151**

Type: **not specified**

## Discussion

*Saturday, 20 April 2024 10:35 (45 minutes)*

**Presenter:** ALL !

**Session Classification:** Discussion

Contribution ID: 152

Type: **not specified**

## Closing remarks

**Presenter:** DA MOTTA, Helio

**Session Classification:** Closeout