Contribution ID: 51

Type: not specified

An axion-like particle explanation of $B \to \pi K$ puzzle and $B \to K \nu \bar{\nu}$ excess

Tuesday, 4 June 2024 11:00 (30 minutes)

In light of the recent branching fraction measurement of $B^+ \to K^+ \nu \bar{\nu}$ -decay and its deviation from the SM expectation, we analyze the prospect of an axion-like particle (ALP) as the cause of such a departure. We assume a long-lived ALP with a mass of the order of a pion that predominantly decays to two photons. We assess the scenario where the ALP decay length is several meters and therefore has a non-negligible probability to decay outside the detector volume of Belle-II mimicking the $B^+ \to K^+ \nu \bar{\nu}$ -signal. Remarkably, such an arrangement provides a simple explanation to the long-standing $B \to \pi K$ -puzzle by noting that the measured $B^0 \to \pi^0 K^0$ and $B^+ \to \pi^0 K^+$ decays have a $B^0 \to a K^0$ and $B^+ \to a K^+$ component respectively. We also argue based on our results that the axion-photon effective coupling belongs to a region in the parameter space that can be probed in future experiments.

Primary authors: ROY, Shibasis (Chennai Mathematical Institute); Prof. ALTMANNSHOFER, Wolfgang (UC Santa Cruz)

Presenter: ROY, Shibasis (Chennai Mathematical Institute)

Session Classification: Session 3