Update on 3D reconstruction

2024-02-15








Reminder: Last status

- Looked into different issues in 2D reconstruction
- Fixed issues with 'missing' hits at end/start of tracks → extrapolation







Towards 3D matching

Initial idea for matching

- Match ends of tracks if close enough (space + time)
- Put all hits of both tracks into one new
- Optional
 - Add adjacent hits to new track?
 - Combined 3D line from Hough transform or calculate from combined lines?
 - Kalman filter?





Towards 3D matching: Refine plan

Idea for matching

- Match ends of tracks if close enough (space + time)
- Put all hits of both tracks into one new
- Optional
 - Add adjacent hits to new track?

 Not for now
 - Combined 3D line from Hough transform or calculate from combined lines?
 - Kalman filter?





Towards 3D matching: Refine plan

Idea for matching


- Match ends of tracks if close enough (space + time)
 - Close enough at start and end: ±1 plane number(!) and within 10 bars
 - Close enough in time: in same time slice (and within 30 ns, problem with hit time assignment at the moment → Jeffrey?)
- Put all hits of both tracks into one new
 - Copy hits from both tracks into **new 3D track class** (→ Liam)
 - Can contain: hit positions, start hit, end hit, direction of track, length of track, energy deposit, energy range, time
 - Calculate y coordinate per plane number pair (if possible, handle exemptions)





How to calculate y coordinate?

- Layers are tilted against each other by 3°
 - Use x position of the hit scintillator bars → calculate geometry y position
 - y ='Anchor point' in $y \frac{1}{2} tan(90^{\circ}-3^{\circ}) | x_bar1 x_bar2 | (fiber crossing)$
- The true hit can be anywhere in the crossing area though
 - From fiber crossing to top of bar crossing: ~34 cm → uncertainty










How to display this now?

- Event display so far only takes x and z coordinates
- Need to develop new event display for 3D tracks













2D projections of the 3D tracks

- Display **hits** in the **size** that they actually are
 - add in air gaps between steel and scintillator for visible hits
- Use uncertainty of hit position in y for hit areas
- Different views in scale







More examples





Need to **check** that everything is **correct** now

Then make **pull request** to merge nehm_3D_reconstruction branch onto main













Backup

