

# Investigating the future of proton decay searches using paleo detectors

*Tuesday, 17 September 2024 16:55 (20 minutes)*

Paleo-detectors utilize the fact that mineral lattices can retain deformations in their structure caused by charged particle interactions. We consider the search for proton decay,  $p \rightarrow \bar{\nu} K^+$ , in such detectors via the possible crystal damage produced by the endpoint of the charged kaon. Atmospheric neutrino induced backgrounds render this search impossible on Earth, but in a lunar environment with no atmosphere, this background is significantly suppressed. For a 100g,  $10^9$  year-old (100kton-yr exposure) piece of lunar olivine, we expect 0.5 kaon endpoints due to atmospheric neutrino backgrounds. If this lunar sample could be acquired and analyzed, the proton decay sensitivity would be  $\tau_p \approx 10^{34}$  years, which is competitive with Super-Kamiokande's currently published limit of  $\tau_p > 5.9 \times 10^{33}$  years at 90% CL, as well as the projected range of Hyper-Kamiokande and DUNE for the  $p \rightarrow \bar{\nu} K^+$  channel.

## Working Group

WG 5: Neutrinos Beyond PMNS

**Primary author:** LITTLE, Cassandra (University of Michigan)

**Co-authors:** SPITZ, Joshua (University of Michigan); Dr BAUM, Sebastian (Institute for Theoretical Particle Physics and Cosmology); Dr STENGEL, Patrick (INFN Ferrara)

**Presenter:** LITTLE, Cassandra (University of Michigan)

**Session Classification:** Parallel: WG5

**Track Classification:** WG5: Neutrino Beyond PMNS