

BSM Theory

CERN/Fermilab HCPSS

Adam Martin
amarti41@nd.edu



ATLAS Exotics Searches* - 95% CL Upper Exclusion Limits

Status: May 2019

ATLAS Preliminary

$\int \mathcal{L} dt = (3.2 - 139) \text{ fb}^{-1}$


$\sqrt{s} = 8, 13 \text{ TeV}$

	Model	ℓ, γ	Jets†	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Limit	Reference
Extra dimensions	ADD $G_{KK} + g/q$	$0 e, \mu$	1-4 j	Yes	36.1	M_D 7.7 TeV	$n = 2$ 1711.03301
	ADD non-resonant $\gamma\gamma$	2γ	-	-	36.7	M_S 8.6 TeV	$n = 3$ HLZ NLO 1707.04147
	ADD QBH	-	2 j	-	37.0	M_{th} 8.9 TeV	$n = 6$ 1703.09127
	ADD BH high $\sum p_T$	$\geq 1 e, \mu$	$\geq 2 j$	-	3.2	M_{th} 8.2 TeV	$n = 6, M_D = 3 \text{ TeV}$, rot BH 1606.02265
	ADD BH multijet	-	$\geq 3 j$	-	3.6	M_{th} 9.55 TeV	$n = 6, M_D = 3 \text{ TeV}$, rot BH 1512.02586
	RS1 $G_{KK} \rightarrow \gamma\gamma$	2γ	-	-	36.7	G_{KK} mass 4.1 TeV	$k/\overline{M}_{Pl} = 0.1$ 1707.04147
	Bulk RS $G_{KK} \rightarrow WW/ZZ$	multi-channel	-	-	36.1	G_{KK} mass 2.3 TeV	$k/\overline{M}_{Pl} = 1.0$ 1808.02380
	Bulk RS $G_{KK} \rightarrow WW \rightarrow qqqq$	$0 e, \mu$	2 J	-	139	G_{KK} mass 1.6 TeV	$k/\overline{M}_{Pl} = 1.0$ ATLAS-CONF-2019-003
	Bulk RS $g_{KK} \rightarrow tt$	$1 e, \mu$	$\geq 1 b, \geq 1J/2j$	Yes	36.1	g_{KK} mass 3.8 TeV	$\Gamma/m = 15\%$ 1804.10823
	2UED / RPP	$1 e, \mu$	$\geq 2 b, \geq 3 j$	Yes	36.1	KK mass 1.8 TeV	Tier (1,1), $\mathcal{B}(A^{(1,1)} \rightarrow tt) = 1$ 1803.09678
Gauge bosons	SSM $Z' \rightarrow \ell\ell$	$2 e, \mu$	-	-	139	Z' mass 5.1 TeV	$\Gamma/m = 1\%$ CERN-EP-2019-100 ATLAS-CONF-2019-003
	SSM $Z' \rightarrow \tau\tau$	2τ	-	-	36.1	Z' mass 2.42 TeV	
	Leptophobic $Z' \rightarrow bb$	-	2 b	-	36.1	Z' mass 2.1 TeV	
	Leptophobic $Z' \rightarrow tt$	$1 e, \mu$	$\geq 1 b, \geq 1J/2j$	Yes	36.1	Z' mass 3.0 TeV	
	SSM $W' \rightarrow \ell\nu$	$1 e, \mu$	-	Yes	139	W' mass 6.0 TeV	
	SSM $W' \rightarrow \tau\nu$	1τ	-	Yes	36.1	W' mass 3.7 TeV	
	HVT $V' \rightarrow WZ \rightarrow qqqq$ model B	$0 e, \mu$	2 J	-	139	V' mass 3.6 TeV	
	HVT $V' \rightarrow WH/ZH$ model B	multi-channel	-	-	36.1	V' mass 2.93 TeV	
	LRSM $W_R \rightarrow tb$	multi-channel	-	-	36.1	W_R mass 3.25 TeV	
	LRSM $W_R \rightarrow \mu N_R$	2μ	1 J	-	80	W_R mass 5.0 TeV	
CI	CI $qqqq$	-	2 j	-	37.0	Λ 21.8 TeV η_{LL}^-	1703.09127
	CI $\ell\ell qq$	$2 e, \mu$	-	-	36.1	Λ 40.0 TeV η_{LL}^-	1707.02424
	CI $tttt$	$\geq 1 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	Λ 2.57 TeV $ C_{4t} = 4\pi$	1811.02305
DM	Axial-vector mediator (Dirac DM)	$0 e, \mu$	1-4 j	Yes	36.1	m_{med} 1.55 TeV	$g_q=0.25, g_\chi=1.0, m(\chi) = 1 \text{ GeV}$ 1711.03301
	Colored scalar mediator (Dirac DM)	$0 e, \mu$	1-4 j	Yes	36.1	m_{med} 1.67 TeV	$g=1.0, m(\chi) = 1 \text{ GeV}$ 1711.03301
	$VV\chi\chi$ EFT (Dirac DM)	$0 e, \mu$	1 J, $\leq 1 j$	Yes	3.2	M_* 700 GeV	$m(\chi) < 150 \text{ GeV}$ 1608.02372
	Scalar reson. $\phi \rightarrow t\chi$ (Dirac DM)	$0-1 e, \mu$	1 b, 0-1 J	Yes	36.1	m_ϕ 3.4 TeV	$y = 0.4, \lambda = 0.2, m(\chi) = 10 \text{ GeV}$ 1812.09743
LQ	Scalar LQ 1 st gen	$1, 2 e$	$\geq 2 j$	Yes	36.1	LQ mass 1.4 TeV	$\beta = 1$ 1902.00377
	Scalar LQ 2 nd gen	$1, 2 \mu$	$\geq 2 j$	Yes	36.1	LQ mass 1.56 TeV	$\beta = 1$ 1902.00377
	Scalar LQ 3 rd gen	2τ	2 b	-	36.1	LQ_3^q mass 1.03 TeV	$\mathcal{B}(LQ_3^q \rightarrow b\tau) = 1$ 1902.08103
	Scalar LQ 3 rd gen	$0-1 e, \mu$	2 b	Yes	36.1	LQ_3^d mass 970 GeV	$\mathcal{B}(LQ_3^d \rightarrow t\tau) = 0$ 1902.08103
Heavy quarks	VLQ $TT \rightarrow Ht/Zt/Wb + X$	multi-channel	-	-	36.1	T mass 1.37 TeV	SU(2) doublet 1808.02343
	VLQ $BB \rightarrow Wt/Zb + X$	multi-channel	-	-	36.1	B mass 1.34 TeV	SU(2) doublet 1808.02343
	VLQ $T_{5/3} T_{5/3} T_{5/3} \rightarrow Wt + X$	$2(SS) \geq 3 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	$T_{5/3}$ mass 1.64 TeV	$\mathcal{B}(T_{5/3} \rightarrow Wt) = 1, c(T_{5/3} Wt) = 1$ 1807.11883
	VLQ $Y \rightarrow Wb + X$	$1 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	Y mass 1.85 TeV	$\mathcal{B}(Y \rightarrow Wb) = 1, c_R(Wb) = 1$ 1812.07343
	VLQ $B \rightarrow Hb + X$	$0 e, \mu, 2 \gamma$	$\geq 1 b, \geq 1 j$	Yes	79.8	B mass 1.21 TeV	$\kappa_B = 0.5$ ATLAS-CONF-2018-024
VLQ $QQ \rightarrow WqWq$	$1 e, \mu$	$\geq 4 j$	Yes	20.3	Q mass 690 GeV	1509.04261	
Excited fermions	Excited quark $q^* \rightarrow qg$	-	2 j	-	139	q^* mass 6.7 TeV	only u^* and d^* , $\Lambda = m(q^*)$ ATLAS-CONF-2019-007
	Excited quark $q^* \rightarrow q\gamma$	1γ	1 j	-	36.7	q^* mass 5.3 TeV	only u^* and d^* , $\Lambda = m(q^*)$ 1709.10440
	Excited quark $b^* \rightarrow bg$	-	1 b, 1 j	-	36.1	b^* mass 2.6 TeV	1805.09299
	Excited lepton l^*	$3 e, \mu$	-	-	20.3	l^* mass 3.0 TeV	$\Lambda = 3.0 \text{ TeV}$ 1411.2921
	Excited lepton ν^*	$3 e, \mu, \tau$	-	-	20.3	ν^* mass 1.6 TeV	$\Lambda = 1.6 \text{ TeV}$ 1411.2921
Other	Type III Seesaw	$1 e, \mu$	$\geq 2 j$	Yes	79.8	N^0 mass 560 GeV	ATLAS-CONF-2018-020
	LRSM Majorana ν	2μ	2 j	-	36.1	N_R mass 3.2 TeV	
	Higgs triplet $H^{\pm\pm} \rightarrow \ell\ell$	$2, 3, 4 e, \mu$ (SS)	-	-	36.1	$H^{\pm\pm}$ mass 870 GeV	
	Higgs triplet $H^{\pm\pm} \rightarrow \ell\tau$	$3 e, \mu, \tau$	-	-	20.3	$H^{\pm\pm}$ mass 400 GeV	
	Multi-charged particles	-	-	-	36.1	multi-charged particle mass 1.22 TeV	
	Magnetic monopoles	-	-	-	34.4	monopole mass 2.37 TeV	

$\sqrt{s} = 8 \text{ TeV}$ $\sqrt{s} = 13 \text{ TeV}$ partial data $\sqrt{s} = 13 \text{ TeV}$ full data

10⁻¹ 1 10 Mass scale [TeV]

Inclusive Searches	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$	0	2-6 jets	Yes	36.1	\tilde{q}	1.57 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(1^{\text{st}} \text{ gen. } \tilde{q}) = m(2^{\text{nd}} \text{ gen. } \tilde{q})$	1712.02332	
	$\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ (compressed)	mono-jet	1-3 jets	Yes	36.1	\tilde{q}	710 GeV	$m(\tilde{q}) - m(\tilde{\chi}_1^0) < 5 \text{ GeV}$	1711.03301	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	36.1	\tilde{g}	2.02 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}$	1712.02332	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0 \rightarrow q\tilde{q}W^\pm\tilde{\chi}_1^0$	0	2-6 jets	Yes	36.1	\tilde{g}	2.01 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}^\pm) = 0.5(m(\tilde{\chi}_1^0) + m(\tilde{g}))$	1712.02332	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}(\ell\ell)\tilde{\chi}_1^0$	$ee, \mu\mu$	2 jets	Yes	14.7	\tilde{g}	1.7 TeV	$m(\tilde{\chi}_1^0) < 300 \text{ GeV},$	1611.05791	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}(\ell\ell/\nu\nu)\tilde{\chi}_1^0$	3 e, μ	4 jets	-	36.1	\tilde{g}	1.87 TeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$	1706.03731	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}WZ\tilde{\chi}_1^0$	0	7-11 jets	Yes	36.1	\tilde{g}	1.8 TeV	$m(\tilde{\chi}_1^0) < 400 \text{ GeV}$	1708.02794	
	GMSB ($\tilde{\ell}$ NLSP)	1-2 $\tau + 0-1 \ell$	0-2 jets	Yes	3.2	\tilde{g}	2.0 TeV		1607.05979	
	GGM (bino NLSP)	2 γ	-	Yes	36.1	\tilde{g}	2.15 TeV	$c\tau(\text{NLSP}) < 0.1 \text{ mm}$	ATLAS-CONF-2017-080	
	GGM (higgsino-bino NLSP)	γ	2 jets	Yes	36.1	\tilde{g}	2.05 TeV	$m(\tilde{\chi}_1^0) = 1700 \text{ GeV}, c\tau(\text{NLSP}) < 0.1 \text{ mm}, \mu > 0$	ATLAS-CONF-2017-080	
	Gravitino LSP	0	mono-jet	Yes	20.3	\tilde{g}	865 GeV	$m(\tilde{G}) > 1.8 \times 10^{-4} \text{ eV}, m(\tilde{g}) = m(\tilde{q}) = 1.5 \text{ TeV}$	1502.01518	
	3 rd gen. \tilde{g} med.	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow b\tilde{b}\tilde{\chi}_1^0$	0	3 b	Yes	36.1	\tilde{g}	1.92 TeV	$m(\tilde{\chi}_1^0) < 600 \text{ GeV}$	1711.01901
		$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0$	0-1 e, μ	3 b	Yes	36.1	\tilde{g}	1.97 TeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}$	1711.01901
	3 rd gen. squarks direct production	$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$	0	2 b	Yes	36.1	\tilde{b}_1	950 GeV	$m(\tilde{\chi}_1^0) < 420 \text{ GeV}$	1708.09266
$\tilde{b}_1\tilde{b}_1, \tilde{b}_1 \rightarrow t\tilde{\chi}_1^+$		2 e, μ (SS)	1 b	Yes	36.1	\tilde{b}_1	275-700 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}, m(\tilde{\chi}_1^\pm) = m(\tilde{\chi}_1^0) + 100 \text{ GeV}$	1706.03731	
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$		0-2 e, μ	1-2 b	Yes	4.7/13.3	\tilde{t}_1	117-170 GeV, 200-720 GeV	$m(\tilde{\chi}_1^\pm) = 2m(\tilde{\chi}_1^0), m(\tilde{\chi}_1^0) = 55 \text{ GeV}$	1209.2102, ATLAS-CONF-2016-077	
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow Wb\tilde{\chi}_1^0$ or $t\tilde{\chi}_1^0$		0-2 e, μ	0-2 jets/1-2 b	Yes	20.3/36.1	\tilde{t}_1	90-198 GeV, 0.195-1.0 TeV	$m(\tilde{\chi}_1^0) = 1 \text{ GeV}$	1506.08616, 1709.04183, 1711.11520	
$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow c\tilde{\chi}_1^0$		0	mono-jet	Yes	36.1	\tilde{t}_1	90-430 GeV	$m(\tilde{t}_1) - m(\tilde{\chi}_1^0) = 5 \text{ GeV}$	1711.03301	
$\tilde{t}_1\tilde{t}_1$ (natural GMSB)		2 e, μ (Z)	1 b	Yes	20.3	\tilde{t}_1	150-600 GeV	$m(\tilde{\chi}_1^0) > 150 \text{ GeV}$	1403.5222	
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + Z$		3 e, μ (Z)	1 b	Yes	36.1	\tilde{t}_2	290-790 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$	1706.03986	
$\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + h$		1-2 e, μ	4 b	Yes	36.1	\tilde{t}_2	320-880 GeV	$m(\tilde{\chi}_1^0) = 0 \text{ GeV}$	1706.03986	
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	36.1	$\tilde{\ell}$	90-500 GeV	$m(\tilde{\chi}_1^0) = 0$	ATLAS-CONF-2017-039	
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow \tilde{\ell}\nu(\ell\bar{\nu})$	2 e, μ	0	Yes	36.1	$\tilde{\chi}_1^\pm$	750 GeV	$m(\tilde{\chi}_1^0) = 0, m(\tilde{\ell}, \bar{\nu}) = 0.5(m(\tilde{\chi}_1^\pm) + m(\tilde{\chi}_1^0))$	ATLAS-CONF-2017-039	
	$\tilde{\chi}_1^+\tilde{\chi}_1^+/\tilde{\chi}_1^+\tilde{\chi}_1^0, \tilde{\chi}_1^+ \rightarrow \tilde{\tau}\nu(\tau\bar{\nu}), \tilde{\chi}_2^0 \rightarrow \tilde{\tau}\tau(\nu\bar{\nu})$	2 τ	-	Yes	36.1	$\tilde{\chi}_1^\pm$	760 GeV	$m(\tilde{\chi}_1^0) = 0, m(\tilde{\tau}, \bar{\nu}) = 0.5(m(\tilde{\chi}_1^\pm) + m(\tilde{\chi}_1^0))$	1708.07875	
	$\tilde{\chi}_1^+\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\nu\tilde{\ell}\nu(\ell\bar{\nu}\nu), \ell\tilde{\nu}\tilde{\ell}\nu(\ell\bar{\nu}\nu)$	3 e, μ	0	Yes	36.1	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$	1.13 TeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 0, m(\tilde{\ell}, \bar{\nu}) = 0.5(m(\tilde{\chi}_1^\pm) + m(\tilde{\chi}_1^0))$	ATLAS-CONF-2017-039	
	$\tilde{\chi}_1^+\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$	2-3 e, μ	0-2 jets	Yes	36.1	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$	580 GeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 0, \tilde{\ell} \text{ decoupled}$	ATLAS-CONF-2017-039	
	$\tilde{\chi}_1^+\tilde{\chi}_2^0 \rightarrow W\tilde{\chi}_1^0 h\tilde{\chi}_1^0, h \rightarrow b\bar{b}/WW/\tau\tau/\gamma\gamma$	e, μ, γ	0-2 b	Yes	20.3	$\tilde{\chi}_1^\pm, \tilde{\chi}_2^0$	270 GeV	$m(\tilde{\chi}_1^\pm) = m(\tilde{\chi}_2^0), m(\tilde{\chi}_1^0) = 0, \tilde{\ell} \text{ decoupled}$	1501.07110	
	$\tilde{\chi}_2^0\tilde{\chi}_3^0, \tilde{\chi}_2^0 \rightarrow \tilde{\ell}_R\ell$	4 e, μ	0	Yes	20.3	$\tilde{\chi}_2^0$	635 GeV	$m(\tilde{\chi}_2^0) = m(\tilde{\chi}_3^0), m(\tilde{\chi}_1^0) = 0, m(\tilde{\ell}, \bar{\nu}) = 0.5(m(\tilde{\chi}_2^0) + m(\tilde{\chi}_1^0))$	1405.5086	
	GGM (wino NLSP) weak prod., $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$	1 $e, \mu + \gamma$	-	Yes	20.3	\tilde{W}	115-370 GeV	$c\tau < 1 \text{ mm}$	1507.05493	
	GGM (bino NLSP) weak prod., $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$	2 γ	-	Yes	36.1	\tilde{W}	1.06 TeV	$c\tau < 1 \text{ mm}$	ATLAS-CONF-2017-080	
	Long-lived particles	Direct $\tilde{\chi}_1^+\tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^\pm$	Disapp. trk	1 jet	Yes	36.1	$\tilde{\chi}_1^\pm$	460 GeV	$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) \sim 160 \text{ MeV}, \tau(\tilde{\chi}_1^\pm) = 0.2 \text{ ns}$	1712.02118
Direct $\tilde{\chi}_1^+\tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^\pm$		dE/dx trk	-	Yes	18.4	$\tilde{\chi}_1^\pm$	495 GeV	$m(\tilde{\chi}_1^\pm) - m(\tilde{\chi}_1^0) \sim 160 \text{ MeV}, \tau(\tilde{\chi}_1^\pm) < 15 \text{ ns}$	1506.05332	
Stable, stopped \tilde{g} R-hadron		0	1-5 jets	Yes	27.9	\tilde{g}	850 GeV	$m(\tilde{\chi}_1^0) = 100 \text{ GeV}, 10 \mu\text{s} < \tau(\tilde{g}) < 1000 \text{ s}$	1310.6584	
Stable \tilde{g} R-hadron		trk	-	-	3.2	\tilde{g}	1.58 TeV		1606.05129	
Metastable \tilde{g} R-hadron		dE/dx trk	-	-	3.2	\tilde{g}	1.57 TeV	$m(\tilde{\chi}_1^0) = 100 \text{ GeV}, \tau > 10 \text{ ns}$	1604.04520	
Metastable \tilde{g} R-hadron, $\tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0$		displ. vtx	-	Yes	32.8	\tilde{g}	2.37 TeV	$\tau(\tilde{g}) = 0.17 \text{ ns}, m(\tilde{\chi}_1^0) = 100 \text{ GeV}$	1710.04901	
GMSB, stable $\tilde{\tau}, \tilde{\chi}_1^0 \rightarrow \tilde{\tau}(\bar{e}, \bar{\mu}) + \tau(e, \mu)$		1-2 μ	-	-	19.1	$\tilde{\chi}_1^0$	537 GeV	$10 < \tan\beta < 50$	1411.6795	
GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$		2 γ	-	Yes	20.3	$\tilde{\chi}_1^0$	440 GeV	$1 < \tau(\tilde{\chi}_1^0) < 3 \text{ ns}, \text{SPS8 model}$	1409.5542	
$\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow e\bar{e}\nu/\mu\bar{\mu}\nu/\mu\bar{\mu}\nu$		displ. $ee/\mu\mu/\mu\mu$	-	-	20.3	$\tilde{\chi}_1^0$	1.0 TeV	$7 < c\tau(\tilde{\chi}_1^0) < 740 \text{ mm}, m(\tilde{g}) = 1.3 \text{ TeV}$	1504.05162	
RPV		LFV $pp \rightarrow \tilde{\nu}_\tau + X, \tilde{\nu}_\tau \rightarrow e\mu/\tau\mu/\mu\tau$	$e\mu, e\tau, \mu\tau$	-	-	3.2	$\tilde{\nu}_\tau$	1.9 TeV	$\lambda'_{311} = 0.11, \lambda'_{132/133/233} = 0.07$	1607.08079
	Bilinear RPV CMSSM	2 e, μ (SS)	0-3 b	Yes	20.3	\tilde{q}, \tilde{g}	1.45 TeV	$m(\tilde{q}) = m(\tilde{g}), c\tau_{\text{LSP}} < 1 \text{ mm}$	1404.2500	
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow e\bar{e}\nu, \mu\bar{\mu}\nu$	4 e, μ	-	Yes	13.3	$\tilde{\chi}_1^\pm$	1.14 TeV	$m(\tilde{\chi}_1^0) > 400 \text{ GeV}, \lambda_{12k} \neq 0 (k = 1, 2)$	ATLAS-CONF-2016-075	
	$\tilde{\chi}_1^+\tilde{\chi}_1^-, \tilde{\chi}_1^+ \rightarrow W\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \tau\nu\tau, e\nu e, \tau\nu\tau$	3 $e, \mu + \tau$	-	Yes	20.3	$\tilde{\chi}_1^\pm$	450 GeV	$m(\tilde{\chi}_1^0) > 0.2 \times m(\tilde{\chi}_1^\pm), \lambda_{133} \neq 0$	1405.5086	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow q\tilde{q}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow q\tilde{q}\tilde{\chi}_1^0$	0	4-5 large-R jets	-	36.1	\tilde{g}	1.875 TeV	$m(\tilde{\chi}_1^0) = 1075 \text{ GeV}$	SUSY-2016-22	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow t\tilde{t}\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow q\tilde{q}\tilde{\chi}_1^0$	1 e, μ	8-10 jets/0-4 b	-	36.1	\tilde{g}	2.1 TeV	$m(\tilde{\chi}_1^0) = 1 \text{ TeV}, \lambda_{112} \neq 0$	1704.08493	
	$\tilde{g}\tilde{g}, \tilde{g} \rightarrow \tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{s}$	1 e, μ	8-10 jets/0-4 b	-	36.1	\tilde{g}	1.65 TeV	$m(\tilde{t}_1) = 1 \text{ TeV}, \lambda_{323} \neq 0$	1704.08493	
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{s}$	0	2 jets + 2 b	-	36.7	\tilde{t}_1	100-470 GeV, 480-610 GeV		1710.07171	
	$\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\tilde{\ell}$	2 e, μ	2 b	-	36.1	\tilde{t}_1	0.4-1.45 TeV	$\text{BR}(\tilde{t}_1 \rightarrow b\ell/\mu) > 20\%$	1710.05544	
Other	Scalar charm, $\tilde{c} \rightarrow c\tilde{\chi}_1^0$	0	2 c	Yes	20.3	\tilde{c}	510 GeV	$m(\tilde{\chi}_1^0) < 200 \text{ GeV}$	1501.01325	



Lower mass limit at 95% CL [TeV]

0-fermion and 2-fermion operators in “Warsaw basis”

X^3		φ^6 and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
Q_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	Q_φ	$(\varphi^\dagger \varphi)^3$	$Q_{e\varphi}$	$(\varphi^\dagger \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$Q_{\varphi\Box}$	$(\varphi^\dagger \varphi)\Box(\varphi^\dagger \varphi)$	$Q_{u\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p u_r \tilde{\varphi})$
Q_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	$Q_{\varphi D}$	$(\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$	$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^\dagger \varphi G_{\mu\nu}^A G^{A\mu\nu}$	Q_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi l}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^\dagger \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	Q_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\varphi l}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi) (\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^\dagger \varphi W_{\mu\nu}^I W^{I\mu\nu}$	Q_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\varphi e}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^\dagger \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	Q_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^\dagger \varphi B_{\mu\nu} B^{\mu\nu}$	Q_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi) (\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^\dagger \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	Q_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{u}_p \gamma^\mu u_r)$
$Q_{\varphi WB}$	$\varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	Q_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi) (\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W}B}$	$\varphi^\dagger \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	Q_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi) (\bar{u}_p \gamma^\mu d_r)$

Here, $\varphi = H$; $p, r, \dots =$ flavor indices (so $p = r \rightarrow$ flavor universal)

4-fermion operators in “Warsaw basis”


$(\bar{L}L)(\bar{L}L)$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
Q_{ll}	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	Q_{ee}	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	Q_{le}	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{uu}	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{lu}	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qq}^{(3)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{dd}	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	Q_{ld}	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{lq}^{(1)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	Q_{eu}	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	Q_{qe}	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{lq}^{(3)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	Q_{ed}	$(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
		$Q_{ud}^{(8)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qd}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t)$
				$Q_{qd}^{(8)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$
$(\bar{L}R)(\bar{R}L)$ and $(\bar{L}R)(\bar{L}R)$		B -violating			
Q_{ledq}	$(\bar{l}_p^j e_r)(\bar{d}_s^j q_t^j)$	Q_{duq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(d_p^\alpha)^T C u_r^\beta] [(q_s^{\gamma j})^T C l_t^k]$		
$Q_{quqd}^{(1)}$	$(\bar{q}_p^j u_r) \varepsilon_{jk} (\bar{q}_s^k d_t)$	Q_{qqqu}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(q_p^{\alpha j})^T C q_r^{\beta k}] [(u_s^\gamma)^T C e_t]$		
$Q_{quqd}^{(8)}$	$(\bar{q}_p^j T^A u_r) \varepsilon_{jk} (\bar{q}_s^k T^A d_t)$	$Q_{qqq}^{(1)}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} \varepsilon_{mn} [(q_p^{\alpha j})^T C q_r^{\beta k}] [(q_s^{\gamma m})^T C l_t^n]$		
$Q_{lequ}^{(1)}$	$(\bar{l}_p^j e_r) \varepsilon_{jk} (\bar{q}_s^k u_t)$	$Q_{qqq}^{(3)}$	$\varepsilon^{\alpha\beta\gamma} (\tau^I \varepsilon)_{jk} (\tau^I \varepsilon)_{mn} [(q_p^{\alpha j})^T C q_r^{\beta k}] [(q_s^{\gamma m})^T C l_t^n]$		
$Q_{lequ}^{(3)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r) \varepsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$	Q_{duu}	$\varepsilon^{\alpha\beta\gamma} [(d_p^\alpha)^T C u_r^\beta] [(u_s^\gamma)^T C e_t]$		

Implemented in MadGraph UFO models via SMEFTsim, SMEFT@NLO

SMEFT approach is a global approach



$$\sigma_{tot} = \sigma_{SM} + \sum_i \frac{c_i}{\Lambda^2} \sigma_{SM*dim-6} + \sum_{i,j} \frac{c_i c_j}{\Lambda^4} \sigma_{(dim-6)^2}$$





**Operators impact multiple processes:
Global approach needed**

LHC SMEFT analysis goal:


from pattern in deviations, **determine Λ**

SMEFT approach is a global approach




Lots of work in the area!
Tightest constraints on
operators that affect $\bar{f}fV$
couplings or $h \rightarrow \gamma\gamma$

SMEFT approach is a global approach




[2105.00006]



[1910.14012]


Run II, ATLAS+CMS, 68% and 95% C.L.



[1910.03606]

Watch for different assumptions (flavor/CP), coefficient conventions, linear vs. quadratic SMEFT, NLO vs. LO

SMEFT approach is a global approach





Looking for heavy new physics





SMEFT

Looking for heavy new physics

