



# Experimental measurements of Standard Model Properties HCP Summer School 2024

Matthew Herndon

2024-07-29,30

# In this talk, QCD+EW

- **Measurement of SM Properties**

- Introduction to the SM
- Predictions using the SM
- Experimental measurements

- **QCD measurements**

- Jets, PDFs,  $\alpha_s$

- **more QCD**

- W and Z + jets

- **High Precision EW measurement**

- W and Z cross sections
- Parameters:  $\sin^2\theta_{\text{eff}}^l$ ,  $m_W$


- **Multiboson physics**


- di-boson, tri-boson, VBS, polarization, TGC, QGC

- **Summary**

Standard Model Production Cross Section Measurements

Status: June 2024





Nearly every diagram you find like this is wrong!

In this case the  $Z\gamma WW$  vertex not represented

# Measurement of SM Properties

and other introductory material


# The Standard Model of Particle Physics

Exploring the fundamental theory that explains 3 (of 4) of the forces of nature.


This talk will focus on selected measurements QCD and EW physics performed using high transverse momentum ( $p_T$ ) and/or massive SM particles.

Only covers a tiny fraction of the SM measurements made at the LHC (and elsewhere)



See recent reviews from: [ATLAS](#), [CMS](#)



# As measured by the ATLAS and CMS Experiments (+1 from LHCb)



Choice of images is biased – I work on muon systems 😊.



# Some qualitative remarks on the performance of these detectors

The efficiency, angular coverage, granularity, resolution and identification capabilities of the detectors are good enough that most activity from a collision can often be attributed to individual: electrons, muon, taus, photons, charged hadrons or neutral hadrons.

## Electrons and muons

- Identified with low background and near 100% efficiency over large ranges of pseudo-rapidity and momentum
- Often as much as twice the pseudo-rapidity compared to previous collider detectors.
- Enables the precision measurements in single and multi-boson physics and searches for rare processes

## Photons



- Identified efficiently with low background
- Enables the precise cross section measurements in modes with photons and searches for rare processes

## Jets

- Particle flow jets measure and identify most of the jet constituents
- Near hadron level jets
- Excellent jet energy scale and resolution calibrations
- Corrections between reconstructed and hadron level jets fairly diagonal
- Enables an extensive program of comparison with theory, PDF and  $\alpha_s$  determinations, and tuning of MC performance

Missing energy resolution is decent with reduced tails

All this in turn enables many precision measurements:  $m_W$ ,  $\sin^2\theta_{\text{eff}}^l$  ...



The performance of the previous generations of detectors were nothing like this!

and the accelerator and theory communities have accomplished equivalent improvements!

# Predictions - Measurements

## Scientific exploration

- build an experiment → take data → formulate a theory
- theory → make predictions
- build a new experiment → test the predictions


The theory is not really that useful if it's not predictive!

The Standard Model is one of the ultimate predictive theories

We need a strong understanding of the SM and calculations within the SM:

- What is interesting to measure
- How precisely can we predict measurables
- How precisely can/should we measure them
- How to build our detector

Theory/Experiment - pushing each other to higher precision



# Elements of a SM Prediction: the cross section

The basic measurable prediction of the SM at a collider is a production cross section

**Hadron collider: colliding partons: quarks and gluons**

- QCD interactions are ubiquitous
- QCD factorization allows us to separate perturbative and non/semi-perturbative physics and make precisions – introduces factorization scale

**Perturbative calculation hard collision**

- Evaluation of Feynman diagrams -  $\alpha_s, \alpha$
- LO: Almost useless: can be off by factors of 3 ( $gg \rightarrow H$ )
- NLO QCD: large increase in cross sections: 10-100%
  - New initial/final states (involving jets), loop diagrams
  - Renormalization necessary - renormalization scale
  - Minimum necessary for reasonable accuracy
- NNLO QCD: large increase in cross section: 5%-50%
  - Generally, achieves several percent accuracy
  - Necessary for many analyses
- NLO EW: increase (or decrease) the high energy tails of distributions: 5-30%
- Can also incorporating a logarithmic resummation calculation - helps low  $p_T$  or  $p_T$  thresholds of objects, systems,

$$\underbrace{\sigma_{pp \rightarrow \text{jet} + X}}_{\text{experimental data}} = \sum_{ij \in gq\bar{q}} \overbrace{f_i(x_i, \mu_F^2) \otimes f_j(x_j, \mu_F^2)}^{\text{PDFs}} \otimes \underbrace{\hat{\sigma}_{ij \rightarrow \text{jet} + X} \left( x_i, x_j, \frac{Q^2}{\mu_F^2}, \frac{Q^2}{\mu_R^2}, \alpha_S(\mu_R^2) \right)}_{\text{SM(EFT)}}$$

**SM calculation of the inclusive jet cross section**



# Elements of a SM Prediction: the cross section

## Non/semi-perturbative physics

### Colliding partons

- Proton distribution functions, PDFs, structure of the proton
- Non-perturbative at lowest energies
- Evolved up to the perturbative collision scale, factorization scale,  $\mu_F$

### QCD N<sup>X</sup>LO → Jets

- Fragmentation functions and partons showers (PS)
- Hadronization
- Introduces a matching scale, jets produced in the hard collision vs PS jets

### Proton remnants

- Underlying events (UE)
- Multiparton interactions (MPI)


## PS, UE, MPI treated with tunes and/or transverse momentum dependent TMD-PDFs


These issues must be corrected for when measuring a cross section and represent important ancillary measurements

## SM calculation of the inclusive jet cross section

$$\underbrace{\sigma_{pp \rightarrow \text{jet} + X}}_{\text{experimental data}} = \sum_{ij \in gq\bar{q}} \overbrace{f_i(x_i, \mu_F^2) \otimes f_j(x_j, \mu_F^2)}^{\text{PDFs}} \otimes \underbrace{\hat{\sigma}_{ij \rightarrow \text{jet} + X} \left( x_i, x_j, \frac{Q^2}{\mu_F^2}, \frac{Q^2}{\mu_R^2}, \alpha_S(\mu_R^2) \right)}_{\text{SM(EFT)}}$$

# Proceeding down the stairway to discovery!





# QCD Measurements

Jets, PDFs,  $\alpha_s$

CMS xs review


# SM measurement: inclusive jet cross section

## The measurable

- Cross section to produce one jet. Multijet events  $\rightarrow$  multiple contributions per pp collision
- Fiducial:  $p_T(\text{jet})$  and  $y(\text{jet})$  requirements. Typically differential:  $p_T(\text{jet})$  and  $|y(\text{jet})|$  bins

## Defining a jet

- Parton level: original quark or gluon  $p_T$ ,  $y$ 
  - Advantages: What was directly calculated in the cross-section calculation
  - More independent of detector or jet clustering details
  - Solid understanding of perturbative accuracy of the calculation
  - How to consider the jets formed in PS process
  - Disadvantages: Must correct for non/semi-perturbative effects to/from the parton level
- Particle level
  - Cluster jets using MC particles (mostly hadrons)
  - Use infrared/collinear safe jet clustering algorithm.
  - Advantage: closer to what is measured - detector PF objects and/or calorimeter energy clusters
  - Disadvantage: depends on clustering




## Unfolding

- Account for acceptance and migrations in differential measurements – using MC
  - Finite resolutions – events outside of fiducial (or a bin) can migrate into the region and or vice versa
  - Typically iterated. Amount of migration depends on the underlying distribution. Correct the MC to the distribution observed in the data and recalculate migrations.

# Inclusive Jet Cross Section - CMS


JHEP 02 142, JHEP 12 035 (2022)



## Prediction: NNLO QCD with NLO EW corrections

- Corrected for NP effects to particle level jets
- NP corrections based on several MCs and tunes


**Data: Unfolded to particle level using full detector simulation**



**Unfolding Matrix:  
Reconstructed to  
hadron level gen jets**

**Excellent agreement seen with  
NNLO QCD + NLO EW prediction**

# Inclusive Jet Cross Section, NLO vs NNLO



Predictions:

**NNLO QCD with NLO EW corrections**

**NLO+NLL QCD with NLO EW corrections**

**(since NLL was used for PS have to account for that)**

**NLO, even with NLL, is not enough to achieve agreement with the data**

**10-20% disagreements unsurprising.**

**Different predictions from NLO PDFs inconsistent**

**NNLO calculations necessary to demonstrate consistency with SM at the precision of the LHC data for many analyses.**

# The “QCD Analysis”

## Requirements for comparison of SM predictions to LHC data


- NNLO QCD calculations
- NNLO PDF determinations
- NNLO  $\alpha_S$  determination

## PDFs

- LHC probes  $Q^2$  and  $x$  regions of PDFs not accesses by previous accelerators
- Needs HERA data, fix lower  $Q^2$  and  $x$ , where the PDF parameterization is typically defined

## One solution, to assemble these elements: “QCD Analysis”


- Analysis of LHC QCD Jets data (and often other useful data sets)
- Determination PDFs and  $\alpha_S$
- Using HERA I, II datasets with LHC Jets data
- fitting framework available as open-source software - [xFitter](#)



# QCD Analysis, PDFs - ATLAS

## Combined PDF fit of ATLAS data

- Inclusive Jets  $\rightarrow$  strongly impacts valance quark and gluon distributions
- $W, Z/\gamma^*$   $\rightarrow$  strange and anti-quark,  $W$ +Jets,  $Z$ +jets  $\rightarrow$  strange and anti-quark
- $t\bar{t}$   $\rightarrow$  high- $x$  gluon distribution
- Inclusive isolated photon – well fit by data (has often not been the case)




## xFitter framework: ATLASpdf21

- NNLO QCD + NLO EW
- Parameterization at initial  $Q^2$  evolved up to relevant scale using DGLAP equations
- Correlations carefully treated
  - Luminosity, jet related uncertainties

Significant improvement over HERAPDF (where expected). Equivalent performance to Global PDF fits

[Eur. Phys. J. C 82 \(2022\) 438](https://arxiv.org/abs/2107.12788)





# QCD Analysis, PDFs and $\alpha_s$ - CMS

## Combined PDF and $\alpha_s$ fits


- Separate analysis with sensitivity to specific PDFs and/or  $\alpha_s$

Most precise  $\alpha_s$  results from NNLO QCD Analysis of 13 TeV inclusive jet and differential di-Jet data

In general, the ultimate measurements of SM parameters that depend on PDFs will benefit from simultaneous QCD Analysis of the PDFs using appropriately chosen datasets for the parameter(s) of interest –  $m_W$ ,  $\sin\theta_{W\text{eff}}$  ...

CMS


Summary of  $\alpha_s(M_Z)$



CMS xs review

13 TeV Dijets 2D/3D Sub. EPJC 7/29/24





# More QCD Measurements

CMS xs review

## W+jets and Z+jets



# W+jets, Z+jets

## Excellent data-sets for QCD studies

- Easily triggered pure data samples ( $W \rightarrow l\nu$ ,  $Z \rightarrow l^+l^-$ )
- Interesting array of final states
  - Large numbers of jets
  - Heavy flavor jets
  - Topologies important in  $t\bar{t}$ , Higgs, NP searches
- Study of the recoiling jets system via well measured vector boson (Z) properties - tunes

## SM predictions of V+jets





- Interested in exclusive final states (V + njets) or inclusive ( $V + \geq n$  jets)
- Calculation of all diagrams with final states up to V + n jets. If using MC, additional jets from PS

## What does this mean? Some examples

- NLO ZZ+"0" jets production (I work on ZZ physics, so these diagrams were available)
  - LO ZZ prediction poor, neglects large contributions from NLO  $gq$  initial states
  - Inclusive ZZ production NLO QCD accuracy
  - Z+0 at NLO QCD accuracy
  - Renormalized NLO  $\alpha_s$  captures some higher order behavior
  - Neglects contributions from NNLO  $gg$  initial states (add separately)
  - ZZ+1jet production LO QCD accuracy, +  $\geq 2$  jets depends on accuracy of PS (if used) tune
  - Portions of the phase space of a calculation (ZZ+1jet) less accurate than the full inclusive calculation
  - Typically normalize the cross section after PS back to NLO results

LO QDC

Interference between loop and LO diagram is NLO



New initial states

Real emission. Negative contribution from loop diagram cancels infrared divergence in this NLO diagram

# W+jets, Z+jets

## Excellent data-sets for QCD studies

- Easily triggered pure data samples ( $W \rightarrow l\nu$ ,  $Z \rightarrow l^+l^-$ )
- Interesting array of final states
  - Large numbers of jets
  - Heavy flavor jets
  - Topologies important in  $t\bar{t}$ , Higgs, NP searches
- Study of the recoiling jets system via well measured vector boson (Z) properties - tunes

## SM predictions of V+jets





- Interested in exclusive final states (V + njets) or inclusive (V +  $\geq n$  jets)
- Calculation of all diagrams with final states up to V + n jets. If using MC, additional jets from PS

## What does this mean? Some examples

- NLO Z+0,1,2 jets production (three samples generated separately and combined)
  - Inclusive Z production NLO QCD accuracy
  - Z+0,1,2 at NLO QCD accuracy
  - Renormalized NLO  $\alpha_s$  captures some higher order behavior
  - Neglects contributions from higher order initial states
  - Z+3jet production LO QCD accuracy, +  $\geq 4$  jets depends on accuracy of PS (if used) tune
  - Portions of the phase space of a calculation (ZZ+3jet) less accurate than the full inclusive calculation
  - Typically normalize the cross section after PS back to NLO results
  - This is the limit of a reasonable computing time budget

LO QDC

Interference between loop and LO diagram is NLO



New initial states

Real emission. Negative contribution from loop diagram cancels infrared divergence in this NLO diagram

# W+jets, Z+jets

## Excellent data-sets for QCD studies

- Easily triggered pure data samples ( $W \rightarrow l\nu$ ,  $Z \rightarrow l^+l^-$ )
- Interesting array of final states
  - Large numbers of jets
  - Heavy flavor jets
  - Topologies important in  $t\bar{t}$ , Higgs, NP searches
- Study of the recoiling jets system via well measured vector boson (Z) properties - tunes

## SM predictions of V+jets





- Interested in exclusive final states (V + njets) or inclusive (V +  $\geq n$  jets)
- Calculation of all diagrams with final states up to V + n jets. If using MC, additional jets from PS

## What does this mean? Some examples

- NNLO ZZ+0 Jets (three samples generated separately and combined)
  - NNLO overall, Z+0 Jets; NLO Z+1jets; LO Z+2 Jets; +  $\geq 3$  jets depends on accuracy of PS tune if used. Few NNLO + PS implementations available.
  - Should include most important new initial states
  - This is the limit of a reasonable computing time budget

LO QDC

Interference between loop and LO diagram is NLO



New initial states now include a gg diagram

Real emission. Negative contribution from loop diagram cancels infrared divergence in this NLO diagram


# Exclusive Z+jets


## Comparison to LO, NLO and NNLO calculations/MCs

- (if there is a preference for CMS results it's only because I know where to find the plots to make my points – no reflection on the quality of the results)
- LO Magraph 0-4 jets with Pythia 8 PS
  - Why does this look so good? It's up to 4 jets. Most new initial states included.
  - Pythia tunes of the non/semi-perturbative physics heavily leverages Z data
    - UE, MPI and the PS behaviors
  - Note that only statistical uncertainties are shown
  - It's difficult to assess the uncertainty of the MC simulation or interpret the source of differences when observed.
- NLO Magraph 0,1,2 jets with Pythia 8 PS
  - Consistent with data and reasonable uncertainties to up to 2 jets
  - Third hard jet calculated at LO looks good
  - Does not interface well with Pythia 8 PS after that
  - Needs an NLO Tune (see ATLAS AZNLO later!)
- NNLO+NNLL Geneva "0" jets with Pythia 8 PS
  - Better uncertainty in the zero jets bin. Inclusive calculation is great!
  - Consistent with the data with reasonable uncertainties for 1 Jet (NLO) and 2 jets (LO)

To see good agreement and well understood uncertainty you need NLO up to the number of jets of interest

NNLO with jets would be better, but is often prohibitive to generate





# W and Z cross sections

The hadron collider precision cross section frontier

CMS xs review

# Precision pp collider cross sections

## W and Z cross sections



- Easily triggered pure data samples ( $W \rightarrow l\nu$ ,  $Z \rightarrow l^+l^-$ )
- Performance of lepton triggering, reconstruction, and Id calibrated using tag and probe on large data samples.
- Can define both fiducial and total cross-section measurements.
- Dominant uncertainties
  - luminosity uncertainty
  - Extrapolation from measurement phase space to fiducial region (small) or total cross section (larger)
  - Ratios can reduce uncertainties: especially luminosity uncertainty

## The SM calculations

- NNLO, N3LO, N3LO+N3LL, even approximate N4LO+N4LL ([DYTurbo](#))
- Logarithmic resummation of vector boson pT
- Dominant uncertainties
  - PDF and scale uncertainty
  - Especially when phase space limited to fiducial cross-section region

## The precision frontier in hadron collider cross-section measurements and predictions

Interesting dichotomy of uncertainties fiducial vs. total/ exp vs. theory cross section measurement comparisons.






# W and Z cross sections - CMS

5, 13 TeV CMS: to be submitted to JHEP

CMS xs review

## CMS




Inclusive cross section comparison to N3LO QCD with approximate N3LO PDFs

5 energies shown. 13.6 W,Z [PLB 854 \(2024\) 138725 \(ATLAS\)](#) and 13.6 Z ([CMS](#)) results newly available.

# W and Z cross sections historical Context

Takes a few minutes to collect a UA1 sized dataset at the LHC



# W and Z cross sections - CMS

## Inclusive cross sections, N3LO calculation

$\sqrt{s}$ (TeV)	$\sigma(Z)$ (pb)	Tot. exp. unc.	$\sigma^{\text{SM}}(Z)$ (pb)
2.76 [218]	$298 \pm 10$ (stat) (syst) $\pm 11$ (lumi)	5.0%	$313_{-2}^{+1}$
5.02 [219]	$669 \pm 2$ (stat) $\pm 6$ (syst) $\pm 13$ (lumi)	2.2%	$674.7_{-7.4}^{+7.1}$
7 [221]	$986 \pm 22$ (syst) $\pm 22$ (lumi)	3.1%	$968_{-7}^{+6}$
8 [223]	$1138 \pm 26$ (syst) $\pm 30$ (lumi)	3.5%	$1124_{-2}^{+7}$
13 [219]	$1952 \pm 4$ (stat) $\pm 18$ (syst) $\pm 45$ (lumi)	2.5%	$1940_{-21}^{+15}$

## Fiducial cross sections, NNLO calculations

$\sqrt{s}$ (TeV)	$\sigma_{\text{fid.}}(Z)$ (pb)	Tot. exp. unc.	$\sigma_{\text{fid.}}^{\text{SM}}(Z)$ (pb)
5.02 [219]	$319.8 \pm 0.9$ (stat) $\pm 1.2$ (syst) $\pm 6.2$ (lumi)	2.0%	$319.5 \pm 3.7$
7 [221]	$524.7 \pm 0.4$ (stat) $\pm 5.2$ (syst) $\pm 11.5$ (lumi)	2.4%	$525 \pm 6$
8 [222]	$410.0 \pm 10.0$ (stat) $\pm 10.0$ (syst) $\pm 10.0$ (lumi)	4.2%	$400 \pm 10$
13 [219]	$754 \pm 2$ (stat) $\pm 3$ (syst) $\pm 17$ (lumi)	2.3%	$743 \pm 18$

[CMS xs review](#)

## Experimental results

Additional theory uncertainty to extrapolate to total cross-section substantial

2% fiducial Z result at 5TeV

Special short run with reduced instantaneous luminosity

still dominated by luminosity uncertainty

## Theory Calculations

Calculations in a limited phase space less accurate

Exp.-theory pattern of uncertainties best seen in measurements with the same data set

New ATLAS Run 2 luminosity measurement achieves 0.83% uncertainty. Sub 1% cross section measurements possible!



# W and Z cross section ratios - CMS

CMS xs review

## Inclusive cross section ratios, NNLO calculation

$\sqrt{s}$ (TeV)	Ratio	$R_{\text{exp}}$	Tot. exp. unc.	$R_{\text{SM}}$
5.02 [219]	$R_{W^+/W^-}$	$1.519 \pm 0.002$ (stat) $\pm 0.010$ (syst)	0.67%	$1.5240^{+0.33\%}_{-0.31\%}$
7 [220]	$R_{W^+/W^-}$	$1.421 \pm 0.006$ (stat) $\pm 0.032$ (syst)	1.8%	$1.43 \pm 0.7\%$
8 [222]	$R_{W^+/W^-}$	$1.39 \pm 0.01$ (stat) $\pm 0.02$ (syst)	1.6%	$1.41 \pm 0.7\%$
13 [219]	$R_{W^+/W^-}$	$1.3615 \pm 0.0018$ (stat) $\pm 0.0094$ (syst)	0.70%	$1.3536^{+0.37\%}_{-0.33\%}$
5.02 [219]	$R_{W/Z}$	$10.905 \pm 0.032$ (stat) $\pm 0.054$ (syst)	0.58%	$10.777^{+0.33\%}_{-0.34\%}$
7 [220]	$R_{W/Z}$	$10.54 \pm 0.07$ (stat) $\pm 0.18$ (syst)	2.3%	$10.74 \pm 0.4\%$
8 [222]	$R_{W/Z}$	$10.63 \pm 0.11$ (stat) $\pm 0.25$ (syst)	2.6%	$10.74 \pm 0.4\%$
13 [219]	$R_{W/Z}$	$10.491 \pm 0.024$ (stat) $\pm 0.083$ (syst)	0.82%	$10.341^{+0.41\%}_{-0.38\%}$

Experimental measurements

Luminosity uncertainty cancels out. Other uncertainties reduced.

Theory calculation

Scale uncertainties reduced

## Fiducial cross sections ratios, NNLO calculations


$\sqrt{s}$ (TeV)	Ratio	$R_{\text{exp}}$	Tot. exp. unc.	$R_{\text{SM}}$
5.02 [219]	$R_{W^+/W^-}$	$1.6232 \pm 0.0026$ (stat) $\pm 0.0065$ (syst)	0.43%	$1.631 \pm 0.98\%$
8 [222]	$R_{W^+/W^-}$	$1.40 \pm 0.01$ (stat) $\pm 0.02$ (syst)	1.6%	$1.42 \pm 1.4\%$
13 [219]	$R_{W^+/W^-}$	$1.3159 \pm 0.0017$ (stat) $\pm 0.0053$ (syst)	0.43%	$1.307 \pm 1.3\%$
5.02 [219]	$R_{W/Z}$	$12.505 \pm 0.037$ (stat) $\pm 0.032$ (syst)	0.39%	$12.51 \pm 0.96\%$
8 [222]	$R_{W/Z}$	$13.26 \pm 0.15$ (stat) $\pm 0.21$ (syst)	1.9%	$13.49 \pm 2.1\%$
13 [219]	$R_{W/Z}$	$12.078 \pm 0.028$ (stat) $\pm 0.032$ (syst)	0.35%	$12.02 \pm 2.3\%$

Both achieve sub 0.5% uncertainty in best cases!

Not the same best cases.

Fid exp best - Total theory best





CMS xs review



# Measurements of EW Parameters

$m_W, \sin^2\theta_{\text{eff}}^l$

# W mass – ATLAS Improved measurement of $m_W$ and also $\Gamma_W$

- Profile likelihood fit of  $p_T(l)$  and  $m_T(W)$
- Best modeling of W production kinematics and hadronic recoil
- NLO Powheg reweighted with NNLO calculations, NNLO PDFs, Pythia 8 with AZNLO tune based on Z data
- Predicts W  $p_T$  distribution well at 5.02 and 13 TeV, Critical element for extracting  $m_W$  from  $p_T(l)$  and  $m_T(W)$



It would be interesting to see if AZNLO does well with nJet. Focused on  $p_T(Z)$



# W mass - ATLAS

submitted to EPJC

Overview of  $m_W$  measurements



Unc. [MeV]	Total	Stat.	Syst.	PDF	$A_i$	Backg.	EW	$e$	$\mu$	$u_T$	Lumi	$\Gamma_W$	PS
$p_T^e$	16.2	11.1	11.8	4.9	3.5	1.7	5.6	5.9	5.4	0.9	1.1	0.1	1.5
$m_T$	24.4	11.4	21.6	11.7	4.7	4.1	4.9	6.7	6.0	11.4	2.5	0.2	7.0
Combined	15.9	9.8	12.5	5.7	3.7	2.0	5.4	6.0	5.4	2.3	1.3	0.1	2.3

Uncertainties –  $p_T(l)$  measurement dominates  
 Statistical component substantial  
 Systematic distributed over several components  
 Probably difficult, but possible, to match CDF uncertainty

# Effective weak mixing angle – CMS, LHCb

## Drell Yan, $pp \rightarrow l^+l^-$ forward backward asymmetry

- $A_{FB}$  used to determine  $\sin^2\theta_{\text{eff}}^l$

$$\frac{d\sigma}{d\cos\theta} \sim 1 + \cos^2\theta + \frac{1}{2}A_0(1 - 3\cos^2\theta) + A_4\cos\theta$$



- $A_{FB}$  due to  $A_4$  term which depends on  $\sin^2\theta_{\text{eff}}^l$
- Quark direction in the hadron plane inferred from the DY pair rapidity
- Events with forward rapidity leptons more useful for determining  $\sin^2\theta_{\text{eff}}^l$

**CMS includes forward HF electrons  $3.14 < |\eta| < 4.36$**

**LHCb muon reconstruction  $2.0 < |\eta| < 4.5$**



[CMS Preliminary PAS-SMP-22-010](#)

[LHCb preliminary - paper in preparation](#)





# Effective weak mixing angle – CMS, LHCb




Nearing precision of SLD and LEP combination results

Hadron collider results midway between  $e^+e^-$  collider results ☺

Difference in results from various global PDF sets

PDF were profiled but a more global simultaneous QCD

Analysis of PDFs likely beneficial



CMS xs review

# Multi-boson Physics


Di-boson

Weak boson polarization, TGCs

# Di-boson production cross sections

## Di-boson cross sections also near the pp precision frontier

- Same advantages as W and Z physics
- Easily triggered pure data samples (typically at least one  $Z \rightarrow l^+l^-$ )
- Performance of lepton triggering, reconstruction, and Id calibrated using large tag and probe on large data samples.
- Good statistics due to large integrated luminosities
- Dominant uncertainties
  - luminosity uncertainty
  - Extrapolation from measurement phase space to fiducial region (small) or total cross section (larger)



## The SM calculations




- NNLO QCD + NLO EW
- Dominant uncertainties
  - PDF and scale uncertainty
  - Especially when extrapolating to fiducial cross-section

**ZZ,  $Z\gamma$  and WZ measurement achieving precision near that of W and Z cross sections**

# Di-boson production cross sections

## Leading precision from ATLAS $Z\gamma$ , ZZ and CMS WZ measurements

- ZZ: 2.6% precision:  $49.3 \pm 0.8(\text{stat}) \pm 0.8(\text{stat}) \pm 0.8(\text{lumi})$  (1.3 total) fb
- $Z\gamma$ : 2.9% precision:  $533.7 \pm 2.1(\text{stat}) \pm 12.4(\text{stat}) \pm 9.1(\text{lumi})$  fb
- WZ: 3.6% precision:  $298.9 \pm 4.8(\text{stat}) \pm 7.7(\text{syst}) \pm 5.4(\text{lumi}) \pm 2.7(\text{theo})$  fb
- **Luminosity precision the second dominant uncertainty,**  
**ZZ: 2.2,  $Z\gamma$ : 2.5% achievable with new ATLAS luminosity. (my calculation)**




	Full	$Z \rightarrow 4\ell$	Region $H \rightarrow 4\ell$	Off-shell ZZ	On-shell ZZ
Measured	88.9	22.1	4.76	12.4	49.3
fiducial	$\pm 1.1$ (stat.)	$\pm 0.7$ (stat.)	$\pm 0.29$ (stat.)	$\pm 0.5$ (stat.)	$\pm 0.8$ (stat.)
cross-section	$\pm 2.3$ (syst.)	$\pm 1.1$ (syst.)	$\pm 0.18$ (syst.)	$\pm 0.6$ (syst.)	$\pm 0.8$ (syst.)
[fb]	$\pm 1.5$ (lumi.)	$\pm 0.4$ (lumi.)	$\pm 0.08$ (lumi.)	$\pm 0.2$ (lumi.)	$\pm 0.8$ (lumi.)
	$\pm 3.0$ (total)	$\pm 1.3$ (total)	$\pm 0.35$ (total)	$\pm 0.8$ (total)	$\pm 1.3$ (total)
SHERPA	$86 \pm 5$	$23.6 \pm 1.5$	$4.57 \pm 0.21$	$11.5 \pm 0.7$	$46.0 \pm 2.9$
POWHEG + PYTHIA8	$83 \pm 5$	$21.2 \pm 1.3$	$4.38 \pm 0.20$	$10.7 \pm 0.7$	$46.4 \pm 3.0$

**ATLAS should recalculate their Run 2 cross sections with updated luminosities! 😊**


# Di-boson production cross sections

Good precision achieved over a wide array of di-boson final states

Diboson measurements vs. NNLO theory



Diboson Cross Section Measurements



Also now 13.6 TeV WZ and WW results

# Di-boson differential production cross sections

Example CMS: ZZ



Disagreement in high mass tail  $m_{4l}$  distribution

Improved by EW corrections

Disagreement in  $N_{jet}$  distribution

Improved by nNNLO+PS MC. NNLO MC combined with PS using the MiNNLO method

QCD now extensively investigated in di-boson physics also





These disagreements were observed in previous analysis and were improved by advances in theory techniques

# Di-boson production, weak boson polarization

## Vector boson polarization in di-boson production

- An important step toward using polarization to establish longitudinal vector boson scattering
- Longitudinal polarization state is a basic property of the weak bosons from EW symmetry breaking
- Correlations in polarization can test quantum entanglement
- CMS: Observation of individually longitudinally polarized W and Z bosons in WZ production
- ATLAS: Observation longitudinally polarized boson pairs in WZ and evidence in ZZ production ( $4.3\sigma$ )



Single boson polarization extracted from  $\cos(\theta)$



$\theta$  of the lepton relative to boson flight direction in the boson CM frame

Joint polarization required a multivariate discriminant

# Di-boson production, WZ, polarization, RAZ, TGCs

## WZ Radiation Amplitude Zero (RAZ)


- LO WZ production occurs via interfering t channel and triple gauge coupling (TGC) vertex s channel diagrams
- In the transverse-transverse polarization state there is an exact RAZ when W boson is scattered at 90° with respect to the incoming antiquark direction in the WZ rest frame
- This occurs because of an exact cancelation due to interference with the TGC diagram.
- The RAZ is inexact at NLO
- **Most easily observed as a dip near zero in  $\Delta Y(WZ)$**
- This represents one of the few ways in di-boson production to directly observe a TGC and thus a predicted effect of the gauge structure of the SM



**TT signal only – TL, LT, LL subtracted as a background,  $p_T(WZ) < 20 \text{ GeV}$  (more LO like)**

**Approximate RAZ clearly observed!**







# Multi-boson Physics

## Tri-boson

# Tri-boson production

## Tri-boson production many states only accessible at LHC

- Most interesting states, the most difficult to observe, are VVV. Three weak bosons
- Collectively observed by the CMS experiment. WWW observed at ATLAS. WWW, WWZ evidence CMS.
- Extensive analyses pursuing all accessible leptonic and hadronic decay modes





VVV production collectively observed. CMS  
WWW observed ATLAS.

# Tri-boson production

Several states with two weak bosons and all but one state with three yet to be observed.


- CMS  $WW\gamma$ :  $6.0 \pm 0.8$  (stat)  $\pm 0.7$  (syst)  $\pm 0.6$  (modeling) fb,  $5.6\sigma$
- ATLAS  $WZ\gamma$ :  $2.01 \pm 0.30$ (stat)  $\pm 0.16$ (stat) fb,  $6.3\sigma$



CMS :  $WW\gamma$  observed!

ATLAS:  $WZ\gamma$  observed!





# Multi-boson Physics



Vector boson Scattering

Weak boson polarization, QGCs

# Vector boson scattering

## Vector boson scattering (VBS) physics currently unique to the LHC



- Cross sections as small as fb or a fraction of a fb require LHC luminosities to produce observable signals
- VBS diagrams - purely EW interactions.
- Possible to isolate areas of phase space where interference with "QCD" diagrams is minimal
- Production Diagrams include
  - Double TGC in t (and s) channel
  - Quartic gauge coupling (QGC) diagrams: possible to directly measure quartic couplings
  - Higgs scattering: scattering via the Higgs necessary to unitarize the cross section of longitudinal VBS
  - Interfering QCD diagrams
- Final state: forward scattered jets with a large rapidity difference and centrally produces boson pair



First observed  
CMS  $W^\pm W^\pm$   $5.5\sigma$

# Vector boson scattering

With the full Run 2 data set all VBS final states observed






Onward to polarized VBS



# Search for polarized VBS in $W^\pm W^\pm$ - CMS

## Searches now starting for longitudinal VBS

- Use  $W^\pm W^\pm$ , final state used for first observation
- Distinctive same charge final state
- Smaller background of QCD induced  $W^\pm W^\pm$
- Multivariate discriminant needed to maximize sensitivity






Expected (observed) significance for one longitudinal boson  $3.1(2.3)\sigma$   
 Expected  $\sigma_{LLSM} = 0.44$  fb  
 Limit 95% C.L.  $\sigma_{LL} < 1.06$  fb,

# Summary

## CMS and ATLAS have measured a wide array of SM cross section

- Almost every basic QCD+EW final state you expect to produce at the LHC
- The highest precision cross section measurement have reached 2% uncertainty and 1% is likely achievable
- Measurements of fundamental SM parameters are becoming competitive and will soon surpass those of previous experiments
- The complex gauge structure of the SM is being explored with many new measurements
- There are no substantial hints of deviations from the SM in the current set of SM measurements
  - Deviations are often seen in complex final states involving one or many vector bosons and multiple jets
  - These are areas we expect our current calculations to be inadequate
  - However, these calculations and techniques are advancing and resolving observed discrepancies







# ATLAS SM Summary

Status: June 2024

## Standard Model Production Cross Section Measurements



# CMS SM Summary



# CMS SM Summary (QCD+EW)

