


NDFD translation: Paired Training Dataset Update

Colin Weber, Alex Wilkinson

Reco to reco translation

- PRISM analysis currently relies on an interaction model to correct for ND and FD effects.
- These do not completely cancel out between the ND and FD due to detector differences.
- Using the paired data we can perform a data driven *unfolding* using all of the reco information.


Slide from Radi Radev



Reco to reco translation

Slide from Radi Radev


- PRISM analysis currently relies on an interaction model to correct for ND and FD effects.
- These do not completely cancel out between the ND and FD due to detector differences.
- Using the paired data we can perform a data driven *unfolding* using all of the reco information.
- This might be easier to implement in an analysis (and introduce systematics) since we only use reco variables.



Reco to reco translation


Slide from Radi Radev

- PRISM analysis currently relies on an interaction model to correct for ND and FD effects.
- These do not completely cancel out between the ND and FD due to detector differences.
- Using the paired data we can perform a data driven *unfolding* using all of the reco information.
- This might be easier to implement in an analysis (and introduce systematics) since we only use reco variables.




Paired near-to-far dataset

Diagram from Alex Wilkinson




Paired near-to-far dataset

Diagram from Alex Wilkinson




Initial validation showed
discrepancies between ND
simulation, paired data


Unexpected features in paired data



ND-Lar




TMS



No Edep

LAr Edep


Shwr




Other Edep

Likely solution: resimulate muon


Muon contained




ND-Lar



Muon tracker-matched




TMS




Paired near-to-far dataset

Diagram from Alex Wilkinson





Paired near-to-far dataset

Diagram from Alex Wilkinson



Re-validating paired dataset w/ resim. Muon: ND

Effect of Muon Resimulation





Slide from Alex Wilkinson

New Paired Data Validation



2/9

Old simulation





Elep
True

Muon
resimulation





Old simulation





Elep
Reco

Muon
resimulation





Old simulation





Enu
True

Muon
resimulation





Old simulation



Enu
True


Muon
resimulation



Re-validating paired dataset w/ resim. Muon: FD

FD Validation - Nu Energy (Selected)

True energy bins normalised to 1




Slide from Alex Wilkinson

New Paired Data Validation

6/9

FD Validation - Lep Energy (Selected)

True energy bins normalised to 1



Slide from Alex Wilkinson


New Paired Data Validation

7/9

FD Validation - Had Energy (Selected)



True energy bins normalised to 1



Slide from Alex Wilkinson

New Paired Data Validation

8/9

Next steps

- Retrain transformer on new paired dataset
 - Check how robust transformer is to changes in input data
- Repeat for RHC numubar events
- Start exploring paired datasets with oscillations
 - Simulate “tripled” or “paired-paired” datasets with numu and nue